

THE SHI ARRANGEMENT AND THE ISH ARRANGEMENT

DREW ARMSTRONG AND BRENDON RHOADES

ABSTRACT. This paper is about two arrangements of hyperplanes. The first — the **Shi arrangement** — was introduced by Jian-Yi Shi [11, Chapter 7] to describe the Kazhdan-Lusztig cells in the affine Weyl group of type A . The second — the **Ish arrangement** — was recently defined by the first author [1] who used the two arrangements together to give a new interpretation of the q, t -Catalan numbers of Garsia and Haiman. In the present paper we will define a mysterious “combinatorial symmetry” between the two arrangements and show that this symmetry preserves a great deal of information. For example, the Shi and Ish arrangements share the same characteristic polynomial, the same numbers of regions, bounded regions, dominant regions, regions with c “ceilings” and d “degrees of freedom”, etc. Moreover, all of these results hold in the greater generality of “deleted” Shi and Ish arrangements corresponding to an arbitrary subgraph of the complete graph. Our proofs are based on nice combinatorial labelings of Shi and Ish regions and a new set partition-valued statistic on these regions.

1. INTRODUCTION

A **hyperplane arrangement** is a finite collection of affine hyperplanes in Euclidean space. Some of the nicest arrangements come from the reflecting hyperplanes of Coxeter groups. In particular, the **Coxeter arrangement of type A** (also known as the **braid arrangement**) is the arrangement in \mathbb{R}^n defined by


$$\text{Cox}(n) := \{x_i - x_j = 0 : 1 \leq i < j \leq n\}.$$

Here $\{x_1, \dots, x_n\}$ are the standard coordinate functions on \mathbb{R}^n .

Postnikov and Stanley [9] introduced the idea of a **deformation** of the Coxeter arrangement — this is an affine arrangement each of whose hyperplanes is parallel to some hyperplane of the Coxeter arrangement. In the present paper we will study two specific deformations of the Coxeter arrangement and we will observe a deep similarity between them. The first is the **Shi arrangement** which was one of Postnikov and Stanley’s motivating examples:

$$\text{Shi}(n) := \text{Cox}(n) \cup \{x_i - x_j = 1 : 1 \leq i < j \leq n\}.$$

Date: September 10, 2010.

FIGURE 1.1. The arrangements $\text{Shi}(3)$ (left) and $\text{Ish}(3)$ (right)

This arrangement was defined by Jian-Yi Shi [11, Chapter 7] in the study of the Kazhdan-Lusztig cellular structure of the affine Weyl group of type A . The second is the Ish arrangement, recently defined by the first author [1]:

$$\text{Ish}(n) := \text{Cox}(n) \cup \{x_1 - x_j = i : 1 \leq i < j \leq n\}.$$

He used the Shi and Ish arrangements to give a new description of the q, t -Catalan numbers of Garsia and Haiman in terms of the affine Weyl group of type A . Figure 1.1 displays the arrangements $\text{Shi}(3)$ and $\text{Ish}(3)$. (Note that the normals to the hyperplanes of either $\text{Shi}(n)$ or $\text{Ish}(n)$ span the hyperplane $x_1 + x_2 + \cdots + x_n = 0$. Hence we will always draw their restrictions to this space.)

The heart of this paper is the following correspondence between Shi and Ish hyperplanes. The correspondence is natural to state but we find it geometrically mysterious. We will call this a “combinatorial symmetry”:

$$x_i - x_j = 1 \longleftrightarrow x_1 - x_j = i \quad \text{for} \quad 1 \leq i < j \leq n$$

This symmetry allows us to define **deleted** versions of the Shi and Ish arrangements. Let $\binom{[n]}{2}$ denote the set of pairs ij satisfying $1 \leq i < j \leq n$ and consider a simple loopless graph $G \subseteq \binom{[n]}{2}$. The **deleted Shi** and **deleted Ish** arrangement are defined as follows:

$$\text{Shi}(G) := \text{Cox}(n) \cup \{x_i - x_j = 1 : ij \in G\},$$

$$\text{Ish}(G) := \text{Cox}(n) \cup \{x_1 - x_j = i : ij \in G\}.$$

The arrangement $\text{Shi}(G)$ was first considered by Athanasiadis [3]. Note that $\text{Shi}(G)$ (resp. $\text{Ish}(G)$) interpolates between the Coxeter arrangement and the Shi (resp. Ish)




FIGURE 1.2. The arrangements $\text{Shi}(G)$ (left) and $\text{Ish}(G)$ (right) corresponding to the “chain” $G = \{12, 23\} \subseteq \binom{[3]}{2}$

arrangement. That is, if $\emptyset \in \binom{[n]}{2}$ is the “empty” graph and $K_n = \binom{[n]}{2}$ is the “complete” graph, we have

$$\text{Shi}(\emptyset) = \text{Ish}(\emptyset) = \text{Cox}(n), \quad \text{Shi}(K_n) = \text{Shi}(n) \quad \text{and} \quad \text{Ish}(K_n) = \text{Ish}(n).$$

Figure 1.2 displays the arrangements $\text{Shi}(G)$ and $\text{Ish}(G)$ corresponding to the “chain” $G = \{12, 23\} \subseteq \binom{[3]}{2}$.

In order to state our main results right away we need a few definitions.

Let \mathcal{A} be either $\text{Shi}(G)$ or $\text{Ish}(G)$. The connected components of $\mathbb{R}^n - \cup_{H \in \mathcal{A}} H$ are called **regions**. We say that a region is **dominant** if it lies within the **dominant cone**, defined by the coordinate inequalities

$$x_1 > x_2 > \cdots > x_n.$$

The topological closure \bar{R} of a region R is decomposed by the arrangement \mathcal{A} into **faces** of various dimensions. We say that the hyperplane H is a **wall** of R if it is the affine span of a codimension-1 face of R . The wall H is called a **ceiling** if H does not contain the origin and if the region R and the origin lie in the same half-space of H . Since every region R is convex, it determines a **recession cone** (which is closed under non-negative linear combinations):

$$\text{Rec}(R) := \{v \in \mathbb{R}^n : v + R \subseteq R\}.$$

Note that the region R is bounded if and only if $\text{Rec}(R) = 0$. We call the dimension of $\text{Rec}(R)$ the number of **degrees of freedom** of R .

Finally, let $\mathfrak{L}(\mathcal{A})$ denote the collection of intersections of hyperplanes from \mathcal{A} , partially ordered by *reverse*-inclusion of subspaces:

$$\mathfrak{L}(\mathcal{A}) := \{\cap_{H \in S} H : S \subseteq \mathcal{A}\}.$$

This poset has the structure of a geometric semilattice (see [14]) with a unique minimum element \mathbb{R}^n (corresponding to the empty intersection). The **characteristic polynomial** (or **chromatic polynomial**) $\chi_{\mathcal{A}}(p) \in \mathbb{Z}[p]$ of the arrangement \mathcal{A} is defined by

$$\chi_{\mathcal{A}}(p) = \sum_{X \in \mathfrak{L}(\mathcal{A})} \mu(\mathbb{R}^n, X) p^{\dim(X)},$$

where $\mu : \mathfrak{L}(\mathcal{A}) \times \mathfrak{L}(\mathcal{A}) \rightarrow \mathbb{Z}$ is the Möbius function of the poset $\mathfrak{L}(\mathcal{A})$ (see [12]).

Main Theorem. *Let $G \subseteq \binom{[n]}{2}$ be a simple loopless graph on n vertices; let c and d be nonnegative integers. The deleted Shi and Ish arrangements $\mathbf{Shi}(G)$ and $\mathbf{Ish}(G)$ share the following properties in common:*

- (1) *the characteristic polynomial;*
- (2) *the number of **dominant** regions with c ceilings;*
- (3) *the number of regions with c ceilings and d degrees of freedom.*

Proof. Parts (1), (2), (3) are Theorems 3.2, 4.5, and 5.1, respectively. □

For example, here are the joint distributions of ceilings (c) and degrees of freedom (d) for the arrangements in Figures 1.1 and 1.2, respectively.

		d		
		1	2	3
	0			6
c	1	3	6	
	2	1		

		d		
		1	2	3
	0			6
c	1	2	4	
	2	1		

We find it surprising that the symmetry $x_i - x_j = 1 \leftrightarrow x_1 - x_j = i$ preserves so much information. However, there are important properties that it does not preserve. For example, one may observe from Figures 1.1 and 1.2 that the intersection poset is not preserved. One can also show that the Tutte polynomials of $\mathbf{Shi}(3)$ and $\mathbf{Ish}(3)$ differ, and that the Orlik-Solomon algebras of $\mathbf{Shi}(G)$ and $\mathbf{Ish}(G)$ are not graded-isomorphic for $G = \{12, 23\}$ (even though the equality of characteristic polynomials implies that these algebras have the same Hilbert series). Is there a unifying concept that could simplify the statement of the Main Theorem?

The paper is structured as follows.

In **Section 2** we establish some language for set partitions. We define G -partitions — which for the complete graph are just partitions of $[n] = \{1, 2, \dots, n\}$ — and discuss various kinds: connected, nonnesting. We define the **(a, b)** endpoint notation

for partitions which seems to be the correct language for comparing Shi and Ish arrangements.

In **Section 3** we show that $\text{Shi}(G)$ and $\text{Ish}(G)$ have the same characteristic polynomial, which has a formula involving G -Stirling numbers. This proves part (1) of the Main Theorem. Our tool is the finite field method Crapo and Rota [5]. By a standard result of Zaslavsky this implies that $\text{Shi}(G)$ and $\text{Ish}(G)$ share the same numbers of total regions and relatively bounded regions (regions with one degree of freedom).

In **Section 4** we modify a labeling of the regions of $\text{Shi}(G)$ due to Athanasiadis and Linusson [2] and we call the result **Shi ceiling diagrams**. Similarly, we define **Ish ceiling diagrams** for the regions of $\text{Ish}(G)$. We give a bijective proof of part (2) of the Main Theorem by observing that **dominant** regions of $\text{Shi}(G)$ and $\text{Ish}(G)$ correspond to order ideals in isomorphic posets.

In **Section 5** we define the **ceiling partition** for a region of $\text{Shi}(G)$ or $\text{Ish}(G)$. This is a (possibly nesting) G -partition that encodes the ceilings of the region. Given a G -partition π with k blocks and an integer $1 \leq d \leq k$, we show that the number of regions of either $\text{Shi}(G)$ or $\text{Ish}(G)$ with ceiling partition π and d degrees of freedom is equal to

$$\frac{d(n-d-1)!(k-1)!}{(n-k-1)!(k-d)!},$$

which proves part (3) of the Main Theorem. This formula is remarkable, and it is new even for the Shi arrangement. The proof of the formula for Ish regions is direct, whereas the proof for Shi regions uses a new formula due to the second author (see [10] or Lemma 2.3) which counts nonnesting partitions with a fixed number of connected components and fixed block size multiplicities. This suggests an open problem: Find a bijection between Shi regions and Ish regions with ceiling partition π and d degrees of freedom. This bijection **cannot** preserve the property of being dominant, since $\text{Shi}(G)$ and $\text{Ish}(G)$ do **not** share the same number of dominant regions with d degrees of freedom.

We end with an observation:

The Ish arrangement is something of a “toy model” for the Shi arrangement (and other Catalan objects). That is, for any property P that $\text{Shi}(G)$ and $\text{Ish}(G)$ share, the proof that $\text{Ish}(G)$ satisfies P is easier than the proof that $\text{Shi}(G)$ satisfies P .

2. SET PARTITIONS

All of the formulas in this paper are phrased in terms of set partitions. In this section we will give some background on these and establish notation. In particular, for each graph $G \subseteq \binom{[n]}{2}$ we will define G -**partitions** of the set $[n] = \{1, 2, \dots, n\}$. In the case of the complete graph this corresponds to the usual notion of partitions.




FIGURE 2.1. A partition of $[8]$ with type $(1, 0, 1, 1, 0, 0, 0, 0)$

2.1. The endpoint notation. We say that $\pi = \{B_1, B_2, \dots, B_k\}$ is a partition of $[n]$ into k blocks if the following disjoint union holds:

$$[n] = B_1 \sqcup B_2 \sqcup \dots \sqcup B_k.$$

The **type** of the partition π is the sequence (r_1, r_2, \dots, r_n) where r_i is the number of blocks of π with size i . We draw the **arc diagram** of π as follows: Place the numbers $1, 2, \dots, n$ on a line and draw an arc between each pair $i < j$ such that

- i and j are in the same block of π ; and
- there is no $i < \ell < j$ such that i, ℓ, j are in the same block of π .

Figure 2.1 displays the arc diagram for the partition $\{\{1, 2, 5, 6\}, \{3, 7, 8\}, \{4\}\}$, which has type $(1, 0, 1, 1, 0, 0, 0, 0)$.

In this paper we will use a special notation for partitions, based on the arc diagram. First note that a partition π has $n - k$ blocks if and only if its diagram has k arcs. This is because each new arc reduces the number of blocks by one. Now suppose that the arcs of π are $a_1b_1, a_2b_2, \dots, a_kb_k$, with the left endpoints in increasing order: $a_1 < a_2 < \dots < a_k$. We will associate π with its pair (\mathbf{a}, \mathbf{b}) of **endpoint vectors**:

$$\mathbf{a} = a_1a_2 \dots a_k \quad \text{and} \quad \mathbf{b} = b_1b_2 \dots b_k.$$

We call (\mathbf{a}, \mathbf{b}) the **endpoint notation** for π . For example, the endpoint notation for the partition in Figure 2.1 is $(12357, 25768)$. It is straightforward to check that partitions of $[n]$ are in bijection with pairs of vectors (\mathbf{a}, \mathbf{b}) such that

- \mathbf{a} and \mathbf{b} have the same length (called the **length** of the pair (\mathbf{a}, \mathbf{b})),
- $a_i < b_i$ for all i ,
- the entries of \mathbf{a} are increasing, and
- the entries of \mathbf{b} are distinct.

In particular, the empty pair (\emptyset, \emptyset) corresponds to the partition $\{\{1\}, \{2\}, \dots, \{n\}\}$ and the longest pair $(12 \dots (n-1), 23 \dots n)$ corresponds to the partition $\{[n]\}$. We will see that the endpoint notation is the best language for comparing Shi and Ish arrangements.

2.2. Nonnesting partitions. A partition π of $[n]$ is called **nonnesting** if it does not contain arcs ij and $k\ell$ such that $i < k < \ell < j$ — that is, no arc of π “nests” inside another. The partition in Figure 2.1 is *not* nonnesting (it is **nesting**) because the arc

56 nests inside the arc 37. The number of nonnesting partitions of $[n]$ is famously given by the Catalan number $\frac{1}{n+1}\binom{2n}{n}$.

The property of nonnesting agrees well with the endpoint notation for partitions. That is, a partition (\mathbf{a}, \mathbf{b}) is nonnesting if and only if its right endpoint vector \mathbf{b} is *increasing*. In fact, the number of pairs of nesting arcs in (\mathbf{a}, \mathbf{b}) is equal to the number of pairs $b_i > b_j$ such that $i < j$.

2.3. G -partitions and G -Stirling numbers. Now we define a version of set partitions for any graph $G \subseteq \binom{[n]}{2}$:

We say that a partition π of $[n]$ is a G -partition if all of its arcs are contained in the graph G . The G -Stirling number $\text{Stir}(G, k)$ is the number of G -partitions with k blocks.

In particular, when G is the complete graph $K_n = \binom{[n]}{2}$ the G -partitions are unrestricted partitions of $[n]$ and the G -Stirling numbers are the classical Stirling numbers (of the second kind).

2.4. Connectivity. Finally, we mention an auxiliary (nontrivial) result which we need for the proof of the Main Theorem. For $i \leq j$ we say that a partition π of the set $\{i, i+1, \dots, j\}$ is **connected** if there does not exist $i \leq k < j$ such that π refines the partition

$$\{\{i, i+1, \dots, k\}, \{k+1, \dots, j-1, j\}\}.$$

Equivalently, π is connected if its arc diagram has no holes when seen from space. The partition in Figure 2.1 *is* connected. Moreover, a partition π of $[n]$ has d **connected components** if there exist numbers $1 < i_1 < \dots < i_{d-1} < n$ such that π refines the partition

$$\{\{1, 2, \dots, i_1 - 1\}, \{i_1, i_1 + 1, \dots, i_2 - 1\}, \dots, \{i_{d-1}, i_{d-1} + 1, \dots, n\}\}$$

and if its restriction to each block of this partition is connected. Equivalently, the arc diagram of a partition with d connected components has $d - 1$ holes when seen from space. For example, the partition $\{\{1, 2\}, \{3, 5, 7\}, \{4, 6\}, \{8\}\}$ has 3 connected components.

The second author has recently established an enumerative formula for nonnesting partitions (and other Catalan objects) that takes account of the type of the partition and its number of connected components. The prototype for this formula is the following theorem of Kreweras [7, Theorem 4]. Kreweras stated his formula in terms of noncrossing partitions; however, type-preserving bijections between noncrossing and nonnesting partitions have been observed by several authors.

Lemma 2.1. *Let $n > 0$ and suppose that the sequence (r_1, \dots, r_n) of nonnegative integers satisfies $\sum_i ir_i = n$ and $\sum_i r_i = k$. The number of nonnesting partitions of $[n]$ with type (r_1, \dots, r_n) is*

$$\frac{n!}{(n - k + 1)!r_1!r_2! \dots r_n!}.$$

We will need the following formula of the second author [10] in our proof of part (3) of the Main Theorem. The proof of this result is combinatorial and relies on the enumeration of words in certain monoids.

Lemma 2.2. [10, Theorem 2.3, Part 2] *Let $n > 0$ and suppose that the sequence (r_1, \dots, r_n) of nonnegative integers satisfies $\sum_i ir_i = n$ and $\sum_i r_i = k$. Let $k \geq d$ and assume that $(r_1, \dots, r_n) \neq (n, 0, \dots, 0)$. The number of nonnesting partitions of $[n]$ with type (r_1, \dots, r_n) and d connected components is*

$$\frac{d(n - d - 1)!(k - 1)!}{(n - k - 1)!(k - d)!r_1!r_2! \dots r_n!}.$$

When $k < d$, it is clear that there are no partitions of $[n]$ of type (r_1, \dots, r_n) with $\sum_i r_i = k$ and d connected components; there is a unique (nonnesting) partition of $[n]$ with type $(n, 0, \dots, 0)$ and it has n connected components.

We remark that the product formula in Lemma 2.3 was predicted from the formula (5.1) for Ish arrangements, **not** by studying nonnesting partitions directly. This is one case in which the Ish arrangement acted as a “toy model” for other Catalan objects.

3. CHARACTERISTIC POLYNOMIALS

In this section we explicitly compute the characteristic polynomials of $\text{Shi}(G)$ and $\text{lsh}(G)$ and observe that they are equal. The formula is expressed in terms of G -Stirling numbers $\text{Stir}(G, k)$. Our tools are the finite field method of Crapo and Rota and the principle of inclusion-exclusion. Zaslavsky’s theorem then implies that $\text{Shi}(G)$ and $\text{lsh}(G)$ have the same number of regions and the same number of relatively bounded regions (regions with one degree of freedom).

3.1. The method. Let \mathcal{A} be a finite hyperplane arrangement in \mathbb{R}^n and suppose that the defining equations for hyperplanes in \mathcal{A} have coefficients in \mathbb{Z} . Then the finite field method of Crapo and Rota [5] is a useful way to compute the characteristic polynomial of \mathcal{A} without having to know its intersection poset. Let $p \in \mathbb{Z}$ be prime and consider a hyperplane $H \subseteq \mathbb{R}^n$ with fixed defining equation $a_1x_1 + \dots + a_nx_n = b$,

where $a_i, b \in \mathbb{Z}$. Then we define the following subset H_p of the *finite* vector space \mathbb{F}_p^n by reducing the coefficients of H modulo p :

$$H_p := \{(x_1, x_2, \dots, x_n) \in \mathbb{F}_p^n : a_1x_1 + \dots + a_nx_n = b\}.$$

Observe that H_p may not be a hyperplane in \mathbb{F}_p^n when p is small, and that H_p in general depends on the defining equation chosen. However: If p is large enough then each H_p is a hyperplane in \mathbb{F}_p^n and the characteristic polynomial of \mathcal{A} has a nice relationship to the **reduced hyperplane arrangement** $\mathcal{A}_p := \{H_p : H \in \mathcal{A}\}$ in \mathbb{F}_p^n .

Theorem 3.1. [5] *Let $p \in \mathbb{Z}$ be a large prime, and let \mathcal{A} be a finite collection of hyperplanes in \mathbb{R}^n whose hyperplanes have defining equations with coefficients in \mathbb{Z} . Then the characteristic polynomial of \mathcal{A} satisfies*

$$\chi_{\mathcal{A}}(p) = \#(\mathbb{F}_p^n - \cup_{H \in \mathcal{A}} H_p).$$

That is, $\chi_{\mathcal{A}}(p)$ counts the number of points in the complement of the reduced arrangement \mathcal{A}_p in the finite vector space \mathbb{F}_p^n .

3.2. The calculation. Now we use the finite field method to compute the characteristic polynomials of $\text{Shi}(G)$ and $\text{Ish}(G)$. We observe that they are equal.

Theorem 3.2. *Let $G \subseteq \binom{[n]}{2}$ be a graph on n vertices. The characteristic polynomials of the deleted Shi and Ish arrangements are given by:*

$$\chi_{\text{Shi}(G)}(p) = \chi_{\text{Ish}(G)}(p) = p \sum_{k=0}^{n-1} (-1)^k \text{Stir}(G, n-k) \frac{(p-k-1)!}{(p-n)!}.$$

Proof. Let $p \in \mathbb{Z}$ be a large prime. We will show that the reduced complements $\mathbb{F}_p^n - \text{Shi}(G)_p$ and $\mathbb{F}_p^n - \text{Ish}(G)_p$ (forgive the abuse of notation) contain the same number of points, counted by the above formula.

To do this, we identify $\{0, 1, \dots, p-1\} = \mathbb{F}_p$ with the vertices of a regular p -gon, ordered clockwise. (That is, $i+1$ is just clockwise of i .) Then a vector $v = (v_1, \dots, v_n) \in \mathbb{F}_p^n$ is a labeling of the vertices: if $v_i = j$ then we place the label v_i on the vertex j . Note that $v \in \mathbb{F}_p^n$ is in the complement of the (reduced) Coxeter arrangement $\text{Cox}(n)_p$ precisely when $v_i - v_j \neq 0$ for all $1 \leq i < j \leq n$. That is, the points of $\mathbb{F}_p^n - \text{Cox}(n)_p$ correspond to *injective* labelings $\{v_1, \dots, v_n\} \hookrightarrow \mathbb{F}_p$. The complements of $\text{Shi}(G)_p$ and $\text{Ish}(G)_p$ are both contained in $\mathbb{F}_p^n - \text{Cox}(n)_p$, so we must count certain kinds of injective labelings.

First we deal with $\text{Shi}(G)_p$. For any set of edges $S \subseteq G$ let $f(S)$ denote the number of vectors $v \in \mathbb{F}_p^n - \text{Shi}(G)_p$ such that $v_i - v_j = 1$ for all edges $ij \in S$ (this notation implies $i < j$). By the principle of inclusion-exclusion (see for example [12, Chapter

2]) we observe that the number of points in $\mathbb{F}_p^n - \text{Shi}(G)_p$ is equal to

$$(3.1) \quad \sum_{S \subseteq G} (-1)^{|S|} f(S).$$

Now suppose that S contains edges ij and $i\ell$ with the same left endpoint. The conditions $v_i - v_j = 1$ and $v_i - v_\ell = 1$ imply that $v_j = v_\ell$ which cannot be satisfied on $\mathbb{F}_p^n - \text{Cox}(n)_p$, hence $f(S) = 0$. Similarly $f(S) = 0$ whenever S contains two edges with the same right endpoint. That is, the sets S that contribute to the sum (3.1) are precisely the arc sets of G -partitions.

Let $S \subseteq G$ correspond to a G -partition with $n - k$ blocks (that is, $|S| = k$). To compute $f(S)$ note that the conditions $v_i - v_j = 1$ for all $ij \in S$ imply that the p -gon \mathbb{F}_p gets labeled by $n - k - 1$ (given) contiguous strings of labels with spaces between. There are $(n - k - 1)!$ ways to cyclically permute the strings; there are $\binom{p-k-1}{n-k-1}$ ways to place $p - n$ empty spaces between the strings; and there are p ways to choose the “origin” (the location of 0). Hence:

$$(3.2) \quad f(S) = \frac{p(p - k - 1)!}{(p - n)!}.$$

Combining (3.1) and (3.2) with the finite field method gives the desired formula.

We use a parallel argument to deal with $\text{lsh}(G)_p$. For any set of edges $S \subseteq G$ let $g(S)$ be the number of vectors $v \in \mathbb{F}_p^n - \text{lsh}(G)_p$ such that $v_1 - v_j = i$ for all $ij \in S$. As above, the points of $\mathbb{F}_p^n - \text{lsh}(G)_p$ are counted by

$$(3.3) \quad \sum_{S \subseteq G} (-1)^{|S|} g(S),$$

and one can check that $g(S) = 0$ unless S is the arc set of a G -partition. We let S correspond to a G -partition with $n - k$ blocks (that is, $|S| = k$) and compute $g(S)$ as follows. First choose v_1 in p ways. Then for each $ij \in S$ the condition $v_1 - v_j = i$ uniquely determines the value of v_j . The remaining $n - k - 1$ labels must be placed injectively in the remaining $p - k - 1$ positions and there are $(p - k - 1)(p - k - 2) \cdots (p - n + 1)$ ways to do this. Thus we get the desired formula:

$$(3.4) \quad g(S) = \frac{p(p - k - 1)!}{(p - n)!}.$$

□

Notice that the counting argument for computing $\chi_{\text{lsh}(G)}(p)$ was more straightforward than the argument for $\chi_{\text{Shi}(G)}(p)$. This again agrees with our observation that lsh is a toy model for Shi . It is somewhat surprising that the two inclusion-exclusion arguments result in the same expression. It may be interesting to find a direct bijection between the points of the complements $\mathbb{F}_p^n - \text{Shi}(G)_p$ and $\mathbb{F}_p^n - \text{lsh}(G)_p$.

3.3. Remarks. A simplified version of the above argument shows that the characteristic polynomials of $\text{Shi}(n)$ and $\text{Ish}(n)$ (the case of the complete graph) are both equal to $p(p-n)^{n-1}$. This result was obtained earlier by Headley [6] and Athanasiadis [3] (for the Shi arrangement) and by the first author [1, Theorem 1] (for the Ish arrangement). Moreover, Athanasiadis described a special family of graphs G for which the characteristic polynomial of $\text{Shi}(G)$ splits. His result [4, Theorem 2.2] together with Theorem 3.2 implies the following.

Corollary 3.3. *Suppose the graph $G \subseteq \binom{[n]}{2}$ has the following property: if $i < j < k$ and $ij \in G$, then $ik \in G$. Then we have*

$$\chi_{\text{Shi}(G)}(p) = \chi_{\text{Ish}(G)}(p) = p \prod_{i=1}^{n-1} (p - d_i - i),$$

where $d_i := \#\{j : ij \in G\}$ is the outdegree of vertex i in G .

In the same paper, Athanasiadis showed that the arrangements $\text{Shi}(G)$ of the Corollary are **free** in the sense of Terao [13] (see also [8]). This is an open problem for the corresponding Ish arrangements $\text{Ish}(G)$.

We remark that the characteristic polynomial of an arrangement allows us to count certain kinds of regions. Some notation: Let \mathcal{A} be a finite collection of hyperplanes in \mathbb{R}^n and suppose that the normals to the hyperplanes span a space $V \subseteq \mathbb{R}^n$ of dimension r . This r is called the **rank** of the arrangement. If $r < n$ then the arrangement \mathcal{A} has no bounded regions; in this case we say that a region of \mathcal{A} is **relatively bounded** if its intersection with V is bounded. The following is a classic theorem of Zaslavsky.

Theorem 3.4. [15] *Let \mathcal{A} be a hyperplane arrangement in \mathbb{R}^n with rank r . Then:*

- *The number of regions of \mathcal{A} is $(-1)^n \chi_{\mathcal{A}}(-1)$;*
- *The number of relatively bounded regions of \mathcal{A} is $(-1)^r \chi_{\mathcal{A}}(1)$.*

Corollary 3.5. *The arrangements $\text{Shi}(G)$ and $\text{Ish}(G)$ have the same number of regions and the same number of relatively bounded regions.*

Observe that the normals to either $\text{Shi}(G)$ or $\text{Ish}(G)$ span the hyperplane $x_1 + x_2 + \cdots + x_n = 0$. Hence each of these arrangements has rank $n - 1$. It follows that neither arrangement has bounded regions and its relatively bounded regions have one degree of freedom. In the case of the complete graph, we find that the arrangements $\text{Shi}(n)$ and $\text{Ish}(n)$ both have $(n + 1)^{n-1}$ regions and $(n - 1)^{n-1}$ regions with one degree of freedom.

The fact that the Shi arrangement $\text{Shi}(n)$ has $(n + 1)^{n-1}$ regions was first proved by Jian-Yi Shi (see [11]). This beautiful result has motivated more than a few research papers since 1985 (including the present one).

4. LABELING THE REGIONS

Fix a graph $G \subseteq \binom{[n]}{2}$. In this section we devise combinatorial labels for the regions of the deleted arrangements $\text{Shi}(G)$ and $\text{Ish}(G)$; we call these labels **Shi ceiling diagrams** and **Ish ceiling diagrams**, respectively. (Something like “Shi floor diagrams” appeared earlier in Athanasiadis and Linusson [2].) Essentially, each diagram encodes the ceilings of a given region, from which we can easily determine its recession cone.

4.1. Shi ceiling diagrams. Recall that the regions (cones) of the Coxeter arrangement $\text{Cox}(n)$ correspond to elements of the symmetric group $\mathfrak{S}(n)$. If $C \subseteq \mathbb{R}^n$ is the **dominant cone** — defined by the coordinate inequalities $x_1 > x_2 > \cdots > x_n$ — then the collection of regions of $\text{Cox}(n)$ is $\{wC : w \in \mathfrak{S}(n)\}$, where wC is defined by the coordinate inequalities

$$(4.1) \quad x_{w(1)} > x_{w(2)} > \cdots > x_{w(n)}.$$

Now let R be a region of the deleted Shi arrangement $\text{Shi}(G)$. Since $\text{Cox}(n) \subseteq \text{Shi}(G)$, R is contained in some cone wC . In this case, what are the possible ceilings of R ? We note that the hyperplanes of $\text{Shi}(G)$ that intersect wC are precisely

$$\Phi^+(G, w) := \{x_{w(i)} - x_{w(j)} = 1 : ij \in G \text{ and } w(i) < w(j)\}.$$

(We can think of these as the *non-inversions* of w contained in G .) Furthermore, suppose that the region R is “below” some hyperplane $x_{w(i)} - x_{w(j)} = 1$ — that is, suppose that each $v \in R$ satisfies $v_{w(i)} - v_{w(j)} < 1$. Then, considering (4.1), R is also below any hyperplane of the form $x_{w(i')} - x_{w(j')} = 1$ such that

$$(4.2) \quad w(i) \leq w(i') < w(j') \leq w(j).$$

That is, if we declare a partial order on $\Phi^+(G, w)$ by saying that $x_{w(i')} - x_{w(j')} = 1$ is “less than” $x_{w(i)} - x_{w(j)} = 1$ when condition (4.2) holds, then the collection of $\text{Shi}(G)$ -hyperplanes above R forms a down-closed set.

Theorem 4.1. *There is a bijection between regions of $\text{Shi}(G)$ in the cone wC and order ideals (down-closed sets) in the poset $\Phi^+(G, w)$. This map sends a region R to the set of hyperplanes in $\text{Shi}(G)$ that are “above” R (contain R and the origin in the same half space). The maximal elements of the ideal are the ceilings of R .*

Proof. Let R be a region of $\text{Shi}(G)$ contained in wC . We showed above that the collection of hyperplanes above R is an order ideal in $\Phi^+(G, w)$. The map is injective since these hyperplanes uniquely determine R . Observe that the ceilings of R are the elements of the ideal that may be individually removed to obtain another ideal, and these are precisely the maximal elements. We refer to Athanasiadis and Linusson [2] for the proof that every ideal corresponds to a non-empty region. \square




FIGURE 4.1. An order ideal (left) and a Shi ceiling diagram (right)


To express this combinatorially, **we note that order ideals in $\Phi^+(G, w)$ are equivalent to nonnesting G -partitions whose blocks are “increasing” with respect to w .** Indeed, there is a bijection between ideals and antichains (sets of pairwise-incomparable elements), since an ideal is uniquely determined by its antichain of maximal elements. By sending the hyperplane $x_{w(i)} - x_{w(j)} = 1$ to the arc ij , each antichain in $\Phi^+(G, w)$ corresponds to a G -partition of $[n]$ whose arcs $i < j$ satisfy $w(i) < w(j)$. Finally, note that two arcs nest if and only if they are comparable in the poset $\Phi^+(G, w)$.

Following these remarks, we draw a diagram for each region of $\text{Shi}(G)$.

Definition 4.1. Let R be a region of $\text{Shi}(G)$ contained in the cone wC . We associate R with the pair (w, π) where π is an order ideal in the poset $\Phi^+(G, w)$ of non-inversions of w contained in G . Equivalently, π is a nonnesting G -partition whose blocks are increasing with respect to w . We draw (w, π) by placing the arc diagram for π above the numbers $w(1), \dots, w(n)$, and we call this the **Shi ceiling diagram** of R .

For example, let $K_8 = \binom{[8]}{2}$ be the complete graph on 8 vertices and consider the permutation $w = 51286347 \in \mathfrak{S}(8)$. Figure 4.1 displays the ideal in $\Phi^+(K_8, w)$ (left) and the ceiling diagram (right) corresponding to a region R of $\text{Shi}(K_8) = \text{Shi}(8)$ contained in the cone wC . The squares are elements of the poset $\Phi^+(K_8, w)$ and the circles are elements of the ideal (closed to the right and down). The hollow circles (maximal elements) indicate the ceilings of the region: $x_5 - x_8 = 1$, $x_1 - x_6 = 1$, $x_3 - x_7 = 1$. The corresponding nonnesting partition is $\pi = \{\{1, 4\}, \{2, 5\}, \{3\}, \{6, 8\}, \{7\}\}$.

Figure 4.2 displays the whole arrangement $\text{Shi}(3)$ with its regions labeled by ceiling diagrams. Observe that we can read the degrees of freedom from the ceiling diagram (w, π) : the corresponding region has d degrees of freedom if and only if the nonnesting partition π has d connected components. This is a general phenomenon.

FIGURE 4.2. The Shi arrangement $\text{Shi}(3)$ labeled by ceiling diagrams

Lemma 4.2. *Let R be a region of $\text{Shi}(G)$ with ceiling diagram (w, π) . This region has d degrees of freedom if and only if the nonnesting partition π of $[n]$ has d connected components.*

Proof. Suppose that π has d connected components. That is, there exist $1 < i_1 < \dots < i_{d-1} < n$ such that π refines the partition

$$\{\{1, 2, \dots, i_1 - 1\}, \{i_1, i_1 + 1, \dots, i_2 - 1\}, \dots, \{i_{d-1}, i_{d-1} + 1, \dots, n\}\},$$

and its restriction to any block of this partition is connected. We compute the recession cone of $\text{Rec}(R) \subseteq \mathbb{R}^n$ of R as follows.

Consider $v = (v_1, \dots, v_n) \in \text{Rec}(R)$. Since R is in the cone wC we must have $v_{w(1)} \geq v_{w(2)} \geq \dots \geq v_{w(n)}$. Moreover, if $ij \in G$ with i and j in the same block of π then the coordinate inequality $x_{w(i)} - x_{w(j)} < 1$ holds on R and we must have $v_{w(i)} = v_{w(i+1)} = \dots = v_{w(j)}$. Since these are the only constraints on v , we conclude that the recession cone $\text{Rec}(R)$ consists of all vectors of the form $w \cdot (a_1, a_2, \dots, a_n)$, where $a_1 \geq \dots \geq a_n$ and where $a_i = a_j$ if i and j are in the same connected component of π . The dimension of the cone is therefore d . \square

For example, consider the ceiling diagram (w, π) in Figure 4.1 and the corresponding region R of $\text{Shi}(8)$. The connected components of π are $\{1, 2, 3, 4, 5\}, \{6, 7, 8\}$ and their images under w are $\{5, 1, 2, 8, 5\}, \{3, 4, 7\}$. Hence the recession cone $\text{Rec}(R)$ consists of all vectors of the form $(a, a, b, b, a, a, b, a) \in \mathbb{R}^8$ with $a \geq b$, and it has dimension 2.

4.2. Ish ceiling diagrams. In order to compare the two arrangements, we now define an Ish analogue of Shi ceiling diagrams.

Since $\text{Cox}(n) \subseteq \text{Ish}(G)$, each region R of $\text{Ish}(G)$ is contained in wC for some permutation $w \in \mathfrak{S}(n)$, in which case each vector $v = (v_1, \dots, v_n) \in R$ satisfies

$$(4.3) \quad v_{w(1)} > v_{w(2)} > \dots > v_{w(n)}.$$

Which $\text{Ish}(G)$ -hyperplanes are the possible ceilings of this region? If the hyperplane $x_1 - x_j = i$ intersects the cone wC it must be true that $x_1 > x_j$ on wC (since i is positive). Considering (4.3), this means that j must occur to the right of 1 in the list $w(1), \dots, w(n)$ — that is, we must have $w^{-1}(1) < w^{-1}(j)$. We conclude that the Ish hyperplanes that intersect the cone wC are precisely

$$\Psi^+(G, w) := \{x_1 - x_j = i : ij \in G \text{ and } w^{-1}(1) < w^{-1}(j)\}.$$

Now let R be a region of $\text{Ish}(G)$ in the cone wC and suppose that R is below $x_1 - x_j = i$ — that is, each $v \in R$ satisfies $v_1 - v_j < i$. Then it is easy to check that R is also below the hyperplane $x_1 - x_{j'} = i'$, where

$$(4.4) \quad \text{either } i < i' \text{ or } w^{-1}(j') < w^{-1}(j).$$

By analogy with the Shi case, we define a partial order on $\Psi^+(G, w)$ by declaring that the hyperplane $x_1 - x_j = i$ is “less than” the hyperplane $x_1 - x_{j'} = i'$ whenever (4.4) holds. This leads to a useful characterization of $\text{Ish}(G)$ regions.

Theorem 4.3. *There is a bijection between regions of $\text{Ish}(G)$ in the cone wC and order filters (up-closed sets) in the poset $\Psi^+(G, w)$. This map sends a region R to the set of hyperplanes in $\text{Ish}(G)$ that are “above” R (contain R and the origin in the same half space). The minimal elements of the filter are the ceilings of R .*

Proof. Let R be a region of $\text{Ish}(G)$ in the cone wC . By the above remarks we know that the collection of $\text{Ish}(G)$ -hyperplanes above R is an order filter in $\Psi^+(G, w)$. These hyperplanes together with the fact that R lies in wC uniquely determine R , so the map is injective. The ceilings of R are precisely the elements of this filter that may be individually removed to obtain another filter — that is, they are the minimal elements.

To show that the map is surjective, we must show that each filter in $\Psi^+(G, w)$ corresponds to a non-empty region of $\text{Ish}(G)$. Let $F \subseteq \Psi^+(G, w)$ be an order filter and let $A \subseteq F$ be its set of minimal elements. For $1 \leq i \leq n$ define

$$z_i = -\max\{j : x_1 - x_k = j \in A \text{ and } w^{-1}(k) \leq w^{-1}(i)\},$$

where we adopt the convention that $\max(\emptyset) = 0$. One may check that the point $(z_{w(1)}, \dots, z_{w(n)}) \in \mathbb{R}^n$ lies on the boundary of a region of $\text{Ish}(G)$ which maps to the filter F . (Alternatively, note that Theorems 3.2 and 5.1 imply, respectively, that the




FIGURE 4.3. An order filter (left) and an Ish ceiling diagram (right)

number of regions of $\text{Ish}(G)$ and the number of filters in $\Psi^+(G, w)$ (summed over w) are both equal to

$$\sum_{k=0}^{n-1} \text{Stir}(G, n-k) \frac{n!}{(k+1)!}.$$

Hence any injective map between them must be surjective.) \square

It is convenient to express this situation with a picture. Given $w \in \mathfrak{S}(n)$ we draw $w(1), w(2), \dots, w(n)$ on a line. For each j to the right of 1 we draw $j-1$ boxes above the symbol j . If we identify the i th box above j with the hyperplane $x_1 - x_j = i$ then the collection of boxes is exactly $\Psi^+(G, w)$ (we erase the boxes that are not in G); the partial order on boxes increases up and to the left.

In Figure 4.3 (left side) we have drawn the poset $\Psi^+(G, w)$ for the complete graph $G = K_8$ and the permutation $w = 51286347 \in \mathfrak{S}(8)$. The circles (closed up and to the left) indicate an order filter in this poset. This filter defines a region R of $\text{Ish}(K_8) = \text{Ish}(8)$ in the cone wC and its ceilings are the antichain of minimal elements (hollow circles): $x_1 - x_8 = 1$, $x_1 - x_4 = 3$, $x_1 - x_7 = 5$. To simplify the diagram further (right side), we just draw i hollow circles above the symbol j for each ceiling $x_1 - x_j = i$. This is the **Ish ceiling diagram** of the region. We will encode it with the pair (w, ε) where ε_i is the number of circles above the symbol $w(i)$. For the example in Figure 4.3 we have

$$(w, \varepsilon) = (51286347, (0, 0, 0, 1, 0, 0, 3, 5)).$$

Figure 4.4 displays the full arrangement $\text{Ish}(3)$ with regions labeled by ceiling diagrams.

In order to count regions later, here is a purely combinatorial characterization of Ish ceiling diagrams.




FIGURE 4.4. The Ish arrangement $\text{Ish}(3)$ labeled by ceiling diagrams

Definition 4.2. Let $G \subseteq \binom{[n]}{2}$ be a graph and consider a permutation $w \in \mathfrak{S}(n)$. We call the pair (w, ε) an **lsh ceiling diagram** if the vector $\varepsilon = (\varepsilon_1, \dots, \varepsilon_n)$ satisfies:

- $0 \leq \varepsilon_i < w(i)$;
- $\varepsilon_i = 0$ unless $w^{-1}(1) < w^{-1}(i)$;
- If $\varepsilon_i > 0$ then $\varepsilon_i < w(i)$ is an edge in G ;
- the nonzero entries of ε strictly increase.

We will draw the pair (w, ε) by placing $w(1), \dots, w(n)$ on a line and drawing ε_i circles above $w(i)$. By the above remarks, the pair (w, ε) corresponds to a unique region of $\text{lsh}(G)$ with a ceiling $x_1 - x_{w(i)} = i$ for each $\varepsilon_i \neq 0$.

Finally, we can read the recession cone of a region directly from its Ish ceiling diagram.

Lemma 4.4. *Let R be a region of $\text{Ish}(G)$ in the cone wC with ceiling diagram (w, ε) . If k is the maximum index such that $\varepsilon_k \neq 0$ (or $k = w^{-1}(1)$ if ε is the zero vector) then R has $n - k + w^{-1}(1)$ degrees of freedom. In particular, the region R is relatively bounded (has 1 degree of freedom) if and only if $w(1) = 1$ and $\varepsilon_n \neq 0$.*

Proof. Consider $v = (v_1, \dots, v_n) \in \text{Rec}(R)$. Since R is in the cone wC we must have $v_{w(1)} \geq v_{w(2)} \geq \dots \geq v_{w(n)}$. If $\varepsilon_j \neq 0$ then we must also have $v_1 - v_{w(j)} < \varepsilon_j$, which implies that $v_{w^{-1}(1)} = v_{w^{-1}(1)+1} = \dots = v_{w^{-1}(j)}$. Since these are the only constraints on v , we conclude that the recession cone $\text{Rec}(R)$ consists of all vectors of the form


$w \cdot (a_1, a_2, \dots, a_n)$ with $a_1 \geq \dots \geq a_n$ and $a_{w^{-1}(1)} = a_{w^{-1}(1)+1} = \dots = a_k$. The dimension of the cone is therefore $n - (k - w^{-1}(1)) = n - k + w^{-1}(1)$. \square

For example, consider the Ish ceiling diagram (w, ε) in Figure 4.3 and the corresponding region R of $\text{Ish}(8)$. In this case we have $w(1) = 5$ and $k = 8$ is the largest index such that $\varepsilon_k \neq 0$. Hence the recession cone $\text{Rec}(R)$ consists of all vectors $(a, a, a, a, b, a, a, a) \in \mathbb{R}^8$ with $a \geq b$, and it has dimension 2.

4.3. A bijection between dominant regions. The Shi and Ish ceiling diagrams immediately give us a bijection between **dominant** regions of $\text{Shi}(G)$ and $\text{Ish}(G)$ with the same number of ceilings. This bijection does **not** preserve degrees of freedom because it can't: in general $\text{Shi}(G)$ and $\text{Ish}(G)$ have different numbers of relatively bounded dominant regions. For example, $\text{Shi}(3)$ has 2 (see Figure 4.2) and $\text{Ish}(3)$ has 3 (see Figure 4.4).

Theorem 4.5. *Consider a graph $G \subseteq \binom{[n]}{2}$ and an integer c . The deleted arrangements $\text{Shi}(G)$ and $\text{Ish}(G)$ have the same number of **dominant** regions with c ceilings.*

Proof. This is essentially a picture proof. For the identity permutation $w = \mathbf{1}$ we observe that the posets $\Phi^+(G, \mathbf{1})$ and $\Psi^+(G, \mathbf{1})$ look exactly the same, except that one is reflected in a line of slope 1. For example, here are the posets corresponding to the graph $G = \binom{[8]}{2} - \{14, 34, 48, 58\}$; Shi on the left, Ish on the right:



This reflection is an order-reversing bijection between $\Phi^+(G, \mathbf{1})$ and $\Psi^+(G, \mathbf{1})$. Hence it induces a bijection between **ideals in $\Phi^+(G, \mathbf{1})$ with c maximal elements** (dominant $\text{Shi}(G)$ -regions with c ceilings) and **filters in $\Psi^+(G, \mathbf{1})$ with c minimal elements** (dominant $\text{Ish}(G)$ -regions with c ceilings). \square

The number of dominant regions with c ceilings equals the Narayana number $\frac{1}{n} \binom{n}{c} \binom{n-1}{c}$ when G is the complete graph, and equals the binomial coefficient $\binom{n-1}{c}$ when G is the chain $\{12, 23, \dots, (n-1)n\}$. We do not know a closed formula for general G .

Note that the bijection in Theorem 4.5 does **not** extend to other cones wC , since in general the posets $\Phi^+(G, w)$ and $\Psi^+(G, w)$ look very different. Indeed, $\text{Shi}(G)$ and $\text{lsh}(G)$ do **not** have the same number of regions in a given cone wC . (Consider Figures 4.2 and 4.4 with the permutation $w = 132$.)

However, we gain something by summing over the cones wC . Not only do $\text{Shi}(G)$ and $\text{lsh}(G)$ have the same number of (unrestricted) regions with c ceilings, they have the same number of regions with c ceilings and d degrees of freedom. We prove this in the next section using a *non-bijective* method.

5. COUNTING THE REGIONS

In this section we introduce a partition-valued statistic on the regions of $\text{Shi}(G)$ and $\text{lsh}(G)$, and in each case we call this the **ceiling partition** of the region. (This concept is new even for the Shi arrangement.) It turns out that $\text{Shi}(G)$ and $\text{lsh}(G)$ have the same number of regions R with a given ceiling partition π ; moreover, when the partition π has k blocks (i.e. R has $n - k$ ceilings), this number has a beautiful formula: $n!/(n - k + 1)!$. The partition π does **not** determine the degrees of freedom of R . However, we still have a nice formula: The number of regions of $\text{Shi}(G)$ or $\text{lsh}(G)$ with ceiling partition π (with k blocks) and d degrees of freedom equals

$$\frac{d(n - d - 1)!(k - 1)!}{(n - k - 1)!(k - d)!}.$$

This completes the proof of the Main Theorem. At the end we make comments and suggestions for future research.

5.1. Ceiling partitions. To each region R of $\text{Shi}(G)$ or $\text{lsh}(G)$ we associate a partition of the set $[n]$, called its **ceiling partition**. We note that this partition may be *nesting*, and in general every partition of $[n]$ will occur. The ceiling partition is determined by the ceilings of R and the cone wC in which R occurs; thus we can read it from the ceiling diagram. We will see that the correct language for ceiling partitions is the **endpoint notation** (\mathbf{a}, \mathbf{b}) , discussed in Section 2.

Definition 5.1.

- (1) Let R be a region of $\text{Shi}(G)$ with ceiling diagram (w, π) . Then the **ceiling partition** of R is $w \cdot \pi$ (w acting on π). That is, the ceiling partition has $w(i)$ and $w(j)$ in a block whenever i and j are in a block of π . For example, the region in Figure 4.1 has ceiling partition $\{\{1, 6\}, \{2\}, \{3, 7\}, \{4\}, \{5, 8\}\}$, with endpoint notation $(\mathbf{a}, \mathbf{b}) = (135, 678)$. Note that the ceiling partition (\mathbf{a}, \mathbf{b}) has c arcs if and only if R has c ceilings.

- (2) Let R be a region of $\text{Ish}(G)$ with ceiling diagram (w, ε) . We define a pair of vectors (\mathbf{a}, \mathbf{b}) such that a_i is the i th nonzero entry of ε , which occurs in position $w^{-1}(b_i)$. The conditions of Definition 4.2 guarantee that (\mathbf{a}, \mathbf{b}) is the endpoint notation for a partition, which we call the **ceiling partition** of R . For example, the region shown in Figure 4.3 has ceiling partition $(\mathbf{a}, \mathbf{b}) = (135, 847)$ since there is one circle above 8, three above 4, and five above 7. Again, the ceiling partition has c arcs if and only if R has c ceilings.

5.2. Counting Shi and Ish regions. Let c and d be integers. We separately count the regions of $\text{Shi}(G)$ and $\text{Ish}(G)$ with c ceilings and d degrees of freedom, and observe that they are the same. This completes the proof of the Main Theorem.

Theorem 5.1. *Fix a graph $G \subseteq \binom{[n]}{2}$ and let \mathcal{A} be either $\text{Shi}(G)$ or $\text{Ish}(G)$. Let (\mathbf{a}, \mathbf{b}) be a partition of $[n]$ with k blocks ($n - k$ arcs) and consider an integer $1 \leq d \leq k$. There exists a region of \mathcal{A} with ceiling partition (\mathbf{a}, \mathbf{b}) if and only if we have $a_i b_i \in G$ for all i , in which case:*

- (1) *The number of regions of \mathcal{A} with ceiling partition (\mathbf{a}, \mathbf{b}) is*

$$\frac{n!}{(n - k + 1)!}.$$

- (2) *The number of regions of \mathcal{A} with ceiling partition (\mathbf{a}, \mathbf{b}) and d degrees of freedom is*

$$(5.1) \quad \frac{d(n - d - 1)!(k - 1)!}{(n - k - 1)!(k - d)!}.$$

To obtain the number of regions with c ceilings and d degrees of freedom, sum (5.1) over G -partitions (\mathbf{a}, \mathbf{b}) with $k = n - c$ blocks.

Proof. First we deal with $\mathcal{A} = \text{Shi}(G)$. Recall that a Shi ceiling diagram (w, π) is a nonnesting partition π whose blocks are increasing with respect to the permutation w . Thus, to create a ceiling diagram (region) with ceiling partition (\mathbf{a}, \mathbf{b}) , we must first choose a nonnesting partition π_0 with the same block sizes as (\mathbf{a}, \mathbf{b}) and then put the labels from each block of (\mathbf{a}, \mathbf{b}) (increasingly) in a block of π_0 . So suppose that (\mathbf{a}, \mathbf{b}) has r_i blocks of size i . By Lemma 2.1 there are

$$\frac{n!}{(n - k + 1)!r_1!r_2!\cdots r_n!}$$

ways to choose π_0 . Then, there are $r_1!r_2!\cdots r_n!$ ways to map each block of (\mathbf{a}, \mathbf{b}) to a block of π_0 with the same size. This proves (1). To prove (2), note that the region (w, π) has d degrees of freedom if and only if the nonnesting partition π has d connected components. Apply the same argument as above, but use Lemma 2.2.

Next we deal with $\mathcal{A} = \text{Ish}(G)$. We wish to create an Ish region (w, ε) with ceiling partition (\mathbf{a}, \mathbf{b}) . To do this, we choose $w(1), \dots, w(n)$ and then place a_i circles above the symbol b_i . This will be an Ish ceiling diagram as long as the symbols b_i occur in order, to the right of 1. That is, the permutation w must satisfy

$$(5.2) \quad w^{-1}(1) < w^{-1}(b_1) < \dots < w^{-1}(b_{n-k}).$$

There are $\binom{n}{n-k+1}$ ways to place these symbols and then $(k-1)!$ ways to place the remaining symbols, proving (1). To prove (2), recall that (w, ε) has $n - j + w^{-1}(1)$ degrees of freedom, where j is the largest index such that $\varepsilon_j \neq 0$. In our case $j = w^{-1}(b_{n-k})$, so w must satisfy the condition $w^{-1}(b_{n-k}) - w^{-1}(1) = n - d$. First we can choose the pair $(w^{-1}(1), w^{-1}(b_{n-k}))$ in d ways. Having done this, the rest of the permutation is subject to (5.2). There are $\binom{n-d-1}{n-k-1}$ ways to place symbols b_1, \dots, b_{n-k-1} (left to right) in the $n - d - 1$ positions between 1 and b_{n-k} , and then there are $(k-1)!$ ways to place the remaining $k-1$ symbols. The result follows. \square

Once again, note that this proof was more direct for Ish than for Shi. In fact, the calculation of formula (5.1) for Ish was the inspiration for Lemma 2.2.

5.3. Concluding remarks. The notion of a ceiling partition has independent interest, beyond the proof of Theorem 5.1. In particular, it leads to a new proof that the Shi arrangement $\text{Shi}(n)$ has $(n+1)^{n-1}$ regions. Consider the collection of maps from $[n]$ into a set of size x . On one hand, there are x^n of these. On the other hand, there are $\text{Stir}(n, k)x(x-1)\cdots(x-k+1)$ such maps with image of size k , where $\text{Stir}(n, k)$ is the number of partitions of $[n]$ into k blocks (fibers). This proves the famous polynomial identity:

$$x^n = \sum_{k=1}^n \text{Stir}(n, k)x(x-1)\cdots(x-k+1).$$

Dividing by x and substituting $x = n+1$ yields

$$(n+1)^{n-1} = \sum_{k=1}^n \text{Stir}(n, k) \frac{n!}{(n-k+1)!},$$

where the right hand side counts regions of $\text{Shi}(n)$ by the number k of blocks in their ceiling partition.

Finally, here are some problems for future research.

- (1) The original motivation for this paper was to find a bijection between regions of Shi and Ish. We solved this problem for dominant regions, but not in general. Based on Theorem 5.1, one should look for a bijection between Shi ceiling diagrams (w, π) and Ish ceiling diagrams (w, ε) with a fixed ceiling

partition (\mathbf{a}, \mathbf{b}) and d degrees of freedom. Note that this bijection **cannot** preserve the permutation w .

- (2) Following Theorem 3.2, find a direct bijection between points of the finite vector space \mathbb{F}_p^n in the complements of the Shi arrangement $\text{Shi}(G)_p$ and the Ish arrangement $\text{Ish}(G)_p$.
- (3) The Shi arrangement is a famous example of a **free hyperplane arrangement**. Investigate the freeness of Ish arrangements.
- (4) Define and study an extended Ish arrangement corresponding to the **extended Shi arrangement**:

$$\text{Shi}(n, m) := \{x_i - x_j = a : 1 \leq i < j \leq n, -m + 1 \leq a \leq m\}.$$

- (5) To what extent do the results of this paper apply to other deformations of the Coxeter arrangement?
- (6) The deleted Shi arrangements exist for arbitrary crystallographic reflection groups. Define and study Ish arrangements for other reflection groups. Ish arrangements were invented to study q, t -Catalan numbers; this feature should extend to other types.

6. ACKNOWLEDGEMENTS

The authors are grateful to Christos Athanasiadis, Susanna Fishel, Christian Krattenthaler, Vic Reiner, and Richard Stanley for helpful conversations. This project arose out of an AMS sectional meeting held at Penn State in October 2009.

REFERENCES

- [1] D. ARMSTRONG. Hyperplane arrangements and diagonal harmonics. In preparation, 2010.
- [2] C. ATHANASIADIS AND S. LINUSSON. A simple bijection for the regions of the Shi arrangement of hyperplanes. *Discrete Math.*, **204** (1999) pp. 27–39.
- [3] C. A. ATHANASIADIS. Characteristic polynomials of subspace arrangements and finite fields. *Adv. Math.*, **122** (1996) pp. 193–233.
- [4] C. A. ATHANASIADIS. On free deformations of the braid arrangement. *European J. Combin.*, **19** (1997) pp. 7–18.
- [5] H. CRAPO AND G.-C. ROTA. *On the Foundations of Combinatorial Theory: Combinatorial Geometries*. MIT Press, Cambridge, MA (1970). Preliminary edition.
- [6] P. HEADLEY. On a family of hyperplane arrangements related to affine Weyl groups. *J. Alg. Combin.*, **6**, 4 (1997) pp. 331–338.
- [7] G. KREWERAS. Sur les partitions non croisées d’un cycle. *Discrete Math.*, **1**, 4 (1972) pp. 333–350.
- [8] P. ORLIK AND K. TERAOKA. *Arrangements of Hyperplanes*. Springer-Verlag, New York, NY (1992).
- [9] A. POSTNIKOV AND R. STANLEY. Deformations of Coxeter hyperplane arrangements. *J. Combin. Theory Ser. A*, **91** (2000) pp. 544–597.

- [10] B. RHOADES. Enumeration of connected Catalan objects by type (2010). Available at <http://arxiv.org/abs/1005.2553>.
- [11] J.-Y. SHI. *The Kazhdan-Lusztig cells in certain affine Weyl groups*. Lecture Notes in Mathematics, no. 1179, Springer-Verlag, Berlin/Heidelberg/New York (1986).
- [12] R. STANLEY. *Enumerative Combinatorics*, vol. 1. Cambridge University Press, Cambridge (1997).
- [13] H. TERAOKA. Arrangements of hyperplanes and their freeness I, II. *J. Fac. Sci. Univ. Tokyo Sect. IA Math.*, **27** (1980) pp. 293–320.
- [14] M. WACHS AND J. WALKER. On geometric semilattices. *Order*, **2**, 4 (1986) pp. 367–385.
- [15] T. ZASLAVSKY. Facing up to arrangements: a face-count for partitions of space by hyperplanes. *Mem. Amer. Math. Soc.*, **1** (1975).

DREW ARMSTRONG, DEPARTMENT OF MATHEMATICS, UNIVERSITY OF MIAMI, CORAL GABLES, FL, 33146

E-mail address: `armstrong@math.miami.edu`

BRENDON RHOADES, DEPARTMENT OF MATHEMATICS, MIT, CAMBRIDGE, MA, 02139

E-mail address: `brhoades@math.mit.edu`