

Permutation Enumeration and Symmetric Functions

Jeff Remmel

Department of Mathematics
University of California

Joint work with Tony Mendes



$$\sigma = \sigma_1 \dots \sigma_n \in S_n$$

$$Des(\sigma) = \{i : \sigma_i > \sigma_{i+1}\}$$

$$des(\sigma) = |Des(\sigma)|$$

$$maj(\sigma) = \sum_{i \in Des(\sigma)} i$$

$$rlmaj(\sigma) = \sum_{i \in Des(\sigma)} n - i$$

$$inv(\sigma) = \sum_{i < j} \chi(\sigma_i > \sigma_j)$$

$$exc(\sigma) = |\{i : i < \sigma_i\}|$$

$$Rise(\sigma) = \{i : \sigma_i < \sigma_{i+1}\}$$

$$rise(\sigma) = 1 + |Rise(\sigma)|$$

$$comaj(\sigma) = \sum_{i \in Rise(\sigma)} i$$

$$rlcomaj(\sigma) = \sum_{i \in Rise(\sigma)} n - i$$

$$coinv(\sigma) = \sum_{i < j} \chi(\sigma_i < \sigma_j)$$

$$dec(\sigma) = |\{i : i > \sigma_i\}|$$

where for any statement A , $\chi(A) = 1$ if A is true and $\chi(A) = 0$ if A is false. Also if $\alpha^1, \dots, \alpha^k \in S_n$, then we shall write

$$comdes(\alpha^1, \dots, \alpha^k) = |\bigcap_{i=1}^k Des(\alpha^i)|.$$

$$[n]_q = 1 + q + \cdots + q^{n-1} = \frac{1-q^n}{1-q}$$

$$[n]_q! = [n]_q[n-1]_q \cdots [1]_q$$

$${[n] \atop [k]}_q = \frac{[n]_q!}{[k]_q![n-k]_q!}$$

$${n \atop \lambda_1, \dots, \lambda_\ell}_q = \frac{[n]_q!}{[\lambda_1]_q! \cdots [\lambda_\ell]_q!}$$

$$[n]_{p,q} = p^{n-1} + p^{n-2}q + \cdots + p^1q^{n-2} + q^{n-1} = \frac{p^n - q^n}{p - q}.$$

$$(1) \sum_{n=0}^{\infty} \frac{u^n}{n!} \sum_{\sigma \in S_n} x^{des(\sigma)} = \frac{1-x}{-x+e^{u(x-1)}}$$

(2) (Carlitz 1970)

$$\sum_{n=0}^{\infty} \frac{u^n}{(n!)^2} \sum_{(\sigma, \tau) \in S_n \times S_n} x^{comdes(\sigma, \tau)} = \frac{1-x}{-x+J(u(x-1))}.$$

$$(3) \text{ (Stanley 1979)} \quad \sum_{n=0}^{\infty} \frac{u^n}{[n]_q!} \sum_{\sigma \in S_n} x^{des(\sigma)} q^{inv(\sigma)} = \frac{1-x}{-x+e_q(u(x-1))}.$$

$$\sum_{n=0}^{\infty} \frac{u^n}{[n]_q!} \sum_{\sigma \in S_n} x^{des(\sigma)} q^{coinv(\sigma)} = \frac{1-x}{-x+E_q(u(x-1))}.$$

(4) (Fedou and Rawlings 1995)

$$\sum_{n=0}^{\infty} \frac{u^n}{[n]_q! [n]_p!} \sum_{(\sigma, \tau) \in S_n \times S_n} x^{comdes(\sigma, \tau)} q^{inv(\sigma)} p^{inv(\tau)} = \frac{1-x}{-x+J_{q,p}(u(x-1))}.$$

$$J(u) = \sum_{n \geq 0} \frac{u^n}{n! n!}, \quad e_q(u) = \sum_{n=0}^{\infty} \frac{u^n}{[n]_q!} q^{\binom{n}{2}},$$

$$E_q(u) = \sum_{n=0}^{\infty} \frac{u^n}{[n]_q!}, \text{ and} \quad J_{q,p}(u) = \sum_{n=0}^{\infty} \frac{u^n}{[n]_q! [n]_p!} q^{\binom{n}{2}} p^{\binom{n}{2}}.$$

6) Foata-Han (1997)

Let $(x, q)_0 = 1$ and $(x, q)_n = (1 - x)(1 - qx) \cdots (1 - q^{n-1}x)$ for $n > 0$.

$$C_n(z, x, q, y, p) = \sum_{(\sigma, \tau) \in S_n \times S_n} z^{\text{comdes}(\sigma^{-1}, \tau^{-1})} x^{\text{des}(\sigma)} q^{\text{rlmaj}(\sigma)} y^{\text{rise}(\tau)} p^{\text{rlcomaj}(\tau)}.$$

$$\sum_{n \geq 0} t^n \frac{C_n(z, x, q, y, p)}{(x, q)_{n+1}(y, p)_{n+1}} = \sum_{i, j \geq 0} \frac{x^i y^j}{1 + \sum_{n \geq 1} (t(z-1))^{n-1} \begin{bmatrix} i+1 \\ n \end{bmatrix}_q \begin{bmatrix} j+n \\ n \end{bmatrix}_p}.$$

7) Remmel-Mendes

$$R_n(z, x, q, y, p, Q, P) = \sum_{(\alpha, \beta, \gamma, \delta) \in S_n^4} z^{comdes(\alpha^{-1}, \beta^{-1}, \gamma, \delta)} x^{des(\alpha)} q^{rlmaj(\alpha)} y^{rise(\beta)} p^{rlcomaj(\beta)} Q^{inv(\gamma)} P^{coinv(\delta)}$$

and set

$$F^{i,j}(t, q, p, Q, P) = \sum_{n \geq 0} t^n \frac{q^{\binom{n}{2}} Q^{\binom{n}{2}} \left[\begin{array}{c} i+1 \\ n \end{array} \right]_q \left[\begin{array}{c} j+n \\ n \end{array} \right]_p}{[n]_Q! [n]_P!}.$$

Then we can use the combinatorial mechanism described above with 4-tuples of permutations instead of pairs of permutations to prove that

$$\sum_{n \geq 0} \frac{R_n(z, x, q, y, p, Q, P) t^n}{[n]_Q! [n]_P! (x, q)_{n+1} (y, p)_{n+1}} = \sum_{i,j \geq 0} x^i y^j \frac{1-t}{-t + F^{i,j}(t(z-1), q, p, Q, P)}.$$

Elementary & homogeneous symmetric functions

The n^{th} elementary symmetric function e_n is defined by

$$\sum_{n \geq 0} e_n t^n = \prod_i (1 + x_i t).$$

The n^{th} homogeneous symmetric function h_n is defined by

$$\sum_{n \geq 0} h_n t^n = \prod_i \frac{1}{1 - x_i t}.$$

The n^{th} power symmetric function p_n is defined by

$$p_n = \sum_i x_i^n.$$

If $\lambda = (\lambda_1, \dots, \lambda_\ell)$ is a partition, then

$$h_\lambda = \prod_{i=1}^k h_{\lambda_i}, \quad e_\lambda = \prod_{i=1}^k e_{\lambda_i}, \quad \text{and} \quad p_\lambda = \prod_{i=1}^k p_{\lambda_i},$$

Brenti(1993) Define a ring homomorphism $\xi : \Lambda \rightarrow Q[x]$ by setting

$$\xi(e_k) = \frac{(x-1)^{k-1}}{k!}$$

where e_k is the k -th elementary symmetric function and $\xi(e_0) = 1$.


$$n! \xi(h_n) = \sum_{\sigma \in S_n} x^{\text{des}(\sigma)} \text{ and } \frac{n!}{z_\lambda} \xi(p_\lambda) = \sum_{\sigma \in S_n(\lambda)} x^{\text{exc}(\sigma)} \quad (1)$$

where if $\lambda = (1^{m_1}, 2^{m_2}, \dots, n^{m_n})$ is a partition of n , then $S_n(\lambda)$ is the set of permutations in S_n with cycle type λ , and


$$z_\lambda = \prod_{i=1}^n i^{m_i} m_i!$$

λ -Brick Tabloids and Weighted λ -Brick Tabloids. Suppose that $\lambda = (1, 1, 2, 2)$ and $\mu = (2, 4)$.


$$B_{\lambda, \mu} = 4 \text{ and } w(B)_{\lambda, \mu} = 10$$




$$w(\mathbf{T}_1) = 2$$



$$w(\mathbf{T}_2) = 2$$



$$w(\mathbf{T}_3) = 4$$



$$w(\mathbf{T}_4) = 2$$

$$\begin{aligned} h_\mu(\bar{x}) &= \sum_{\mu \vdash n} (-1)^{n-\ell(\lambda)} B_{\lambda,\mu} e_\lambda(\bar{x}) \\ p_\mu(\bar{x}) &= \sum_{\mu \vdash n} (-1)^{n-\ell(\lambda)} w(B)_{\lambda,\mu} e_\lambda(\bar{x}) \end{aligned}$$

(Egecioglu and Remmel 1991)

A link from Λ to permutation enumeration

For $\sigma_1 \cdots \sigma_n \in S_n$,

$des(\sigma)$ is the number of times $\sigma_i > \sigma_{i+1}$,

$ris(\sigma)$ is the number of times $\sigma_i < \sigma_{i+1}$ where $\sigma_{n+1} = n + 1$.

Ex. Let $\sigma = 12 \text{ } 9 \text{ } 7 \text{ } 2 \text{ } 6 \text{ } 8 \text{ } 10 \text{ } 1 \text{ } 3 \text{ } 4 \text{ } 11 \text{ } 5$.

Then $des(\sigma) = 5$ and $ris(\sigma) = 7$.

Let $f_1 : \{0, 1, \dots\} \rightarrow \mathbb{Q}[x, y]$ such that:

$$f_1(n) = \begin{cases} 1 & \text{if } n = 0 \\ y(x - y)^{n-1} & \text{if } n \geq 1. \end{cases}$$

Define $\xi^{f_1} : \Lambda \rightarrow \mathbb{Q}[x, y]$ as a homomorphism such that

$$\xi^{f_1}(e_n) = \frac{(-1)^{n-1}}{n!} f_1(n).$$

Theorem.

$$n! \xi^{f_1}(h_n) = \sum_{\sigma \in S_n} x^{\text{des}(\sigma)} y^{\text{ris}(\sigma)}.$$

Proof.

$$\begin{aligned}
 n! \xi^{f_1}(h_n) &= n! \sum_{\lambda \vdash n} (-1)^{n-\ell(\lambda)} B_{\lambda,n} \xi^{f_1}(e_\lambda) \\
 &= n! \sum_{\lambda \vdash n} (-1)^{n-\ell(\lambda)} B_{\lambda,n} \prod_{i=1}^{\ell(\lambda)} \frac{(-1)^{\lambda_i-1}}{\lambda_i!} f_1(\lambda_i) \\
 &= \sum_{\lambda \vdash n} \binom{n}{\lambda_1, \dots, \lambda_\ell} B_{\lambda,n} f_1(\lambda_1) \cdots f_1(\lambda_\ell).
 \end{aligned}$$

We have

$$n! \xi^{f_1}(h_n) = \sum_{\lambda \vdash n} \binom{n}{\lambda_1, \dots, \lambda_\ell} B_{\lambda, n} f_1(\lambda_1) \cdots f_1(\lambda_\ell)$$

from which we create the following objects:

We have

$$n! \xi^{f_1}(h_n) = \sum_{\lambda \vdash n} \binom{n}{\lambda_1, \dots, \lambda_\ell} B_{\lambda, \textcolor{red}{n}} f_1(\lambda_1) \cdots f_1(\lambda_\ell)$$

from which we create the following objects:

⋮ ⋮	⋮ ⋮ ⋮		⋮ ⋮	⋮ ⋮
-----	-------	--	-----	-----

We have

$$n! \xi^{f_1}(h_n) = \sum_{\lambda \vdash n} \binom{n}{\lambda_1, \dots, \lambda_\ell} B_{\lambda, n} f_1(\lambda_1) \cdots f_1(\lambda_\ell)$$

from which we create the following objects:

11	6	2	10	5	3	1	8	12	9	7	4
----	---	---	----	---	---	---	---	----	---	---	---

We have

$$n! \xi^{f_1}(h_n) = \sum_{\lambda \vdash n} \binom{n}{\lambda_1, \dots, \lambda_\ell} B_{\lambda, n} f_1(\lambda_1) \cdots f_1(\lambda_\ell)$$

from which we create the following objects:

x	x	y	$-y$	x	$-y$	y	y	$-y$	y	x	y
11	6	2	10	5	3	1	8	12	9	7	4

$$f_1(n) = y(x - y)^{n-1}$$

Let \mathcal{T}_{f_1} be the set of objects created in this way.

$T = (B, \sigma, L) \in \mathcal{T}_{f_1}$ if

- (1) $B = (b_1, \dots, b_k)$ is a brick tabloid of shape (n) .
- (2) σ is a permutation which is decreasing within each brick.
- (3) L is a labeling such that the last cell of each brick is labeled with a y and every other cell is labeled with x or $-y$.

The weight $w(T)$ of $T \in \mathcal{T}_{f_1}$, $w(T)$, is the product of $x, -y$, and y labels.

An involution $I : \mathcal{T}_{f_1} \rightarrow \mathcal{T}_{f_1}$ such that

(a) If $I(T) \neq T$, then $w(T) = -w(I(T))$.

(b) If $I(T) = T$, then $w(T)$ is positive.

I will show that

$$\sum_{T \in \mathcal{T}_{f_1}} w(T) = \sum_{T \in \mathcal{T}_{f_1}, I(T)=T} w(T).$$

To define $I(T)$ where $T = (B, \sigma, L)$, scan the cells of T from left to right looking for the first cell c such that either

- (i) c is labeled with $-y$ in which case we break the brick b containing c into two bricks b^* and b^{**} where b^* contains all the cells of b up to and including c and b^{**} consists of the remaining cells of b and then we change the label on cell c from $-y$ to y or
- (ii) c is the last cell of a brick b_i which is followed by another a brick b_{i+1} such that σ is decreasing in all the cells corresponding to b_i and b_{i+1} in which case we replace b_i and b_{i+1} by a single brick b and change the label on cell c from y to $-y$.

If neither (i) or (ii) applies, $I(T) = T$.

x	x	y	$-y$	x	$-y$	y	y	$-y$	y	x	y
11	6	2	10	5	3	1	8	12	9	7	4

is sent to

x	x	y	y	x	$-y$	y	y	$-y$	y	x	y
11	6	2	10	5	3	1	8	12	9	7	4

Fixed points of I : (a) no $-y$'s and (b) increases between bricks.

x	:	x	:	y	x	:	x	:	x	:	y	y	x	:	y	x	:	y
11	:	6	:	2	10	:	5	:	3	:	1	8	12	:	7	9	:	4

A fixed point can be read as an element in S_n :

$$11 \text{ } 6 \text{ } 2 \text{ } 10 \text{ } 5 \text{ } 3 \text{ } 1 \text{ } 8 \text{ } 12 \text{ } 7 \text{ } 9 \text{ } 4$$

Therefore,

$$n! \xi^{f_1}(h_n) = \sum_{T \in \mathcal{T}_{f_1}} w(T) = \sum_{\substack{T \text{ is a} \\ \text{fixed point}}} w(T) = \sum_{\sigma \in S_n} x^{\text{des}(\sigma)} y^{\text{ris}(\sigma)}.$$

□

This gives a generating function:

$$\begin{aligned}
 \sum_{n \geq 0} \frac{t^n}{n!} \sum_{\sigma \in S_n} x^{des(\sigma)} y^{ris(\sigma)} &= \xi^{f_1} \left(\sum_{n \geq 0} h_n t^n \right) \\
 &= \xi^{f_1} \left(\sum_{n \geq 0} e_n (-t)^n \right)^{-1} \\
 &= \left(1 - \sum_{n \geq 1} \frac{t^n}{n!} f_1(n) \right)^{-1} \\
 &= \frac{x - y}{x - ye^{t(x-y)}}.
 \end{aligned}$$

We just

1. Defined f on $\{0, 1, \dots\}$ and ξ^f on Λ such that

$$\xi^f(e_n) = \frac{(-1)^{n-1}}{n!} f(n)$$

2. Applied ξ^f to $n!h_n$ and decorated brick tabloids
3. Performed an involution to find objects corresponding to permutations
4. Found a generating function from the h_n and e_n relationship

More weighting functions

$\sigma_1 \cdots \sigma_n \in S_n$ is alternating if $\sigma_{i-1} > \sigma_i$ and $\sigma_i < \sigma_{i+1}$ for even i .

Ex. 8 6 7 4 5 2 3 1 12 9 11 10 is (even) alternating.

$$\text{Let } f_2(n) = \begin{cases} 0 & n \text{ odd,} \\ (-1)^{\frac{n}{2}-1} & n \text{ even.} \end{cases} \quad \text{and} \quad \xi^{f_2}(e_n) = \frac{(-1)^{n-1}}{n!} f_2(n).$$

Theorem. Let A_n be the number of even alternating permutations of n . Then $n! \xi^{f_2}(h_n) = A_n$.

Proof. We have,

$$n! \xi^{f_2}(h_n) = \sum_{\lambda \vdash n} \binom{n}{\lambda_1, \dots, \lambda_\ell} B_{\lambda, n} f_2(\lambda_1) \cdots f_2(\lambda_\ell)$$

from which we create the following objects:

$$f_2(n) = 0 \text{ if } n \text{ odd and } f_2(n) = (-1)^{\frac{n}{2}-1} \text{ if } n \text{ even.}$$

$$n! \xi^{f_2}(h_n) = \sum_{\lambda \vdash n} \binom{n}{\lambda_1, \dots, \lambda_\ell} B_{\lambda, \textcolor{red}{n}} f_2(\lambda_1) \cdots f_2(\lambda_\ell)$$

from which we create the following objects:

;	;	;	;	;	;	;	;
---	---	---	---	---	---	---	---

$f_2 = 0$ if n odd and $f_2 = (-1)^{\frac{n}{2}-1}$ if n even.

$$n! \xi^{f_2}(h_n) = \sum_{\lambda \vdash n} \binom{n}{\lambda_1, \dots, \lambda_\ell} B_{\lambda, n} f_2(\lambda_1) \cdots f_2(\lambda_\ell)$$

from which we create the following objects:

11	2	10	8	6	5	3	1	12	9	7	4
----	---	----	---	---	---	---	---	----	---	---	---

$f_2 = 0$ if n odd and $f_2 = (-1)^{\frac{n}{2}-1}$ if n even.

$$n! \xi^{f_2}(h_n) = \sum_{\lambda \vdash n} \binom{n}{\lambda_1, \dots, \lambda_\ell} B_{\lambda, n} f_2(\lambda_1) \cdots f_2(\lambda_\ell)$$

from which we create the following objects:

$\begin{array}{ c c } \hline 1 & 1 \\ \hline 11 & 2 \\ \hline \end{array}$	$\begin{array}{ c c c c c c c c } \hline 1 & -1 & 1 & -1 & 1 & 1 \\ \hline 10 & 8 & 6 & 5 & 3 & 1 \\ \hline \end{array}$	$\begin{array}{ c c c c c c c c } \hline 1 & -1 & 1 & 1 \\ \hline 12 & 9 & 7 & 4 \\ \hline \end{array}$
--	--	---

$f_2(n) = 0$ if n odd and $f_2(n) = (-1)^{\frac{n}{2}-1}$ if n even.

Perform a similar involution: scan for -1 or a decrease between two bricks:

1	1	1	-1	1	1	1	1	1	1	1	1
11	2	10	8	6	5	3	1	12	9	7	4

is sent to

1	1	1	1	1	-1	1	1	1	-1	1	1
11	2	10	8	6	5	3	1	12	9	7	4

A fixed point:

1	:	1		1	:	1		1	:	1		1	:	1		1	:	1				
11	:	2		10	:	6		8	:	3		5	:	1		12	:	7		9	:	4

The corresponding permutation:

$$\begin{array}{cccccccccc} \textcolor{blue}{11} & \textcolor{red}{2} & \textcolor{black}{10} & \textcolor{blue}{6} & \textcolor{black}{8} & \textcolor{red}{3} & \textcolor{blue}{12} & \textcolor{red}{7} & \textcolor{black}{9} & \textcolor{blue}{4} \end{array}$$

Summing all fixed points gives the desired result. □

The resultant generating function:

$$\begin{aligned}
 \sum_{n \geq 0} \xi_{f_2}(h_n) &= \sum_{n \geq 0} \frac{t^{2n}}{(2n)!} A_{2n} \\
 &= \frac{1}{1 + \sum_{n \geq 1} (-t)^n \xi_{f_2}(e_n)} \\
 &= \frac{1}{1 + \sum_{n \geq 1} (-t)^{2n} \frac{(-1)^{2n-1}}{(2n)!} (-1)^{n-1}} \\
 &= \frac{1}{1 + \sum_{n \geq 1} \frac{(-1)^n}{(2n)!}} \\
 &= \frac{1}{\cos(t)} \\
 &= \sec(t).
 \end{aligned}$$

Next we want to show that $(2n+1)!\xi_{f_2}(p_{2n+2}) = A_{2n+1}$.

If $\mu = (\mu_1, \dots, \mu_k)$, then $2\mu = (2\mu_1, \dots, 2\mu_k)$.

$$\begin{aligned}
 (2n+1)!\xi_{f_2}(p_{2n+2}) &= (2n+1)! \sum_{\mu \vdash n+1} (-1)^{2n+2-\ell(\mu)} w(B_{2\mu, (2n)}) \xi_{f_2}(e_{2\mu}) = \\
 (2n+1)! \sum_{\mu \vdash n+1} (-1)^{2n+2-\ell(\mu)} &\quad \sum_{B=(b_1, \dots, b_{\ell(\mu)}) \in \mathcal{B}_{\mu, (n+1)}} 2b_{\ell(\mu)} \prod_{i=1}^{\ell(\mu)} \frac{(-1)^{2b_i-1}}{2b_i!} (-1)^{b_i-1} = \\
 \sum_{\mu \vdash n+1} &\quad \sum_{B=(b_1, \dots, b_{\ell(\mu)}) \in \mathcal{B}_{\mu, (n+1)}} \binom{2n+1}{2b_1, \dots, 2b_{\ell(\mu)-1}, 2b_{\ell(\mu)}-1} \prod_{i=1}^{\ell(\mu)} (-1)^{b_i-1}.
 \end{aligned}$$

$f_2 = 0$ if n odd and $f_2 = (-1)^{\frac{n}{2}-1}$ if n even.

$$\sum_{\mu \vdash n+1} \sum_{B=(b_1, \dots, b_{\ell(\mu)}) \in \mathcal{B}_{\mu, (n)}} \binom{2n+1}{2b_1, \dots, 2b_{\ell(\mu)-1}, 2b_{\ell(\mu)} - 1} \prod_{i=1}^{\ell(\mu)} (-1)^{b_i-1}.$$

$$\sum_{\mu \vdash n+1} \sum_{B = (b_1, \dots, b_{\ell(\mu)}) \in \mathcal{B}_{\mu, (n)}} \binom{2n+1}{2b_1, \dots, 2b_{\ell(\mu)-1}, 2b_{\ell(\mu)} - 1} \prod_{i=1}^{\ell(\mu)} (-1)^{b_i-1}.$$

11	·	2	10	·	8	·	6	·	5	·	3	·	1	9	·	7	·	4	·
----	---	---	----	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---

$$\sum_{\mu \vdash n+1} \sum_{B = (b_1, \dots, b_{\ell(\mu)}) \in \mathcal{B}_{\mu, (n)}} \binom{2n+1}{2b_1, \dots, 2b_{\ell(\mu)-1}, 2b_{\ell(mu)} - 1} \prod_{i=1}^{\ell(\mu)} (-1)^{b_i-1}.$$

1	:	1		1	:	-1	:	1	:	-1	:	1	:	1		1	:	-1	:	1	:	1	:	1
11	:	2		10	:	8	:	6	:	5	:	3	:	1		9	:	7	:	4	:	1	:	1

Perform a similar involution: scan for -1 or a decrease between two bricks:

1	1	1	-1	1	1	1	1
11	2	10	8	6	5	3	1

is sent to

1	1	1	1	1	-1	1	1	1	1
11	2	10	8	6	5	3	1	9	7

A fixed point:

1	1	1	1	1	1	1	1	1
11	2	10	6	8	3	5	1	7

The corresponding permutation:

$$\begin{array}{cccccccccc} 11 & 2 & 10 & 6 & 8 & 3 & 7 & 4 & 9 \end{array}$$

Summing all fixed points gives the desired result. □

The resultant generating function:

$$\begin{aligned}
 \sum_{n \geq 1} \xi_{f_2}(p_n) &= \sum_{n \geq 0} \frac{t^{2n+2}}{(2n+1)!} A_{2n+1} \\
 &= \frac{\sum_{n \geq 1} (-1)^{n-1} \nu(n) \xi_{f_2}(e_n)}{1 + \sum_{n \geq 1} (-t)^n \xi_{f_2}(e_n)} \\
 &= \frac{\sum_{n \geq 1} (-1)^{2n-1} 2n \frac{(-1)^{2n-1}}{2n!} (-1)^{n-1}}{1 + \sum_{n \geq 1} (-t)^{2n} \frac{(-1)^{2n-1}}{(2n)!} (-1)^{n-1}} \\
 &= \frac{\sum_{n \geq 1} \frac{(-1)^{n-1} t^{2n}}{(2n-1)!}}{1 + \sum_{n \geq 1} \frac{(-1)^n}{(2n)!}} \\
 &= \frac{t \sin(t)}{\cos(t)}.
 \end{aligned}$$

$$\sum_{n \geq 0} A_{2n+1} \frac{t^{2n+2}}{(2n+1)!} = \frac{t \sin(t)}{\cos(t)}.$$

Dividing both sides by t , we obtain

$$\sum_{n \geq 0} A_{2n+1} \frac{t^{2n+1}}{(2n+1)!} = \frac{\sin(t)}{\cos(t)} = \tan(t).$$

Excedances

For $\sigma_1 \cdots \sigma_n \in S_n$, $exc(\sigma)$ is the number of times $i < \sigma_{i+1}$.

Example. Let $\sigma = (1, 5, 3, 7) \ (2, 6, 8, 10) \ (4) \ (9, 12, 11)$

Then $exc(\sigma) = 6$.

Let $S_n(\mu)$ denote the set of $\sigma \in S_n$ such that the lengths of the cycles in σ induce the partition μ .

Define $\xi^{f_1} : \Lambda \rightarrow \mathbb{Q}[x, y]$ as a homomorphism such that

$$\xi(e_n) = \frac{(-1)^{n-1}}{n!}(x-1)^{n-1}.$$

If $\mu = (1^{\alpha_1} 2^{\alpha_2} \cdots n^{\alpha_n}) \vdash n$, then $z_\mu = 1^{\alpha_1} 2^{\alpha_2} \cdots n^{\alpha_n} \alpha_1! \alpha_2! \cdots \alpha_n!$

Theorem.

$$\frac{n!}{z_\mu} \xi^{f_1}(p_\mu) = \sum_{\sigma \in S_n(\mu)} x^{exc(\sigma)}$$


Proof.

$$\begin{aligned}
 \frac{n!}{z_\mu} \xi^{f_1}(p_\mu) &= \frac{n!}{z_\mu} \sum_{\lambda \vdash n} (-1)^{n-\ell(\lambda)} w(B_{\lambda, \mu}) \xi(e_\lambda) \\
 &= \frac{n!}{z_\mu} \sum_{\lambda \vdash n} (-1)^{n-\ell(\lambda)} w(B_{\lambda, \mu}) \prod_{i=1}^{\ell(\lambda)} \frac{(-1)^{\lambda_i-1}}{\lambda_i!} (x-1)^{\lambda_i-1} \\
 &= \frac{1}{z_\mu} \sum_{\lambda \vdash n} \binom{n}{\lambda_1, \dots, \lambda_\ell} w(B_{\lambda, \mu}) \prod_{i=1}^{\ell(\mu)} (x-1)^{\lambda_i-1}.
 \end{aligned}$$

We have

$$\frac{n!}{z_\mu} \xi^{f_1}(p_\mu) = \frac{1}{z_\mu} \sum_{\lambda \vdash n} \binom{n}{\lambda_1, \dots, \lambda_\ell} w(B_{\lambda, \mu}) \prod_{i=1}^{\ell} (x - 1)^{\lambda_i - 1}$$

from which we create the following objects:



We have

$$\frac{n!}{z_\mu} \xi^{f_1}(p_\mu) = \frac{1}{z_\mu} \sum_{\lambda \vdash n} \binom{n}{\lambda_1, \dots, \lambda_\ell} w(B_{\lambda, \mu}) \prod_{i=1}^{\ell} (x - 1)^{\lambda_i - 1}$$

from which we create the following objects:

18	*	5	14		
2		15	*	17	
1	10	13	*	7	8
16	3	*	4	9	11
					12

We make one more transformation.

18	*	5	14			
2	15	*	17			
1	10	13	*	6	7	8
16	3	*	4	9	11	12



*	5	14	18			
2	15	*	17			
1	6	10	7	8	13	
*	4	3	9	11	12	16

We have

$$\frac{n!}{z_\mu} \xi^{f_1}(p_\mu) = \frac{1}{z_\mu} \sum_{\lambda \vdash n} \binom{n}{\lambda_1, \dots, \lambda_\ell} w(B_{\lambda, \mu}) \prod_{i=1}^{\ell} (x-1)^{\lambda_i-1}$$

from which we create the following objects:

*						
5	14	18				
-1	x	-1				
			*			
2	15	17				
-1	-1	-1				
	*					
1	6	10	7	8	13	
-1	x	-1	x	x	-1	
*						
4	3	9	11	12	16	
-1	x	x	-1	x	-1	

We have

$$\frac{n!}{z_\mu} \xi^{f_1}(p_\mu) = \frac{1}{z_\mu} \sum_{\lambda \vdash n} \binom{n}{\lambda_1, \dots, \lambda_\ell} w(B_{\lambda, \mu}) \prod_{i=1}^{\ell} (x - 1)^{\lambda_i - 1}$$

from which we create the following objects:

*						
5	14	18				
1	x	1				
			*			
2	15	17				
-1	1	1				
	*					
1	6	10	7	8	13	
-1	x	1	x	x	1	
*						
4	3	9	11	12	16	
1	x	x	-1	x	1	

Let \mathcal{T}_{f_1} be the set of objects created in this way.

The weight of $T \in \mathcal{T}_{f_1}$, $w(T)$, is the product of x , -1 , and 1 labels.

$$\frac{n!}{z_\mu} \xi^{f_1}(p_\mu) = \frac{1}{z_\mu} \sum_{T \in \mathcal{T}_{f_1}} w(T).$$

An involution will rid us of all $T \in \mathcal{T}_{f_1}$ with negative weights.

Scan in rows from left to right and in rows from bottom to top for a “−1” or two consecutive bricks with a increase between them:

If a -1 is found, break the brick in two and change -1 to 1 .

If a increase between two bricks is found, combine the bricks and change 1 to -1 .

*						
5	14	18				
1	x	1				
			*			
2	15	17				
-1	1	1				
	*					
1	6	10	7	8	13	
-1	x	1	x	x	1	
*						
4	3	9	11	12	16	
1	x	x	-1	x	1	



*						
5	14	18				
1	x	1				
			*			
2	15	17				
-1	1	1				
	*					
1	6	10	7	8	13	
-1	x	1	x	x	1	
*						
4	3	9	11	12	16	
1	x	x	1	x	1	

A fixed point under this involution:

*						
14	5	18				
1	x	1				
			*			
2	15	17				
x	x	1				
	*					
1	6	10	7	8	13	
x	x	1	x	x	1	
*						
4	3	9	11	12	16	
1	x	x	x	x	1	

A fixed point can be read as an element σ in $S_n(\lambda)$:

$$(14, 5, 18) \quad (2, 15, 17) \quad (1, 6, 10, 7, 8, 13) \quad (4, 3, 9, 11, 12, 16)$$

If $\mu = (1^{\alpha_1} \cdots n^{\alpha_n})$, then each such permutation gives rise to $1^{\alpha_1} \cdots n^{\alpha_n} \alpha_1! \cdots \alpha_n!$ configurations in \mathcal{T}_{f_1} . That is, there is $1^{\alpha_1} \cdots n^{\alpha_n}$ ways to place the *s and $\alpha_1! \cdots \alpha_n!$ ways to order the rows of the same size. In our example, σ gives rise to $3^2 6^2 (2!) (2!)$ depending on where we place the *'s and how we order the cycles of the same size.

Therefore,

$$\frac{n!}{z_\mu} \xi^{f_1}(p_\mu) = \frac{1}{z_\mu} \sum_{T \in \mathcal{T}_{f_1}} w(T) = \frac{1}{z_\mu} \sum_{\substack{T \text{ is a} \\ \text{fixed point}}} w(T) = \sum_{\sigma \in S_n(\mu)} x^{\text{exc}(\sigma)}.$$

□

B_n is the set of $\sigma \in S_n$ where + or - is placed on each integer. Let $pos(\sigma)$ and $neg(\sigma)$ count the +'s and -'s.

D_n is the subset of B_n where $neg(\sigma)$ is even.

Descents and rises in B_n use the linear order Θ :

$$1 <_{\Theta} \cdots <_{\Theta} n <_{\Theta} -n <_{\Theta} \cdots <_{\Theta} -1.$$

Ex. Let $\sigma = -3 \text{ } -2 \text{ } -6 \text{ } +5 \text{ } -1 \text{ } +4$

Then $pos(\sigma) = 2$, $neg(\sigma) = 4$, $des_B(\sigma) = 3$, and $ris_B(\sigma) = 3$.

Let

$$f_3(n) = \begin{cases} 1 & \text{if } n = 0 \\ u^n y(x-y)^{n-1} + v^n x(y-x)^{n-1} & \text{if } n \geq 1. \end{cases}$$

and $\xi^{f_3}(e_n) = \frac{(-1)^{n-1}}{n!} f_3(n)$ as usual.

Theorem.

$$n! \xi^{f_3}(h_n) = \sum_{\sigma \in B_n} u^{pos(\sigma)} v^{neg(\sigma)} x^{des_B(\sigma)} y^{ris_B(\sigma)}.$$

Proof. We have,

$$n! \xi^{f_3}(h_n) = \sum_{\lambda \vdash n} \binom{n}{\lambda_1, \dots, \lambda_\ell} B_{\lambda, n} (-1)^{\ell(\lambda)} f_3(\lambda_1) \cdots f_3(\lambda_\ell)$$

from which we create the following objects:

$$f_3(n) = u^n y (x - y)^{n-1} + v^n x (y - x)^{n-1}$$

$$n! \xi^{f_3}(h_n) = \sum_{\lambda \vdash n} \binom{n}{\lambda_1, \dots, \lambda_\ell} B_{\lambda, n} (-1)^{\ell(\lambda)} f_3(\lambda_1) \cdots f_3(\lambda_\ell)$$

from which we create the following objects:

$$f_3(n) = u^n y(x-y)^{n-1} + v^n x(y-x)^{n-1}$$

$$n! \xi^{f_3}(h_n) = \sum_{\lambda \vdash n} \binom{n}{\lambda_1, \dots, \lambda_\ell} B_{\lambda, n} f_3(\lambda_1) \cdots f_3(\lambda_\ell)$$

from which we create the following objects:

11	9	8	4	3	12	10	7	5	1	6	2

$$f_3(n) = u^n y (x - y)^{n-1} + v^n x (y - x)^{n-1}$$

$$n! \xi^{f_3}(h_n) = \sum_{\lambda \vdash n} \binom{n}{\lambda_1, \dots, \lambda_\ell} B_{\lambda, n} f_3(\lambda_1) \cdots f_3(\lambda_\ell)$$

from which we create the following objects:

v	v	v	u	u	u	u	u	u	v	v
y	y	x	x	y	x	$-y$	$-y$	x	y	x
11	9	8	4	3	12	10	7	5	1	6

$$f_3(n) = u^n y (x - y)^{n-1} + v^n x (y - x)^{n-1}$$

Scan for the first instance of a “–” sign or two bricks with a decrease and the same u or v label. Break or combine accordingly.

In this way,

v	v	v	u	u	u	u	u	u	v	v
y	y	x	x	y	x	$-y$	$-y$	x	y	y
11	9	8	4	3	12	10	7	5	1	6

is sent to

v	v	v	u	u	u	u	u	u	v	v
y	y	x	x	y	x	y	$-y$	x	y	y
11	9	8	4	3	12	10	7	5	1	6

A fixed point:

v	v	v	u	u	u	u	u	u	v	v
y	y	x	x	y	x	x	x	x	y	x
11	9	8	4	3	12	10	7	5	1	6

Interpret every u as positive and v as negative. The above fixed point corresponds to

$$-11 \ -9 \ -8 \ +4 \ +3 \ +12 \ +10 \ +7 \ +5 \ +1 \ -6 \ -2$$

Summing over all fixed points completes the proof. \square

The generating function:

$$\sum_{n \geq 0} \frac{t^n}{n!} \sum_{\sigma \in B_n} u^{pos(\sigma)} v^{neg(\sigma)} x^{des_B(\sigma)} y^{ris_B(\sigma)} = \frac{x - y}{xe^{tv(y-x)} - ye^{tu(x-y)}}.$$

To find a generating function for D_n , notice that for $\sigma \in B_n$

$$\frac{v^{neg(\sigma)} + (-v)^{neg(\sigma)}}{2} = \begin{cases} v^{neg(\sigma)} & \text{if } \sigma \in D_n, \\ 0 & \text{if } \sigma \notin D_n. \end{cases}$$

Therefore,

$$\begin{aligned} & \sum_{n \geq 0} \frac{t^n}{n!} \sum_{\sigma \in D_n} u^{pos(\sigma)} v^{neg(\sigma)} x^{des_B(\sigma)} y^{ris_B(\sigma)} \\ &= \frac{x - y}{2xe^{tv(y-x)} - 2ye^{tu(x-y)}} + \frac{x - y}{2xe^{tv(x-y)} - 2ye^{tu(x-y)}}. \end{aligned}$$

For $\sigma^1, \dots, \sigma^m \in S_n$,

comdes($\sigma^1, \dots, \sigma^m$) is the number of times $\sigma_i^j > \sigma_{i+1}^j$ for all j

oneris($\sigma^1, \dots, \sigma^m$) is the number of times $\sigma_i^j < \sigma_{i+1}^j$ for at least one j

$$\sigma^1 = 12 \ 10 \ 6 \ 4 \ 3 \ 9 \ 2 \ 11 \ 1 \ 8 \ 5 \ 7$$

Ex. If $\sigma^2 = 10 \ 5 \ 11 \ 2 \ 6 \ 1 \ 7 \ 4 \ 12 \ 9 \ 8 \ 3$, then

$$\sigma^3 = 6 \ 5 \ 12 \ 1 \ 4 \ 9 \ 10 \ 3 \ 2 \ 11 \ 8 \ 7$$

comdes($\sigma^1, \sigma^2, \sigma^3$) = 3 and *oneris*($\sigma^1, \sigma^2, \sigma^3$) = 9.

Let $m \geq 1$ and $g_1(n) = \frac{1}{(n!)^m}$ such that

$$\xi^{f_4}(e_n) = \begin{cases} 0 & \text{if } n = 0 \\ \frac{(-1)^{n-1}}{(n!)^m} y(x-y)^{n-1} & \text{if } n \geq 1 \end{cases}$$

Theorem.

$$(n!)^m \xi^{f_4}(h_n) = \sum_{\sigma \in S_n^m} x^{\text{comdes}(\sigma)} y^{\text{oneris}(\sigma)}.$$

Proof.

$$(n!)^m \xi^{f_4}(h_n)$$

$$\begin{aligned} &= (n!)^m \sum_{\lambda \vdash n} (-1)^{n-\ell(\lambda)} B_{\lambda,n} \xi^{f_4}(e_\lambda) \\ &= (n!)^m \sum_{\lambda \vdash n} (-1)^{n-\ell(\lambda)} B_{\lambda,n} \prod_{i=1}^{\ell(\lambda)} \frac{(-1)^{\lambda_i-1}}{(\lambda_i!)^m} y(x-y)^{\lambda_i-1} \\ &= \sum_{\lambda \vdash n} \binom{n}{\lambda_1, \dots, \lambda_\ell}^m B_{\lambda,n} \prod_{i=1}^{\ell(\lambda)} y(x-y)^{\lambda_i-1}. \end{aligned}$$

We have

$$(n!)^m \xi^{f_4}(h_n) = \sum_{\lambda \vdash n} \binom{n}{\lambda_1, \dots, \lambda_\ell}^m B_{\lambda, n} \prod_{i=1}^{\ell(\lambda)} y(x - y)^{\lambda_i - 1}.$$

which may be used to create:

We have

$$(n!)^m \xi^{f_4}(h_n) = \sum_{\lambda \vdash n} \binom{n}{\lambda_1, \dots, \lambda_\ell}^m B_{\lambda, n} \prod_{i=1}^{\ell(\lambda)} y(x - y)^{\lambda_i - 1}.$$

which may be used to create:

We have

$$(n!)^m \xi^{f_4}(h_n) = \sum_{\lambda \vdash n} \binom{n}{\lambda_1, \dots, \lambda_\ell}^m B_{\lambda, n} \prod_{i=1}^{\ell(\lambda)} y(x - y)^{\lambda_i - 1}.$$

which may be used to create:

11	9	3	12	2	10	6	5	4	1	8	7
12	8	5	11	10	9	7	6	4	3	2	1

We have

$$(n!)^m \xi^{f_4}(h_n) = \sum_{\lambda \vdash n} \binom{n}{\lambda_1, \dots, \lambda_\ell}^m B_{\lambda, n} \prod_{i=1}^{\ell(\lambda)} y(x-y)^{\lambda_i-1}.$$

which may be used to create:

x	x	y	x	y	$-y$	x	$-y$	$-y$	y	x	y
11	9	3	12	2	10	6	5	4	1	8	7
12	8	5	11	10	9	7	6	4	3	2	1

Apply the brick breaking/combining involution where bricks are combined if two bricks have a common descent between them. Then

x	x	$-y$	x	y	$-y$	x	$-y$	$-y$	y	x	y
12	11	9	3	2	10	6	5	4	1	8	7
12	11	10	8	5	9	7	6	4	3	2	1

is sent to

x	x	y	x	y	$-y$	x	$-y$	$-y$	y	x	y
12	11	9	3	2	10	6	5	4	1	8	7
12	11	10	8	5	9	7	6	4	3	2	1

A fixed point:

x	x	y	x	y	x	x	x	x	y	x	y
11	9	3	12	2	10	6	5	4	1	8	7
12	8	5	11	10	9	7	6	4	3	2	1

This fixed point corresponds to:

$$\sigma^1 = 11 \text{ } 9 \text{ } 3 \text{ } 12 \text{ } 2 \text{ } 10 \text{ } 6 \text{ } 5 \text{ } 4 \text{ } 1 \text{ } 8 \text{ } 7$$

$$\sigma^2 = 12 \text{ } 8 \text{ } 5 \text{ } 11 \text{ } 10 \text{ } 9 \text{ } 7 \text{ } 6 \text{ } 4 \text{ } 3 \text{ } 2 \text{ } 1$$

Summing all fixed points gives the desired result. \square

The generating function which follows:

$$\begin{aligned}
 \sum_{n \geq 0} \frac{t^n}{(n!)^m} \sum_{\sigma \in S_n^m} x^{\text{comdes}(\sigma)} y^{\text{oneris}(\sigma)} &= \left(1 + \sum_{n \geq 1} (-t)^n \frac{(-1)^{n-1}}{(n!)^m} y(x-y)^{n-1} \right)^{-1} \\
 &= \frac{x-y}{x-y \sum_{n \geq 0} \frac{(x-y)^n t^n}{(n!)^m}}.
 \end{aligned}$$

For $\sigma_1 \cdots \sigma_n \in S_n$,

$inv(\sigma)$ is the number of times $\sigma_i > \sigma_j$ for $i < j$

$coinv(\sigma)$ is the number of times $\sigma_i < \sigma_j$ for $i < j$

Let $[n]_{p,q} = \frac{p^n - q^n}{p - q}$ and $[n]_{p,q}! = [n]_{p,q} \cdots [2]_{p,q} [1]_{p,q}$.

Let

$$\left[\begin{matrix} n \\ \lambda_1, \dots, \lambda_\ell \end{matrix} \right]_{p,q} = \frac{[n]_{p,q}!}{[\lambda_1]_{p,q}! \cdots [\lambda_\ell]_{p,q}!}$$

Let $\mathcal{R}(1^{\lambda_1}, \dots, \ell^{\lambda_\ell})$ be the set of rearrangements of λ_1 1's, λ_2 2's, ...

A generalization of a theorem of Carlitz:

$$\begin{bmatrix} n \\ \lambda_1, \dots, \lambda_\ell \end{bmatrix}_{p,q} = \sum_{r \in \mathcal{R}(1^{\lambda_1}, \dots, \ell^{\lambda_\ell})} q^{inv(r)} p^{coinv(r)}.$$

Given a rearrangement r of b_1 1s, b_2 2s, \dots , b_l l 's, form a permutation $\sigma(r)$ by labelling the 1s in r from right to left with 1, 2, \dots , b_1 . Then number the 2s from right to left with $b_1 + 1, b_1 + 2, \dots, b_1 + b_2$, and so on.

For example, let $b_1 = 4, b_2 = 2, b_3 = 3$ and

$$\begin{aligned} r &= 1 \ 3 \ \textcolor{blue}{2} \ 1 \ 3 \ 3 \ \textcolor{blue}{1} \ \textcolor{red}{2} \ 1 \ 3 \ 3 \\ \sigma(r) &= 4 \ 11 \ \textcolor{red}{6} \ \textcolor{blue}{3} \ 10 \ 9 \ \textcolor{blue}{2} \ \textcolor{red}{5} \ 1 \ 8 \ 7 \end{aligned}$$

$$\sigma(r)^{-1} = \textcolor{blue}{9} \ \textcolor{red}{7} \ \textcolor{blue}{4} \ 1 \ \textcolor{red}{8} \ \textcolor{blue}{3} \ 11 \ 10 \ 6 \ 5 \ 2$$

$\text{inv}(\sigma(r)) = \text{inv}(r) +$ the number of inversions introduced by changing the 1s in r to $1, 2, \dots, b_1$, the 2s to $b_1 + 1, b_1 + 2, \dots, b_1 + b_2$, and so on.

The number of inversions introduced by changing the 1s to $1, 2, \dots, b_1$ is $(b_1 - 1) + (b_1 - 2) + \dots + 1 = \binom{b_1}{2}$.

$$\begin{aligned}\text{inv}(\sigma(r)) &= \text{inv}(r) + \sum_k \binom{b_k}{2} \\ \text{coinv}(\sigma(r)) &= \text{coinv}(r)\end{aligned}$$

Theorem 0.1. *If $b_1 + b_2 + \cdots + b_l = n$ where each b_i is a positive integer, then*

$$\left[\begin{array}{c} n \\ b_1, b_2, \dots, b_l \end{array} \right]_{p,q} q^{\sum_i \binom{b_i}{2}} = \sum_{\tau \in \text{dec}_n(b_1, \dots, b_l)} q^{\text{inv}(\tau)} p^{\text{coinv}(\tau)}$$

where $\text{dec}_n(b_1, \dots, b_l)$ is the set of permutations in S_n that when read from left to right in one-line notation consist of decreasing sequences in blocks of size b_1, b_2, \dots, b_l .

$$f_1(n) = \begin{cases} 1 & \text{if } n = 0 \\ y(x-y)^{n-1} & \text{if } n \geq 1. \end{cases}$$

Define $\xi^{f_1} : \Lambda \rightarrow \mathbb{Q}[x, y]$ as a homomorphism such that

$$\xi_{p,q}^{f_1}(e_n) = \frac{(-1)^{n-1} q^{\binom{n}{2}}}{[n]_{p,q}!} f_1(n).$$

Theorem.

$$[n]_{p,q}! \xi_{p,q}^{f_1}(h_n) = \sum_{\sigma \in S_n} x^{des(\sigma)} y^{ris(\sigma)} q^{inv(\sigma)} p^{coinv(\sigma)}.$$

Proof.

$$[n]_{p,q}! \xi_{p,q}^{f_1}(h_n)$$

$$\begin{aligned} &= [n]_{p,q}! \sum_{\lambda \vdash n} (-1)^{n-\ell(\lambda)} B_{\lambda,n} \xi_{p,q}^{f_1}(e_\lambda) \\ &= \sum_{\lambda \vdash n} \left[\begin{matrix} n \\ \lambda_1, \dots, \lambda_\ell \end{matrix} \right]_{p,q} q^{\binom{\lambda_1}{2} + \dots + \binom{\lambda_\ell}{2}} B_{\lambda,n} (-1)^{\ell(\lambda)-1} f_1(\lambda_1) \cdots f_1(\lambda_\ell). \end{aligned}$$

$$f_1(n) = y(x-y)^{n-1}$$

$$\sum_{\lambda \vdash n} \left[\begin{matrix} n \\ \lambda_1, \dots, \lambda_\ell \end{matrix} \right]_{p,q} q^{\binom{\lambda_1}{2} + \dots + \binom{\lambda_\ell}{2}} B_{\lambda, n} f_1(\lambda_1) \cdots f_1(\lambda_\ell).$$

which may be used to create:

$$f_1(n) = y(x - y)^{n-1}$$

The term we have to work with is

$$\sum_{\lambda \vdash n} \begin{bmatrix} n \\ \lambda_1, \dots, \lambda_\ell \end{bmatrix}_{p,q} q^{\binom{\lambda_1}{2} + \dots + \binom{\lambda_\ell}{2}} \color{red} B_{\lambda, n} f_1(\lambda_1) \cdots f_1(\lambda_\ell).$$

which may be used to create:

⋮ ⋮ ⋮ ⋮ ⋮ ⋮ ⋮ ⋮ ⋮ ⋮ ⋮ ⋮ ⋮ ⋮ ⋮ ⋮ ⋮ ⋮	⋮ ⋮ ⋮ ⋮ ⋮ ⋮	⋮ ⋮ ⋮ ⋮ ⋮ ⋮ ⋮ ⋮ ⋮ ⋮ ⋮ ⋮	⋮ ⋮ ⋮ ⋮ ⋮ ⋮ ⋮ ⋮ ⋮ ⋮ ⋮ ⋮
--	----------------------------	--	--

$$f_1(n) = y(x - y)^{n-1}$$

$$\sum_{\lambda \vdash n} \begin{bmatrix} n \\ \lambda_1, \dots, \lambda_\ell \end{bmatrix}_{p,q} q^{\binom{\lambda_1}{2} + \dots + \binom{\lambda_\ell}{2}} B_{\lambda,n} f_1(\lambda_1) \cdots f_1(\lambda_\ell).$$

which may be used to create:

q^{11}	q^7	q^6	q^4	q^6	q^5	q^5	q^4	q^2	q^2	q^1	q^0
p^0	p^3	p^3	p^4	p^1	p^1	p^0	p^0	p^1	p^0	p^0	p^0
12	8	7	5	10	9	11	6	3	4	2	1

$$f_1(n) = y(x - y)^{n-1}$$

The term we have to work with is

$$\sum_{\lambda \vdash n} \left[\begin{matrix} n \\ \lambda_1, \dots, \lambda_\ell \end{matrix} \right]_{p,q} q^{\binom{\lambda_1}{2} + \dots + \binom{\lambda_\ell}{2}} B_{\lambda, n} f_1(\lambda_1) \cdots f_1(\lambda_\ell).$$

which may be used to create:

x	x	$-y$	y	x	y	x	$-y$	y	$-y$	$-y$	y
q^{11}	q^7	q^6	q^4	q^6	q^5	q^5	q^4	q^2	q^2	q^1	q^0
p^0	p^3	p^3	p^4	p^1	p^1	p^0	p^0	p^1	p^0	p^0	p^0
12	8	7	5	10	9	11	6	3	4	2	1

$$f_1(n) = y(x - y)^{n-1}$$

Apply the brick breaking/combining involution so that

x	x	$-y$	y	x	y	x	$-y$	y	$-y$	$-y$	y
q^{11}	q^7	q^6	q^4	q^6	q^5	q^5	q^4	q^2	q^2	q^1	q^0
p^0	p^3	p^3	p^4	p^1	p^1	p^0	p^0	p^1	p^0	p^0	p^0
12	8	7	5	10	9	11	6	3	4	2	1

is sent to

x	x	y	y	x	y	x	$-y$	y	$-y$	$-y$	y
q^{11}	q^7	q^6	q^4	q^6	q^5	q^5	q^4	q^2	q^2	q^1	q^0
p^0	p^3	p^3	p^4	p^1	p^1	p^0	p^0	p^1	p^0	p^0	p^0
12	8	7	5	10	9	11	6	3	4	2	1

A fixed point under this involution:

x	x	x	y	x	y	x	x	y	x	x	y
q^{11}	q^7	q^6	q^4	q^6	q^5	q^5	q^4	q^2	q^2	q^1	q^0
p^0	p^3	p^3	p^4	p^1	p^1	p^0	p^0	p^1	p^0	p^0	p^0
12	8	7	5	10	9	11	6	3	4	2	1

The powers of x, y, q and p on a fixed point register the correct statistics.

Summing all fixed points completes the proof. \square

The generating function which follows:

$$\sum_{n \geq 0} \frac{t^n}{[n]!} \sum_{\sigma \in S_n} x^{des(\sigma)} y^{ris(\sigma)} q^{inv(\sigma)} p^{coinv(\sigma)} = \frac{x - y}{x - y \exp_{q,p}^{t(x-y)}}$$

where for a series $r(t) = \sum_{n \geq 0} a_n \frac{t^n}{n!}$, we let $r_{q,p}(t) = \sum_{n \geq 0} a_n \frac{t^n}{[n]_{p,q}!} q^{\binom{n}{2}}$.

Replacing $\frac{(-1)^{n-1}}{n!}$ by $\frac{(-1)^{n-1}q^{\binom{n}{2}}}{[n]_{p,q}!}$ in our other proofs will allow us to prove the following.

$$\sum_{n \geq 0} \frac{t^n}{[n]_{p,q}!} \sum_{\substack{\sigma \in S_n \\ \sigma \text{ is even alt.}}} q^{inv(\sigma)} p^{coinv(\sigma)} = \frac{1}{\cos_{q,p}(t)}$$

and

$$\begin{aligned} \sum_{n \geq 0} \frac{t^n}{[n]_{p,q}!} \sum_{\sigma \in B_n} u^{pos(\sigma)} v^{neg(\sigma)} x^{des_B(\sigma)} y^{ris_B(\sigma)} q^{inv(\sigma)} p^{coinv(\sigma)} \\ = \frac{x - y}{xe_{q,p}^{tv(y-x)} - ye_{q,p}^{tu(x-y)}}. \end{aligned}$$