# FIRST ORDER DIFFERENTIAL EQUATIONS

1. A first order differential equation is an equation of the form

$$F(x, y, y') = 0.$$

A solution of the differential equation is a function y = y(x) that satisfies the equation. A differential equation has **infinitely many** solutions.

- 2. An equilibrium solution is a constant solution, i.e. a constant function y(x) = K that satisfies the equation.
- 3. An initial value problem consists of a differential equation and an initial condition  $y(x_0) = y_0$ . It has a **unique** solution, which has to be a function. For instance  $y = \pm \sqrt{x}$  cannot be the solution of an i.v.p.

## A. Separable Equations

$$\frac{dy}{dx} = f(x)g(y)$$

- 1. Separate the variables:  $\frac{dy}{g(y)} = f(x)dx$ .
- 2. Integrate both sides:  $\int \frac{dy}{g(y)} = \int f(x)dx$  and get G(y) = C + F(x).
- 3. Solve for y (if possible).
- 4. If it is an initial value problem, plug in the given values and solve for C.

#### B. Linear equations

$$\frac{dy}{dx} + P(x)y = Q(x)$$

- 1. Write the equation in the above form.
- 2. Compute the integrating factor  $I(x) = e^{\int P(x)dx}$ .
- 3. Multiply both sides of the equation by this integrating factor:  $I(x)\frac{dy}{dx} + I(x) \cdot P(x)y = I(x) \cdot Q(x)$ .
- 4. Integrate both sides and get  $I(x)y = C + \int I(x)Q(x)dx$ .
- 5. Solve for y.
- 6. If it is an initial value problem, plug in the given values and solve for C.

#### C. Homogeneous Equations

$$\frac{dy}{dx} = F\left(\frac{y}{x}\right)$$

1. Make the substitution  $v = \frac{y}{x}$ . Attention:  $\frac{dy}{dx} = x \cdot \frac{dv}{dx} + v$ .

- 2. Solve the new equation, not forgetting the constant involved. (It will probably be a separable equation.)
- 3. Find y(x) = xv(x).
- 4. If it is an initial value problem, make sure you find the constant.

#### D. Bernoulli Equations

$$\frac{dy}{dx} + P(x)y = Q(x)y^n$$

- 1. Make the substitution  $v = y^{1-n}$ .
- 2. Solve the new linear equation:  $\frac{dv}{dx} + (1-n)P(x)y = (1-n)Q(x)$ .
- 3. Find y(x).
- 4. If it is an initial value problem, make sure you find the constant.

#### E. Other Substitutions

Some equations need some other substitution to transform them in a known type. These substitutions vary greatly and there are no general formulas that can help. However, the more you practice, the better your intuition becomes in these matters.

### F. Linear first order equations with sinusoidal input

$$y' + ky = B\cos(\omega t) \text{ or } B\sin(\omega t)$$
 (\*)

Such an equation can be seen as the real or imaginary part of the complex differential equation

$$z' + kz = Be^{i\omega t} \tag{**}$$

The general idea is that the solutions to (\*) are of the form

$$y = y_p + y_h,$$

where  $y_p$  is a particular solution of the original equation (\*), and  $y_h$  is the general solution to the associated homogeneous equation (see below).

- 1. Solve the corresponding homogeneous equation y' + ky = 0. The solution is  $y_h = Ce^{-kt}$ . This is where the arbitrary constant appears!
- 2. Find a particular solution  $z_p$  of (\*\*). It will usually be of the same form as the input, namely  $Ae^{i\omega t}$ . So try that and determine A.
- 3. Write  $y_p$  by taking either the real or imaginary part of  $z_p$ . If the input in (\*) is cos, you should take the real part, if it is sin, the imaginary part.
- 4. Write down the general solution to (\*),  $y = y_p + y_h$
- 5. If it is an initial value problem, find the constant.

### G. General strategy

If you are asked to solve a first order ODE and no method is specified, go through the following steps in order to determine what method to apply.

- 1. Is it separable?
- 2. Is it linear? Does it have sinusoidal or exponential input?
- 3. Is it homogeneous? Reducible?
- 4. Is it a Bernoulli equation?
- 5. Can you think of a substitution that would put it in a form you recognize?