

Math 20C. Lecture Examples.

Sections 15.1 and 15.2. Double integrals[†]

Example 1 What is the value of $\iint_R \sqrt{9 - x^2 - y^2} \, dx \, dy$ where R is the disk $x^2 + y^2 \leq 9$?

Answer: $\iint_R \sqrt{9 - x^2 - y^2} \, dx \, dy = [\text{Volume of the hemisphere in Figure A1}] = 18\pi$




Figure A1

Example 2 Evaluate $\iint_R 2xy \, dx \, dy$, where R is the region bounded by the curve $y = x^2$ and the lines $y = 0$ and $x = 2$ by performing a y -integration first.

Answer: Figure A2 • $\iint_R 2xy \, dx \, dy = \frac{32}{3}$




Figure A2

Example 3 Evaluate the integral of Example 2 by performing an x -integration first.

Answer: Figure A3 • $\iint_R 2xy \, dx \, dy = \frac{32}{3}$




Figure A3

[†]Lecture notes to accompany Sections 15.1 and 15.2 of *Calculus, Early Transcendentals* by Rogawski.

Example 4 Evaluate $\int_0^8 \int_{x^{1/3}}^2 \sin(y^4) dy dx$ by reversing the order of integration.

Answer: Figures A4a and A4b • $\int_0^8 \int_{x^{1/3}}^2 \sin(y^4) dy dx = \int_0^2 \int_0^{y^3} \sin(y^4) dx dy = \frac{1}{4} - \frac{1}{4} \cos(16)$




Figure A4a




Figure A4b

Example 5 What is the volume of the solid V bounded by the surface $z = \sin x \sin y$ and the plane $z = -2$ for $0 \leq x \leq \pi, 0 \leq y \leq \pi$?

Answer: The projection of the solid on the xy -plane is the square in Figure A5. • [Volume] = $4 + 2\pi^2$




Figure A5

Example 6 A plate that occupies the region $R : 0 \leq x \leq 2, 0 \leq y \leq x^2$ with distances measured in feet has density $\rho(x, y) = 3xy^2$ pounds per square foot at (x, y) . How much does the plate weigh?

Answer: Figure A6 • [Weight] = $\iint_R 3xy^2 dx dy = 32$ pounds





Figure A6


Example 7 Figure 1 shows level curves of the population density $z = p(x, y)$ (people per square mile) in a city. Find the approximate population in the region $R: 0 \leq x \leq 4, 0 \leq y \leq 4$. Use a Riemann sum corresponding to a partition into four squares.

Level curves of $z = p(x, y)$
FIGURE 1



Answer: [Population] = $\iint_R p(x, y) \, dy \, dx$ • One approach: Approximate the integral with the Riemann sum corresponding to the partition of R into four equal square subregions of Figure A7 with the population density evaluated at their midpoints. • The population is approximately 58,000.


Figure A7



Example 8 What is the average value of $f(x, y) = ye^{y^2}$ on the region $R: 0 \leq y \leq \sqrt{x}, 0 \leq x \leq 1$?

Answer: Figure A8 • [Area of R] = $\frac{2}{3}$ • $\iint_R ye^{y^2} \, dx \, dy = \frac{1}{2}(e - 2)$ • [Average value] = $\frac{3}{4}(e - 2)$

Figure A8



Interactive Examples

Work the following Interactive Examples on Shenk's web page, <http://www.math.ucsd.edu/~ashenk/>:[‡]

Section 16.1: Examples 1 through 5

Section 16.2: Examples 1, 2a, and 3

[‡]The chapter and section numbers on Shenk's web site refer to his calculus manuscript and not to the chapters and sections of the textbook for the course.