

Math 240C Homework Problem List for S2018

0.1 Homework C1. Due Friday, April 6, 2018

- **Hand in:** 1.3, 1.4, 1.5, 1.25, 1.26
- **Look at:** 1.23

0.2 Homework C2. Due Friday, April 13, 2018 (L^p inequalities)

- **Hand in:** 1.1, 1.2, 1.6, 1.7, 1.19, 1.20
- **Look at:** 1.21, 1.22

0.3 Homework C3. Due Friday, April 20, 2018 (Fourier Series problems)

- **Hand in:** 1.8, 1.10, 1.11, 1.12, 1.13, 1.14, 1.16
- **Look at:** 1.24 (done in class), 1.9, 1.15

0.4 Homework C4. Due Friday, April 27, 2018 (Fourier Transform problems)

- **Hand in:** 1.27, 1.29, 1.30, 1.31, 1.32, 1.33
- **Look at:** 1.28

Problems to Solve

Exercise 1.1. If (X, ρ) is a metric space and μ is a **finite** measure on (X, \mathcal{B}_X) , then for all $A \in \mathcal{B}_X$ and $\varepsilon > 0$ there exists a closed set F and open set V such that $F \subset A \subset V$ and $\mu(V \setminus F) = \mu(F \triangle V) < \varepsilon$.

You may find information in the supplement helpful for this problem. Here are some more suggestions.

1. Let \mathcal{B}_0 denote those $A \subset X$ such that for all $\varepsilon > 0$ there exists a closed set F and open set V such that $F \subset A \subset V$ and $d_\mu(F, V) = \mu(V \setminus F) < \varepsilon$.
2. Show \mathcal{B}_0 contains all closed (or open if you like).
3. Show \mathcal{B}_0 is a σ -algebra.
4. Explain why this proves the result.

Exercise 1.2. Let (X, ρ) be a metric space and μ be a measure on (X, \mathcal{B}_X) . If there exists open sets, $\{V_n\}_{n=1}^\infty$, of X such that $V_n \uparrow X$ and $\mu(V_n) < \infty$ for all n , then for all $A \in \mathcal{B}_X$ and $\varepsilon > 0$ there exists a closed set F and open set V such that $F \subset A \subset V$ and $d_\mu(F, V) = \mu(V \setminus F) < \varepsilon$. **Hints:**

1. Show it suffices to prove; for all $\varepsilon > 0$ and $A \in \mathcal{B}_X$, there exists an open set $V \subset X$ such that $A \subset V$ and $\mu(V \setminus A) < \varepsilon$.
2. Now you must verify the assertion above holds. For this, you may find it useful to apply Exercise 1.1 to the measures, $\mu_n : \mathcal{B}_X \rightarrow [0, \mu(V_n)]$, defined by $\mu_n(A) := \mu(A \cap V_n)$ for all $A \in \mathcal{B}_X$. The ε in Exercise 1.1 should be replaced by judiciously chosen small quantities depending on n .

Exercise 1.3 (Folland Problem 2.62 on p. 80.). Rotation invariance of surface measure on S^{n-1} .

Exercise 1.4 (Folland Problem 2.64 on p. 80.). On the integrability of $|x|^\alpha |\log|x||^b$ for x near 0 and x near ∞ in \mathbb{R}^n .

Exercise 1.5. Show, using Problem 1.3 that

$$\int_{S^{d-1}} \omega_i \omega_j d\sigma(\omega) = \frac{1}{d} \delta_{ij} \sigma(S^{d-1}).$$

Hint: show $\int_{S^{d-1}} \omega_i^2 d\sigma(\omega)$ is independent of i and therefore

$$\int_{S^{d-1}} \omega_i^2 d\sigma(\omega) = \frac{1}{d} \sum_{j=1}^d \int_{S^{d-1}} \omega_j^2 d\sigma(\omega).$$

Exercise 1.6 (Folland 6.38 on p. 199.). Suppose (X, \mathcal{M}, μ) is a measure space, $f : X \rightarrow \mathbb{C}$ is a measurable function, $0 < p < \infty$, $\lambda_f(\alpha) := \mu(|f| > \alpha)$ for all $\alpha \in (0, \infty)$, and

$$M_p(f) := \sum_{k=-\infty}^{\infty} 2^{kp} \lambda_f(2^k).$$

Show

$$(1 - 2^{-p}) M_p(f) \leq \int_X |f|^p d\mu \leq 2^p M_p(f) \quad (1.1)$$

which then implies $f \in L^p(\mu)$ iff $M_p(f) < \infty$.

Hint: first note that

$$\int_X |f|^p d\mu = \sum_{k \in \mathbb{Z}} \int_{\{2^k < |f| \leq 2^{k+1}\}} |f|^p d\mu. \quad (1.2)$$

Exercise 1.7 (Folland 6.39 on p. 199.). Suppose (X, \mathcal{M}, μ) is a measure space, $f : X \rightarrow \mathbb{C}$ is a measurable function, $0 < p < \infty$, and $f \in L^p(\mu)$. Show

$$\lim_{\alpha \rightarrow 0} \alpha^p \lambda_f(\alpha) = 0 = \lim_{\alpha \rightarrow \infty} \alpha^p \lambda_f(\alpha).$$

Hint: for the limit, $\lim_{\alpha \rightarrow 0} \alpha^p \lambda_f(\alpha) = 0$, start with the special case where f is a simple function.

Exercise 1.8. Show $\sum_{k=1}^{\infty} k^{-2} = \pi^2/6$, by taking $f(x) = x$ on $[-\pi, \pi]$ and computing $\|f\|_2^2$ directly and then in terms of the Fourier Coefficients \tilde{f} of f .

Exercise 1.9 (Riemann Lebesgue Lemma for Fourier Series). Show for $f \in L^1([-\pi, \pi]^d)$ that $\tilde{f} \in c_0(\mathbb{Z}^d)$, i.e. $\tilde{f} : \mathbb{Z}^d \rightarrow \mathbb{C}$ and $\lim_{k \rightarrow \infty} \tilde{f}(k) = 0$. **Hint:** If $f \in L^2([-\pi, \pi]^d)$, this follows from Bessel's inequality. Now use a density argument.

Exercise 1.10. Suppose $f \in L^1([-\pi, \pi]^d)$ is a function such that $\tilde{f} \in \ell^1(\mathbb{Z}^d)$ and set

$$g(x) := \sum_{k \in \mathbb{Z}^d} \tilde{f}(k) e^{ik \cdot x} \text{ (pointwise).}$$

1. Show $g \in C_{per}(\mathbb{R}^d)$.
2. Show $g(x) = f(x)$ for m -a.e. x in $[-\pi, \pi]^d$. **Hint:** Show $\tilde{g}(k) = \tilde{f}(k)$ and apply Exercise ?? or results proved in class.
3. Conclude that $f \in L^1([-\pi, \pi]^d) \cap L^\infty([-\pi, \pi]^d)$ and in particular $f \in L^p([-\pi, \pi]^d)$ for all $p \in [1, \infty]$.

Exercise 1.11 (Smoothness implies decay). We use the following notation below.

Notation: Given a multi-index $\alpha \in \mathbb{Z}_+^d$, let $|\alpha| = \alpha_1 + \dots + \alpha_d$,

$$x^\alpha := \prod_{j=1}^d x_j^{\alpha_j}, \text{ and } \partial_x^\alpha = \left(\frac{\partial}{\partial x} \right)^\alpha := \prod_{j=1}^d \left(\frac{\partial}{\partial x_j} \right)^{\alpha_j}.$$

Further for $k \in \mathbb{N}_0$, let $f \in C_{per}^k(\mathbb{R}^d)$ iff $f \in C^k(\mathbb{R}^d) \cap C_{per}(\mathbb{R}^d)$, $\partial_x^\alpha f(x)$ exists and is continuous for $|\alpha| \leq k$.

Suppose $m \in \mathbb{N}_0$, α is a multi-index such that $|\alpha| \leq 2m$ and $f \in C_{per}^{2m}(\mathbb{R}^d)^1$.

1. Using integration by parts, show (using Notation ??) that

$$(ik)^\alpha \tilde{f}(k) = \langle \partial^\alpha f | \varphi_k \rangle \text{ for all } k \in \mathbb{Z}^d.$$

Note: This equality implies

$$|\tilde{f}(k)| \leq \frac{1}{k^\alpha} \|\partial^\alpha f\|_H \leq \frac{1}{k^\alpha} \|\partial^\alpha f\|_\infty.$$

2. Now let $\Delta f = \sum_{i=1}^d \partial^2 f / \partial x_i^2$, Working as in part 1) show

$$\langle (1 - \Delta)^m f | \varphi_k \rangle = (1 + \|k\|^2)^m \tilde{f}(k). \quad (1.3)$$

where $\|k\|^2 = \sum_{j=1}^d k_j^2$.

Exercise 1.12 (A Sobolev Imbedding Theorem). Suppose $s \in \mathbb{R}$ and $\{c_k \in \mathbb{C} : k \in \mathbb{Z}^d\}$ are coefficients such that

$$\sum_{k \in \mathbb{Z}^d} |c_k|^2 (1 + \|k\|^2)^s < \infty.$$

Show if $s > \frac{d}{2} + m$, the function f defined by

$$f(x) = \sum_{k \in \mathbb{Z}^d} c_k e^{ik \cdot x}$$

¹ We view $C_{per}(\mathbb{R}^d)$ as a subspace of $H = L^2([-\pi, \pi]^d)$ by identifying $f \in C_{per}(\mathbb{R}^d)$ with $f|_{[-\pi, \pi]^d} \in H$.

is in $C_{per}^m(\mathbb{R}^d)$. **Hint:** Work as in the above remark to show

$$\sum_{k \in \mathbb{Z}^d} |c_k| |k^\alpha| < \infty \text{ for all } |\alpha| \leq m.$$

Exercise 1.13 (Poisson Summation Formula). Let $F \in L^1(\mathbb{R}^d)$,

$$E := \left\{ x \in \mathbb{R}^d : \sum_{k \in \mathbb{Z}^d} |F(x + 2\pi k)| = \infty \right\}$$

and set (see Section ?? below for an introduction to the Fourier transform)

$$\hat{F}(k) := (2\pi)^{-d/2} \int_{\mathbb{R}^d} F(x) e^{-ik \cdot x} dx \text{ for } k \in \mathbb{Z}^d.$$

Further **assume** $\hat{F} \in \ell^1(\mathbb{Z}^d)$. [This can be achieved by assuming F is sufficiently differentiable with the derivatives being integrable like in Exercise 1.11.]

1. Show $m(E) = 0$ and $E + 2\pi k = E$ for all $k \in \mathbb{Z}^d$. **Hint:** Compute $\int_{[-\pi, \pi]^d} \sum_{k \in \mathbb{Z}^d} |F(x + 2\pi k)| dx$.
2. Let

$$f(x) := \begin{cases} \sum_{k \in \mathbb{Z}^d} F(x + 2\pi k) & \text{for } x \notin E \\ 0 & \text{if } x \in E. \end{cases}$$

Show $f \in L^1([-\pi, \pi]^d)$ and $\tilde{f}(k) = (2\pi)^{-d/2} \hat{F}(k)$.

3. Using item 2) and the assumptions on F , show

$$f(x) = \sum_{k \in \mathbb{Z}^d} \tilde{f}(k) e^{ik \cdot x} = \sum_{k \in \mathbb{Z}^d} (2\pi)^{-d/2} \hat{F}(k) e^{ik \cdot x} \text{ for } m\text{-a.e. } x,$$

i.e.

$$\sum_{k \in \mathbb{Z}^d} F(x + 2\pi k) = (2\pi)^{-d/2} \sum_{k \in \mathbb{Z}^d} \hat{F}(k) e^{ik \cdot x} \text{ for } m\text{-a.e. } x \quad (1.4)$$

and from this conclude that $f \in L^1([-\pi, \pi]^d) \cap L^\infty([-\pi, \pi]^d)$.

Hint: see the hint for item 2. of Exercise 1.10.

4. Suppose we now assume that $F \in C(\mathbb{R}^d)$ and F satisfies $|F(x)| \leq C(1 + |x|)^{-s}$ for some $s > d$ and $C < \infty$. Under these added assumptions on F , show Eq. (1.4) holds for **all** $x \in \mathbb{R}^d$ and in particular

$$\sum_{k \in \mathbb{Z}^d} F(2\pi k) = (2\pi)^{-d/2} \sum_{k \in \mathbb{Z}^d} \hat{F}(k).$$

Exercise 1.14 (Heat Equation 1). Let $[0, \infty) \times \mathbb{R} \ni (t, x) \rightarrow u(t, x)$ be a continuous function such that $u(t, \cdot) \in C_{per}(\mathbb{R})$ for all $t \geq 0$, $\dot{u} := u_t$, u_x , and u_{xx} exists and are continuous when $t > 0$. Further assume that u satisfies the heat equation $\dot{u} = \frac{1}{2}u_{xx}$. Let $\tilde{u}(t, k) := \langle u(t, \cdot) | \varphi_k \rangle$ for $k \in \mathbb{Z}$. Show for $t > 0$ and $k \in \mathbb{Z}$ that $\tilde{u}(t, k)$ is differentiable in t and $\frac{d}{dt} \tilde{u}(t, k) = -k^2 \tilde{u}(t, k)/2$. Use this result to show

$$u(t, x) = \sum_{k \in \mathbb{Z}} e^{-\frac{t}{2}k^2} \tilde{f}(k) e^{ikx} \tag{1.5}$$

where $f(x) := u(0, x)$ and as above

$$\tilde{f}(k) = \langle f | \varphi_k \rangle = \int_{-\pi}^{\pi} f(y) e^{-iky} dy = \frac{1}{2\pi} \int_{-\pi}^{\pi} f(y) e^{-iky} dm(y).$$

Notice from Eq. (1.5) that $(t, x) \rightarrow u(t, x)$ is C^∞ for $t > 0$.

Exercise 1.15 (Heat Equation 2). Let $q_t(x) := \frac{1}{2\pi} \sum_{k \in \mathbb{Z}} e^{-\frac{t}{2}k^2} e^{ikx}$. Show;

- Eq. (1.5) may be rewritten as

$$u(t, x) = \int_{-\pi}^{\pi} q_t(x - y) f(y) dy$$

and

- $q_t(x)$ may be expressed

$$q_t(x) = \sum_{k \in \mathbb{Z}} p_t(x + k2\pi)$$

where $p_t(x) := \frac{1}{\sqrt{2\pi t}} e^{-\frac{1}{2t}x^2}$.

- Also show $u(t, x)$ may be written as

$$u(t, x) = (p_t * f)(x) := \int_{\mathbb{R}^d} p_t(x - y) f(y) dy.$$

Hint: To show $q_t(x) = \sum_{k \in \mathbb{Z}} p_t(x + k2\pi)$, use the Poisson summation formula (Exercise 1.13) and the Gaussian integration identity,

$$\hat{p}_t(\omega) = \frac{1}{\sqrt{2\pi}} \int_{\mathbb{R}} p_t(x) e^{i\omega x} dx = \frac{1}{\sqrt{2\pi}} e^{-\frac{t}{2}\omega^2}. \tag{1.6}$$

Exercise 1.16 (Wave Equation). Let $u \in C^2(\mathbb{R} \times \mathbb{R})$ be such that $u(t, \cdot) \in C_{per}(\mathbb{R})$ for all $t \in \mathbb{R}$. Further assume that u solves the wave equation, $u_{tt} = u_{xx}$. Let $f(x) := u(0, x)$ and $g(x) = \dot{u}(0, x)$. Show $\tilde{u}(t, k) := \langle u(t, \cdot), \varphi_k \rangle$ for $k \in \mathbb{Z}$ is twice continuously differentiable in t and $\frac{d^2}{dt^2} \tilde{u}(t, k) = -k^2 \tilde{u}(t, k)$. Use this result to show

$$u(t, x) = \sum_{k \in \mathbb{Z}} \left(\tilde{f}(k) \cos(kt) + \tilde{g}(k) \frac{\sin kt}{k} \right) e^{ikx} \tag{1.7}$$

with the sum converging absolutely. Also show that $u(t, x)$ may be written as

$$u(t, x) = \frac{1}{2} [f(x+t) + f(x-t)] + \frac{1}{2} \int_{-t}^t g(x+\tau) d\tau. \tag{1.8}$$

Hint: To show Eq. (1.7) implies (1.8) use

$$\begin{aligned} \cos kt &= \frac{e^{ikt} + e^{-ikt}}{2}, \\ \sin kt &= \frac{e^{ikt} - e^{-ikt}}{2i}, \text{ and} \\ \frac{e^{ik(x+t)} - e^{ik(x-t)}}{ik} &= \int_{-t}^t e^{ik(x+\tau)} d\tau. \end{aligned}$$

Exercise 1.17. Let $f \in L^1((-\pi, \pi])$ which we extend to a 2π -periodic function on \mathbb{R} and continue to denote by f . If there exists $q \in \mathbb{N}$ such that $f\left(x + \frac{2\pi}{q}\right) = f(x)$ for m -a.e. x , then $\tilde{f}(k) = 0$ unless q divides k .

Exercise 1.18. In this problem we assume the notation from subsection ?? with $d = 1$. For simplicity of notation we identify $L^2((-\pi, \pi], d\theta)$ with 2π -periodic functions on \mathbb{R} via,

$$L^2((-\pi, \pi], d\theta) \ni f \longleftrightarrow \sum_{n \in \mathbb{Z}} f(x + n2\pi) 1_{(-\pi, \pi]}(x + n2\pi) \in L^2_{per}(\mathbb{R}).$$

Given $\alpha \in \mathbb{R}$ let $(U_\alpha f)(\theta) = f(\theta + \alpha 2\pi)$ wherein we have used the above identification. If $\alpha \notin \mathbb{Q}$ show

$$M_\alpha = \text{Nul}(U_\alpha - I) = \mathbb{C} \cdot 1.$$

If $\alpha \in \mathbb{Q}$ write $\alpha = \frac{p}{q}$ where $\text{gcd}(q, p) = 1$, i.e. p and q are relatively prime. In this case show $M_\alpha = \text{Nul}(U_\alpha - I)$ consists of those $f \in L^2_{per}(\mathbb{R})$ such that $f\left(x + \frac{2\pi}{q}\right) = f(x)$ for m -a.e. x . [Consequently, combining this exercise with Mean Ergodic Theorem ?? shows,

$$\frac{1}{n} \sum_{k=0}^{n-1} U_\alpha^k \xrightarrow{s} P_{M_\alpha}$$

where M_α depends on α as described above.]

Exercise 1.19 (Folland 6.27 on p. 196. Hilbert's Inequality). **Hint:** See Theorem ?? which is Theorem 6.20 in Folland .

Exercise 1.20 (Folland 6.22). Exercise, Folland 6.22 on p. 192.

Exercise 1.21 (Global Integration by Parts Formula). Suppose that $f, g : \mathbb{R} \rightarrow \mathbb{C}$ are locally absolutely continuous functions² such that $f'g, fg'$, and fg are all Lebesgue integrable functions on \mathbb{R} . Prove the following integration by parts formula;

$$\int_{\mathbb{R}} f'(x) \cdot g(x) dx = - \int_{\mathbb{R}} f(x) \cdot g'(x) dx. \quad (1.9)$$

Similarly show that; if $f, g : [0, \infty) \rightarrow [0, \infty)$ are locally absolutely continuous functions such that $f'g, fg'$, and fg are all Lebesgue integrable functions on $[0, \infty)$, then

$$\int_0^{\infty} f'(x) \cdot g(x) dx = -f(0)g(0) - \int_0^{\infty} f(x) \cdot g'(x) dx. \quad (1.10)$$

Outline: 1. First use the theory developed to see that Eq. (1.9) holds if $f(x) = 0$ for $|x| \geq N$ for some $N < \infty$.

2. Let $\psi : \mathbb{R} \rightarrow [0, 1]$ be a continuously differentiable function such that $\psi(x) = 1$ if $|x| \leq 1$ and $\psi(x) = 0$ if $|x| \geq 2$.³ For any $\varepsilon > 0$ let $\psi_{\varepsilon}(x) = \psi(\varepsilon x)$. Write out the identity in Eq. (1.9) with $f(x)$ being replaced by $f(x)\psi_{\varepsilon}(x)$.

3. Now use the dominated convergence theorem to pass to the limit as $\varepsilon \downarrow 0$ in the identity you found in step 2.

4. A similar outline works to prove Eq. (1.10).

Exercise 1.22 (Heisenberg's Inequality). Suppose that $f : \mathbb{R} \rightarrow \mathbb{C}$ is a locally absolutely continuous function⁴, show

$$\int_{\mathbb{R}} |f(x)|^2 dx \leq 2 \left[\int_{\mathbb{R}} |xf(x)|^2 dx \right]^{1/2} \left[\int_{\mathbb{R}} |f'(x)|^2 dx \right]^{1/2}. \quad (1.11)$$

Hint: assuming the right hand side of the above inequality is finite show

$$\int_{\mathbb{R}} |f(x)|^2 dx = -2 \operatorname{Re} \int_{\mathbb{R}} x \bar{f}(x) f'(x) dx. \quad (1.12)$$

² This means that f and g restricted to any bounded interval in \mathbb{R} are absolutely continuous on that interval.

³ You may assume the existence of such a ψ , we will deal with this later.

⁴ This means that f restricted to any bounded interval in \mathbb{R} are absolutely continuous on that interval.

Exercise 1.23. Let

$$f(t) = \begin{cases} e^{-1/t} & \text{if } t > 0 \\ 0 & \text{if } t \leq 0. \end{cases}$$

Show $f \in C^{\infty}(\mathbb{R}, [0, 1])$. **Hints:** you might start by first showing $\lim_{t \downarrow 0} f^{(n)}(t) = 0$ for all $n \in \mathbb{N}_0$.

Exercise 1.24. Show $C_c^{\infty}(\mathbb{R}^d)$ is dense in $L^p(\mathbb{R}^d, m)$ for any $1 \leq p < \infty$.

Exercise 1.25. If $f \in L^1_{loc}(\mathbb{R}^d, m)$ and $\varphi \in C_c^1(\mathbb{R}^d)$, then $f * \varphi \in C^1(\mathbb{R}^d)$ and $\partial_i(f * \varphi) = f * \partial_i \varphi$. Moreover if $\varphi \in C_c^{\infty}(\mathbb{R}^d)$ then $f * \varphi \in C^{\infty}(\mathbb{R}^d)$.

Exercise 1.26 (Integration by Parts). Suppose that $(x, y) \in \mathbb{R} \times \mathbb{R}^{d-1} \rightarrow f(x, y) \in \mathbb{C}$ and $(x, y) \in \mathbb{R} \times \mathbb{R}^{d-1} \rightarrow g(x, y) \in \mathbb{C}$ are measurable functions such that for each fixed $y \in \mathbb{R}^d$, $x \rightarrow f(x, y)$ and $x \rightarrow g(x, y)$ are continuously differentiable. Also assume $f \cdot g$, $\partial_x f \cdot g$ and $f \cdot \partial_x g$ are integrable relative to Lebesgue measure on $\mathbb{R} \times \mathbb{R}^{d-1}$, where $\partial_x f(x, y) := \frac{d}{dt} f(x+t, y)|_{t=0}$. Show

$$\int_{\mathbb{R} \times \mathbb{R}^{d-1}} \partial_x f(x, y) \cdot g(x, y) dx dy = - \int_{\mathbb{R} \times \mathbb{R}^{d-1}} f(x, y) \cdot \partial_x g(x, y) dx dy. \quad (1.13)$$

(Note: this result and Fubini's theorem proves Lemma ??.)

Hints: Let $\psi \in C_c^{\infty}(\mathbb{R})$ be a function which is 1 in a neighborhood of $0 \in \mathbb{R}$ and set $\psi_{\varepsilon}(x) = \psi(\varepsilon x)$. First verify Eq. (1.13) with $f(x, y)$ replaced by $\psi_{\varepsilon}(x) f(x, y)$ by doing the x -integral first. Then use the dominated convergence theorem to prove Eq. (1.13) by passing to the limit, $\varepsilon \downarrow 0$.

Exercise 1.27 (Wirtinger's inequality, Folland 8.18). Given $a > 0$ and $f \in C^1([0, a], \mathbb{C})$ such that $f(0) = f(a) = 0$, show⁵

$$\int_0^a |f(x)|^2 dx \leq \left(\frac{a}{\pi}\right)^2 \int_0^a |f'(x)|^2 dx.$$

Hint: to use the notation above, let $\pi L = a$ and extend f to $[-a, 0]$ by setting $f(-x) = -f(x)$ for $0 \leq x \leq a$. Now compute $\int_0^a |f(x)|^2 dx$ and $\int_0^a |f'(x)|^2 dx$ in terms of their Fourier coefficients, $\langle f | \varphi_k^L \rangle_L$ and $\langle f' | \varphi_k^L \rangle_L$ respectively.

Exercise 1.28. Let

$$L = \sum_{|\alpha| \leq k} a_{\alpha}(x) \partial^{\alpha} \quad (1.14)$$

with $a_{\alpha} \in \mathcal{P}$. Show $L(\mathcal{S}) \subset \mathcal{S}$ and in particular $\partial^{\alpha} f$ and $x^{\alpha} f$ are back in \mathcal{S} for all multi-indices α .

⁵ This inequality is sharp as is seen by taking $f(x) = \sin(\pi x/a)$.

Exercise 1.29. In this problem let $d = 1$ so that $x, \xi \in \mathbb{R} = \mathbb{R}^1$. For any $m > 0$, show

$$\mathcal{F} \left[e^{-m|x|} \right] (\xi) = \frac{2m}{\sqrt{2\pi}} \frac{1}{m^2 + \xi^2}$$

and

$$\mathcal{F} \left(\frac{1}{m^2 + \xi^2} \right) (x) = \frac{\sqrt{2\pi}}{2m} e^{-m|x|}.$$

More precisely these equations mean;

$$\begin{aligned} \mathcal{F} \left[x \rightarrow e^{-m|x|} \right] (\xi) &= \frac{2m}{\sqrt{2\pi}} \frac{1}{m^2 + \xi^2} \text{ and} \\ \mathcal{F} \left(\xi \rightarrow \frac{1}{m^2 + \xi^2} \right) (x) &= \frac{\sqrt{2\pi}}{2m} e^{-m|x|} \end{aligned}$$

or equivalently,

$$\mathcal{F} \left[e^{-m|\cdot|} \right] = \frac{2m}{\sqrt{2\pi}} \frac{1}{m^2 + (\cdot)^2} \text{ and } \mathcal{F} \left(\frac{1}{m^2 + (\cdot)^2} \right) = \frac{\sqrt{2\pi}}{2m} e^{-m|\cdot|}.$$

Exercise 1.30. Using the identity

$$\frac{1}{\xi^2 + 1} = \int_0^\infty e^{-s(\xi^2 + 1)} ds$$

along with Exercise 1.29 and the known Fourier transform of Gaussians to show

$$e^{-|x|} = \int_0^\infty ds \frac{1}{\sqrt{\pi s}} e^{-s} e^{-\frac{x^2}{4s}} \text{ for all } x \in \mathbb{R}. \quad (1.15)$$

Thus we have written $e^{-|x|}$ as an average of Gaussians.

Exercise 1.31. Now let $x \in \mathbb{R}^d$ and $|x|^2 := \sum_{i=1}^d x_i^2$ be the standard Euclidean norm. Show for all $m > 0$ that

$$\mathcal{F} \left[e^{-m|x|} \right] (\xi) = \frac{2^{d/2}}{\sqrt{\pi}} \Gamma \left(\frac{d+1}{2} \right) \frac{m}{\left(m^2 + |\xi|^2 \right)^{\frac{d+1}{2}}}, \quad (1.16)$$

where $\Gamma(x)$ in the gamma function defined as

$$\Gamma(x) := \int_0^\infty t^x e^{-t} \frac{dt}{t}.$$

Hint: By Exercise 1.30 with x replaced by $m|x|$ we know that

$$e^{-m|x|} = \int_0^\infty ds \frac{1}{\sqrt{\pi s}} e^{-s} e^{-\frac{m^2}{4s}|x|^2} \text{ for all } x \in \mathbb{R}^d.$$

Exercise 1.32. Show for $f \in \mathcal{S}(\mathbb{R})$ that;

1. For all $x \in \mathbb{R}$,

$$|f(x)| \leq \frac{1}{\sqrt{2\pi}} \|\hat{f}\|_1 = \frac{1}{\sqrt{2\pi}} \int_{\mathbb{R}} |\hat{f}(k)| dk$$

and

$$|f(x)| \leq \frac{1}{\sqrt{2}} \left[\int_{\mathbb{R}} |\hat{f}(k)|^2 (1+k^2) dk \right]^{1/2}.$$

2. Use the last displayed inequality and the basic properties of the Fourier transform to prove the ‘‘Sobolev inequality,’’

$$|f(x)|^2 \leq \frac{1}{2} \left[\|f\|_2^2 + \|f'\|_2^2 \right] \text{ for all } x \in \mathbb{R},$$

where

$$\|f\|_2^2 := \int_{\mathbb{R}} |f(x)|^2 dx.$$

Exercise 1.33 (Sampling Theorem). Let

$$\text{sinc } x = \begin{cases} \frac{\sin \pi x}{\pi x} & \text{if } x \neq 0 \\ 1 & \text{if } x = 0 \end{cases}$$

and for any $a \in (0, \infty)$, let

$$\mathcal{H}_a = \{f \in L^2(m) : \hat{f}(\xi) = 0 \text{ a.e. when } |\xi| > \pi a\}.$$

Show

1. Show that every $f \in \mathcal{H}_a$ has a version⁶ $f_0 \in C_0(\mathbb{R})$ and moreover,

$$\|f_0\|_\infty \leq \sqrt{a} \|f\|_{L^2(m)}. \quad (1.17)$$

[We now identify f with this continuous version.] **Hint:** after identifying $L^2([-\pi a, \pi a], \lambda)$ as a subspace of $L^2(\mathbb{R}, \lambda)$ one has

$$\mathcal{H}_a = \mathcal{F}^{-1} L^2([-\pi a, \pi a], \lambda).$$

2. Show by direct computation that

$$\mathcal{F}^{-1} \left[\frac{1}{\sqrt{2\pi a}} e^{-in\xi/a} 1_{|\xi| \leq \pi a} \right] (x) = \text{sinc}(ax - n).$$

⁶ We say that f_0 is a version of f if $f(x) = f_0(x)$ for m -a.e. x .

3. If $f \in \mathcal{H}_a$ then (assuming f is the C_0 - version as in part a), show

$$f(x) = \sum_{k=-\infty}^{\infty} f(k/a) \operatorname{sinc}(ax - k),$$

where the series converges both uniformly and in L^2 . [**Hint:** Start by writing $\hat{f}(\xi)$ for $|\xi| \leq \pi a$ as a Fourier expansion in the orthonormal basis $\{e^{-in\xi/a}\}_{n=-\infty}^{\infty}$ for $L^2\left([-\pi a, \pi a], \frac{m}{2\pi a}\right)$.]

In the terminology of signal analysis, a signal of band width $2\pi a$ is completely determined by sampling its value at a sequence of points $\{k/2\pi a\}$ whose spacing is the reciprocal of the bandwidth.