

Competitive adsorption and ordered packing of counterions near highly charged surfaces: From mean-field theory to Monte Carlo simulations

Jiayi Wen,^{1,*} Shenggao Zhou,^{1,2,†} Zhenli Xu,^{3,‡} and Bo Li^{1,§}

¹*Department of Mathematics, and the NSF Center for Theoretical Biological Physics, University of California, San Diego, 9500 Gilman Drive, Mail Code 0112, La Jolla, California 92093-0112, USA*

²*Department of Mathematics, Zhejiang University, 38 Zheda Road, Hangzhou 310027, People's Republic of China*

³*Department of Mathematics and Institute of Natural Sciences, Shanghai Jiao Tong University, 800 Dongchuan Road, Shanghai 200240, People's Republic of China*

(Received 19 January 2012; published 26 April 2012)

Competitive adsorption of counterions of multiple species to charged surfaces is studied by a size-effect-included mean-field theory and Monte Carlo (MC) simulations. The mean-field electrostatic free-energy functional of ionic concentrations, constrained by Poisson's equation, is numerically minimized by an augmented Lagrangian multiplier method. Unrestricted primitive models and canonical ensemble MC simulations with the Metropolis criterion are used to predict the ionic distributions around a charged surface. It is found that, for a low surface charge density, the adsorption of ions with a higher valence is preferable, agreeing with existing studies. For a highly charged surface, both the mean-field theory and the MC simulations demonstrate that the counterions bind tightly around the charged surface, resulting in a stratification of counterions of different species. The competition between mixed entropy and electrostatic energetics leads to a compromise that the ionic species with a higher valence-to-volume ratio has a larger probability to form the first layer of stratification. In particular, the MC simulations confirm the crucial role of ionic valence-to-volume ratios in the competitive adsorption to charged surfaces that had been previously predicted by the mean-field theory. The charge inversion for ionic systems with salt is predicted by the MC simulations but not by the mean-field theory. This work provides a better understanding of competitive adsorption of counterions to charged surfaces and calls for further studies on the ionic size effect with application to large-scale biomolecular modeling.

DOI: [10.1103/PhysRevE.85.041406](https://doi.org/10.1103/PhysRevE.85.041406)

PACS number(s): 82.70.Dd, 87.10.Rt, 82.60.Lf, 87.15.ak

I. INTRODUCTION

Electrostatic interactions play an important role in many complex systems, such as biological processes, soft matter material, nanofluids, and electrochemical devices [1–8]. Accurate and efficient modeling and computations of such interactions have been challenging due to the inhomogeneity, complicated geometry, multiple scales, and the nature of the many-body interaction of an underlying charged system.

A common scenario of electrostatic interactions is a mixture of crowded mobile ions of multiple species with different valences and sizes in an electrolyte surrounding an external charged surface. Excluded-volume effects or size effects of such mobile ions, in particular effects of different ionic sizes, contribute significantly to the electrostatic free energy and forces, which in turn determine the structure and stability of an underlying system. For instance, the size of monovalent cations can influence the stability of RNA tertiary structures [9], and differences in ionic sizes can also affect how mobile ions bind to nucleic acids [10,11]. Concentrations of ions in an ion channel can reach as high as dozens of mol/L (about 30 mol/L in calcium and sodium channels), and the ionic sizes can affect the ion transport and channel selectivity [12]. Detailed density-functional theory calculations, Monte

Carlo simulations, and integral equations calculations confirm some of these experimentally observed properties due to the nonuniformity of ionic sizes [13–17].

Historically, many theoretical studies of electrostatic interactions have been based on the classical, mean-field, Poisson-Boltzmann (PB) theory [18–21]. In particular, such a theory has been successfully applied in biomolecular modeling and colloidal science (see [1,3,8,22,23] and the references therein). In the PB theory, electrolytes are treated as ideal ionic gases, and the ionic concentrations are related to the electrostatic potential by the Boltzmann distributions. This theory, often very efficient, thus works well for monovalent ions, low surface charge densities, and high solvent dielectric coefficients. The mathematical form of the PB theory is the PB equation, which is Poisson's equation for the electrostatic potential with the equilibrium ionic concentrations given by the Boltzmann distributions via the potential. In a variational setting, such distributions result from the equilibrium conditions for a mean-field electrostatic free-energy functional of ionic concentrations where the potential is determined by Poisson's equation [24–28]. Despite its success in many applications, the classical PB theory is known to fail in capturing the ion-ion correlations and ionic size effects, particularly for highly charged systems at molecular scales [29,30].

For years, attempts have been made to include ionic size effects, particularly nonuniform ionic size effects, into a PB-like efficient approach [26,31–37] (see also [38–45]). One of the key ideas has been to introduce the local concentration of solvent molecules, in addition to those of ions of multiple

*j2wen@ucsd.edu

†s4zhou@math.ucsd.edu

‡xuzl@sjtu.edu.cn

§bli@math.ucsd.edu

species and to incorporate all the ionic and solvent molecular volumes in the entropic part of a mean-field electrostatic free-energy functional. If all linear sizes (including that of solvent molecules) are the same, such a free-energy functional can be derived using a lattice-gas model [32,35,36]. Moreover, there are explicit formulas, the generalized Boltzmann distributions, relating equilibrium ionic concentrations and the corresponding electrostatic potential. These distributions, together with Poisson's equation, lead to the generalized PB equation for the case of a uniform ionic size [26,27,46]. For a system of three ionic species with two different ionic sizes, Chu *et al.* [34] derived a different size-modified PB equation from a similar lattice-gas model and applied this equation to study the ionic size effect in the binding of ions to DNA. For a general system, Tresset [36] derived an expression of the free energy with an effective volume fraction of free space, under the assumption that the ionic excluded volumes are dispersed from each other to a reasonable extent.

For the general case of multiple ionic species with different valences and sizes, Li [26] proposed and analyzed a semiphenomenological free-energy functional of ionic concentrations with Poisson's equation as a constraint for the electrostatic potential. This functional is obtained simply by using different individual ionic sizes instead of a uniform size in the previous functional derived from a lattice-gas model. Equilibrium conditions for the new and general free-energy functional are nonlinear algebraic equations for the equilibrium concentrations. It is shown that such conditions determine completely the dependence of equilibrium ionic concentrations on the corresponding electrostatic potential [26]. Explicit formulas of such dependence and hence Boltzmann-like distributions for the equilibrium concentrations, however, seem unavailable. Therefore, there is no explicit PB-like equation of the electrostatic potential in the general case.

Nevertheless, Zhou *et al.* [37] developed a robust numerical method for minimizing such a functional to obtain the equilibrium ionic concentrations and the corresponding electrostatic potential. The starting point there is to reformulate the variational problem as a constrained optimization problem [47,48]. An augmented Lagrange multiplier method is then constructed and implemented to solve this constrained optimization problem. Extensive numerical results reported in [37] demonstrate that the new mean-field, size-effect-included model can describe many detailed properties of ionic concentrations, including the stratification of concentrations, that have been predicted by other refined models but not by the classical PB theory (cf. [33,36]). In particular, it is found that the ionic valence-to-volume ratio is the key parameter in the stratification [37].

In this work, we study the ionic size or excluded-volume effect to the structure of electrical double layer in the vicinity of a highly charged surface, using both the mean-field model and Monte Carlo (MC) simulations. Our goal is twofold. First, we would like to understand how counterions with different valences and sizes compete in the adsorption to the charged surface and how the ionic valence-to-volume ratio affects the ordering of ion packing near such a surface. Second, we would like to examine the validity of the mean-field theory with nonuniform size effects by comparing it with the MC simulations.

The adsorption of counterions to a charged surface is determined by the competition between the entropic and the energetic contributions of an underlying system of electrolyte. The ionic size effect is quite significant in such adsorption, since the excluded volume of crowded ions reduces the mixed entropy, and thus increases the Helmholtz free energy of the total system. Concentrations of counterions can reach maximal values at the charged surface controlled by the ionic sizes. The competition of entropy and energy results in a stratification of counterions of different species in the electrical double layer, as revealed in both experimental investigations [49] and theoretical predictions [36,37,50,51]. For a low surface charge density, the electrostatic interaction dominates and the ions with higher valence are most likely to stay closest to the charged surface. For a highly charged surface, smaller counterions are stronger in competition to form the first layer of the stratification [16,51]. Our mean-field numerical computations and MC simulations reproduce these results. In particular, our MC simulations validate the prediction by the mean-field theory of the role of ionic valence-to-volume ratios in the counterion stratification.

MC simulations treat an underlying system of electrolyte as a set of discrete particles and provide equilibrium properties of the system with statistical averages [52,53]. Such simulations have been a standard tool in the study of structures of electrical double layer, if the geometry of the charged surface is not too complicated [16,51,54,55]. In MC simulations, ionic size and correlation effects are automatically included, and image charge effects can also be included [56–60]. Therefore, the correlation-induced phenomena, such as charge inversion and like-charge attraction, can be described by MC simulations [29,55].

Our simulation system consists of a spherical macroion immersed centrally in an electrolyte system. There are counterions of multiple species in the electrolyte. The entire system is assumed to be neutral in charge. The parameters of the system include the linear size of the simulation box, the radius and constant surface charge density of the macroion, and the valence, volume, and total number of each species of (micro) mobile ions. The same set of parameters is used in our MC simulations and mean-field computations. We use unrestrictive primitive models of ionic system, treating ions as hard spheres. Based on such a model, we use canonical ensemble MC simulations with Metropolis criterion. We plot the radial particle density function for each species of mobile ions. Such functions are compared with the corresponding equilibrium ionic concentrations predicted by our mean-field theory.

The rest of this paper is organized as follows: In Sec. II, we introduce the mean-field theory and numerical method for nonuniform ionic size effects. In Sec. III, we describe our method of MC simulations. In Sec. IV, we present and discuss the results of our MC simulations and mean-field computations. Finally, in Sec. V, we draw conclusions.

II. MEAN-FIELD THEORY AND METHOD

We consider an electrolyte with M species of ions. For each i ($1 \leq i \leq M$), we denote by z_i the valence and v_i the volume of an ion of the i th species. We also denote by N_i the

total number of ions of the i th species. The total number of all ions is $N = \sum_{i=1}^M N_i$. We assume that there is a spherical colloidal particle—a charged macroion—of radius R inside the electrolyte solution and that its charge effect is described effectively by a constant surface charge density, denoted σ . We assume the system charge neutrality

$$ze + \sum_{i=1}^M N_i z_i e = 0, \quad (2.1)$$

where $z = 4\pi R^2 \sigma / e$ is the valence of the macroion and e is the elementary charge.

We assume that the entire system occupies the cubical region $(-L/2, L/2)^3$ with the linear size $L > 2R$ and that the macroion occupies the spherical region B_R of radius R centered at the origin. Therefore, all the ions are in the region $\Omega = (-L/2, L/2)^3 \setminus B_R$. We denote by $\Gamma = \partial B_R$ the boundary of the sphere B_R , that is, the spherical surface of the macroion.

A. A mean-field theory with nonuniform size effects

For each i ($1 \leq i \leq M$), we denote by $c_i(\mathbf{r})$ the local concentration at a spatial point $\mathbf{r} \in \Omega$ of ions of the i th species. The charge density of solution is then given by $\sum_{i=1}^M z_i e c_i(\mathbf{r})$ ($\mathbf{r} \in \Omega$). All the concentrations $c_i(\mathbf{r})$ are constrained by

$$\int_{\Omega} c_i dV = N_i, \quad i = 1, \dots, M. \quad (2.2)$$

We also denote by v_0 the volume of a solvent molecule. The local concentration $c_0 = c_0(\mathbf{r})$ of the solvent molecules is defined by

$$c_0(\mathbf{r}) = v_0^{-1} \left[1 - \sum_{i=1}^M v_i c_i(\mathbf{r}) \right] \quad \text{for all } \mathbf{r} \in \Omega.$$

For a given set of ionic concentrations $c = (c_1, \dots, c_M)$, a mean-field approximation of the electrostatic free energy is given by

$$F[c] = F_{\text{pot}}[c] + F_{\text{ent}}[c]. \quad (2.3)$$

The first part $F_{\text{pot}}[c]$ is the electrostatic potential energy, defined by

$$F_{\text{pot}}[c] = \int_{\Omega} \frac{1}{2} \left(\sum_{i=1}^M z_i e c_i \right) \Phi dV + \int_{\Gamma} \frac{1}{2} \sigma \Phi dS, \quad (2.4)$$

where Φ is the electrostatic potential. It is determined by Poisson's equation,

$$\nabla \cdot \varepsilon \varepsilon_0 \nabla \Phi = - \sum_{i=1}^M z_i e c_i \quad \text{in } \Omega, \quad (2.5)$$

together with the boundary condition

$$\varepsilon \varepsilon_0 \frac{\partial \Phi}{\partial n} = \begin{cases} \sigma & \text{on } \Gamma, \\ 0 & \text{on } \Gamma_{\text{box}}, \end{cases} \quad (2.6)$$

where ε_0 is the vacuum permittivity, ε is the relative permittivity or dielectric coefficient of the solution, and n is the exterior unit normal at the boundary of Ω that consists of the spherical surface Γ and the boundary, Γ_{box} , of the box $(-L/2, L/2)^3$. We

shall assume that ε is a constant in the entire solution region Ω . Notice that Φ is not an independent variable of the functional $F[c]$.

The second part $F_{\text{ent}}[c]$ is the entropic contribution. It is given by [26,37]

$$F_{\text{ent}}[c] = k_B T \sum_{i=0}^M \int_{\Omega} c_i [\ln(v_i c_i) - 1] dV, \quad (2.7)$$

where k_B is the Boltzmann constant and T is the absolute temperature. Notice that the summation index starts from $i = 0$. Notice also that in the variational approach to the classical PB equation, the solvent entropy is not included and all the ionic linear sizes $v_i^{1/3}$ are replaced by the de Broglie wave length [24,25,27].

The set of equilibrium ionic concentrations $c = (c_1, \dots, c_M)$ is defined to minimize the free-energy functional (2.3), subject to the constraint (2.2). The equilibrium electrostatic potential is determined by the corresponding equilibrium ionic concentrations through Poisson's equation (2.5) and the boundary condition (2.6).

Alternatively, we can introduce for each i the chemical potential μ_i for ions of the i th species, and add the term

$$- \sum_{i=1}^M \int_{\Omega} \mu_i c_i dV \quad (2.8)$$

to the free energy $F[c]$ in (2.3). The chemical potentials μ_i ($i = 1, \dots, M$) can be regarded as Lagrange multipliers accounting for the constraint (2.2). With these chemical potentials, one minimizes the new, total electrostatic free-energy functional that now consists of all the integral terms in Eqs. (2.4), (2.7), and (2.8), without the constraint (2.2).

Taking the variational derivative with respect to each concentration field $c_i(\mathbf{r})$ of the new, total free energy and setting it to 0, we obtain with suitable boundary conditions for Poisson's equation (2.5) the conditions for equilibrium concentrations c_1, \dots, c_M [26]

$$\frac{v_i}{v_0} \ln(v_0 c_0(\mathbf{r})) - \ln(v_i c_i(\mathbf{r})) = \frac{1}{k_B T} [z_i e \Phi(\mathbf{r}) - \mu_i] \quad (2.9)$$

for all $\mathbf{r} \in \Omega$, $i = 1, \dots, M$.

In the special case that $v_0 = v_1 = \dots = v_M$, one can solve this system of nonlinear algebraic equations to obtain explicit formulas of $c_i(\mathbf{r}) = c_i(\Phi(\mathbf{r}))$ ($i = 1, \dots, M$). These are the generalized Boltzmann distributions. For the general case, it is known that the conditions (2.9) determine uniquely $c_i(\mathbf{r}) = c_i(\Phi(\mathbf{r}))$ ($i = 1, \dots, M$); but explicit formulas for such dependence seem unavailable (see Refs. [26,27]).

B. A constrained optimization method

By integration by parts, Poisson's equation (2.5), and the boundary conditions (2.6), we can rewrite the free-energy functional (2.3), which is the sum of $F_{\text{pot}}[c]$ given in Eq. (2.4)

and $F_{\text{ent}}[c]$ given in Eq. (2.7), as

$$F[\Phi, c] = \int_{\Omega} \left\{ \frac{\varepsilon \varepsilon_0}{2} |\nabla \Phi|^2 + k_B T \sum_{i=0}^M c_i [\ln(v_i c_i) - 1] \right\} dV,$$

where Φ solves the boundary-value problem of Poisson's equation (2.5) and (2.6). Notice that the dependence of F on Φ is now explicitly indicated. One can verify mathematically that the minimization of $F[c]$ defined in (2.3) over all c subject to (2.2) is equivalent to that of $F[\Phi, c]$ over all (Φ, c) subject to (2.2), (2.5), and (2.6).

Introduce the Bjerrum length $l_B = e^2 / (4\pi \varepsilon \varepsilon_0 k_B T)$. Define $\Phi' = e\Phi / (k_B T)$, $c'_i = 4\pi l_B c_i$ and $v'_i = (4\pi l_B)^{-1} v_i$ ($0 \leq i \leq M$), $N'_i = 4\pi l_B N_i$ ($1 \leq i \leq M$), $\sigma' = 4\pi l_B \sigma / e$, and $\omega' = (4\pi l_B)^{-1/3} \omega$ for $\omega = \Gamma, \Gamma_{\text{box}}, \text{ or } \Omega$. Then $F[\Phi, c] = \varepsilon \varepsilon_0 (k_B T / e)^2 F'[\Phi', c']$, where

$$F'[\Phi', c'] = \int_{\Omega'} \left\{ \frac{1}{2} |\nabla \Phi'|^2 + \sum_{i=0}^M c'_i [\ln(v'_i c'_i) - 1] \right\} dV, \quad (2.10)$$

and c'_0 is defined similarly using the primed quantities. The constraint (2.2), Poisson's equation (2.5), and the boundary condition (2.6) become now

$$\int_{\Omega'} c'_i dV = N'_i, \quad (2.11)$$

$$\Delta \Phi' = - \sum_{i=1}^M z_i c'_i \quad \text{in } \Omega', \quad (2.12)$$

$$\frac{\partial \Phi'}{\partial n'} = \begin{cases} \sigma' & \text{on } \Gamma', \\ 0 & \text{on } \Gamma'_{\text{box}}, \end{cases} \quad (2.13)$$

respectively, where n' is the unit exterior normal at the boundary of Ω' .

For simplicity, we drop all the primes in what follows.

We apply an augmented Lagrange multiplier method [61,62] to numerically minimize the functional $F[\Phi, c]$ defined in (2.10) subject to (2.11)–(2.13) (with all the primes dropped). Our method is an improved version of that developed in our previous work [37] for minimizing numerically a similar functional formulated using (\mathbf{E}, c) instead of (Φ, c) , where $\mathbf{E} = -\nabla \Phi$ is the electric field. In the augmented Lagrange multiplier formulation, we solve the corresponding saddle-point problem,

$$\min_{(\Phi, c)} \max_{(\Psi, \Lambda, \mathbf{s})} \hat{L}(\Phi, c, \Psi, \Lambda, \mathbf{s}), \quad (2.14)$$

where $\Lambda = (\lambda_1, \dots, \lambda_M) \in \mathbb{R}^M$, $\mathbf{s} = (s_1, \dots, s_M) \in \mathbb{R}^M$ with each $s_i \geq 0$, and

$$\begin{aligned} \hat{L}(\Phi, c, \Psi, \Lambda, \mathbf{s}) &= F[\Phi, c] + \int_{\Omega} \Psi \left(\Delta \Phi + \sum_{i=1}^M z_i c_i \right) dV + \sum_{i=1}^M \lambda_i \left(\int_{\Omega} c_i dV - N_i \right) + \sum_{i=1}^M \frac{s_i}{2} \left(\int_{\Omega} c_i dV - N_i \right)^2 \\ &= \int_{\Omega} \left\{ \frac{1}{2} |\nabla \Phi|^2 + \sum_{i=0}^M c_i [\ln(v_i c_i) - 1] \right\} dV + \int_{\Omega} \Psi \left(\Delta \Phi + \sum_{i=1}^M z_i c_i \right) dV \\ &\quad + \sum_{i=1}^M \lambda_i \left(\int_{\Omega} c_i dV - N_i \right) + \sum_{i=1}^M \frac{s_i}{2} \left(\int_{\Omega} c_i dV - N_i \right)^2. \end{aligned}$$

The function Ψ is the Lagrange multiplier for Poisson's equation (2.5). It satisfies the same boundary conditions as for Φ [cf. (2.13) (no primes)]. The numbers $\lambda_1, \dots, \lambda_M$ are the Lagrange multipliers for the constraint (2.11) (no primes). The last summation term is a penalty term. It is added to stabilize and accelerate our numerical iterations.

The solution $(\Phi, c, \Psi, \Lambda, \mathbf{s})$ to the saddle-point problem (2.14) is determined by the following equations:

$$\frac{\partial \hat{L}}{\partial \Phi} = -\Delta(\Phi - \Psi) = 0 \quad \text{in } \Omega, \quad (2.15)$$

$$\frac{\partial \hat{L}}{\partial \Psi} = \Delta \Phi + \sum_{i=1}^M z_i c_i = 0 \quad \text{in } \Omega, \quad (2.16)$$

$$\frac{\partial \hat{L}}{\partial c_i} = \ln(v_i c_i) - \frac{v_i}{v_0} \ln \left(v_0 \left(1 - \sum_{j=1}^M v_j c_j \right) \right) + z_i \Psi + \lambda_i + s_i \left(\int_{\Omega} c_i dV - N_i \right) = 0 \quad \text{in } \Omega, \quad i = 1, \dots, M, \quad (2.17)$$

$$\frac{\partial \hat{L}}{\partial \lambda_i} = \int_{\Omega} c_i dV - N_i = 0, \quad i = 1, \dots, M. \quad (2.18)$$

Since both Φ and Ψ satisfy the same boundary conditions, Eq. (2.15) implies that they differ by an additive constant. We may choose this constant to be 0 and assume that $\Psi = \Phi$ in Ω . Notice that Eq. (2.16) is Poisson's equation (2.12) (no primes)

and Eq. (2.18) is the constraint (2.11) (no primes). As pointed out before, the nonlinear system of algebraic equations (2.17) has a unique solution $c = (c_1, \dots, c_M)$ but its explicit solution formulas seem unavailable [26].

The entire system of equations is equivalent now to the three sets of Eqs. (2.16)–(2.18) with Ψ in (2.17) replaced with Φ . We solve these equations by the following algorithm.

Step 0. Distribute the total surface charge $4\pi R^2\sigma$ uniformly on the spherical surface by interpolation onto the nearest grids [37,47]. Initialize $\Phi^{(0)}$, $c^{(0)} = (c_1^{(0)}, \dots, c_M^{(0)})$, $\Lambda^{(0)} = (\lambda_1^{(0)}, \dots, \lambda_M^{(0)})$, and $\mathbf{s}^{(0)} = (s_1^{(0)}, \dots, s_M^{(0)})$. Choose a parameter $\gamma > 1$. Set $l = 0$.

Step 1. Solve Eq. (2.16) with c_i replaced with $c_i^{(l)}$, together with the boundary condition (2.13), to obtain the solution $\Phi^{(l+1)}$.

Step 2. Use Newton's method to solve Eq. (2.17) (where Ψ is replaced with Φ) with Φ , Λ , and \mathbf{s} replaced with $\Phi^{(l+1)}$, $\Lambda^{(l)}$, and $\mathbf{s}^{(l)}$, respectively, to obtain the solution $c^{(l+1)}$.

Step 3. Update the Lagrange multipliers

$$\lambda_i^{(l+1)} = \lambda_i^{(l)} + s_i^{(l)} \left(\int_{\Omega} c_i^{(l+1)} dV - N_i \right), \quad i = 1, \dots, M.$$

Update the penalty parameters $s_i^{(l+1)} = \gamma s_i^{(l)}$ ($i = 1, \dots, M$).

Step 4. Test convergence. If not, set $l \leftarrow l + 1$ and go to Step 1.

The parameter $\gamma > 1$ is used only for updating s_i ($i = 1, \dots, M$). Various kinds of approximations can be used to solve the boundary-value problem of Poisson's equation. For instance, we can use the periodic boundary condition instead, and apply the fast Fourier transform. In this case, we have the linear complexity in terms of the number of unknowns of resulting system of linear equations. We note that the matrix-vector multiplication can be avoided in Newton's iteration scheme for solving the system (2.17) (with Ψ replaced by Φ), since the exact formula of the inverse of related Jacobian matrix can be obtained. See Ref. [37] for more details.

III. MONTE CARLO SIMULATIONS

We consider the same system described in the previous section: A macroion occupying the sphere B_R of radius R centered at the origin, with a constant surface charge density σ , is immersed in an electrolyte in the box $(-L/2, L/2)^3$. There are M species of (micro) ions in the region $\Omega = (-L/2, L/2)^3 \setminus B_R$. For each i ($1 \leq i \leq M$), an ion of the i th species has valence z_i and volume v_i . The number of ions of the i th species is N_i ; and the total number of all (micro) ions is $N = \sum_{i=1}^M N_i$. We use an unrestricted primitive model for our underlying electrolyte system and apply the canonical ensemble MC simulations with the Metropolis criterion [50,52,63–66].

In a primitive model of electrolytes, the mobile ions are represented by charged hard spheres and the solvent is modeled through its dielectric permittivity ϵ . We label all the (micro) ions by $k = 1, \dots, N$. We denote by \hat{z}_k and \hat{R}_k the valence and radius of the k th ion. If the k th ion is of the i th type ($1 \leq i \leq M$), then its valence is $\hat{z}_k = z_i$ and its volume is $4\pi \hat{R}_k^3/3 = v_i$. For convenience, we label the spherical macroion by 0 and denote $\hat{R}_0 = R$, the radius of the macroion. We also denote its valence by $\hat{z}_0 = z = 4\pi R^2\sigma/e$.

For a given configuration of the system, the Hamiltonian is defined to be the work needed to bring all the ions from infinity to their current positions. It is the sum of all

pairwise interaction energies between all the ions, including the macroion. We only consider the hard-sphere contribution and the Coulomb interaction. Therefore, we define the total potential energy of the system to be

$$U = \sum_{0 \leq j < k \leq N} u_{jk},$$

where

$$\beta u_{jk} = \begin{cases} \frac{l_B \hat{z}_j \hat{z}_k}{r_{jk}} & \text{if } r_{jk} \geq \hat{R}_j + \hat{R}_k, \\ \infty & \text{if } r_{jk} < \hat{R}_j + \hat{R}_k. \end{cases} \quad (3.1)$$

Here, $\beta = (k_B T)^{-1}$, $l_B = e^2 \beta / (4\pi \epsilon \epsilon_0)$ is the Bjerrum length, and r_{jk} is the center-center distance between the j th and k th ions. Notice that, in the case $r_{jk} \geq \hat{R}_j + \hat{R}_k$, u_{jk} is just the Coulomb interaction energy between the j th and k th ions in the solvent with the relative dielectric permittivity ϵ . We consider the water solvent at room-temperature and thus take $l_B = 7 \text{ \AA}$.

Our MC simulations consist of a sequence of single-particle moves with the periodical boundary condition. In each move, we randomly select an individual particle (i.e., mobile ion). Let us assume that the selected particle is centered at \mathbf{p} . We then randomly generate a positive number, denoted a , from the interval $[0, \Delta_{\max}]$ for some parameter $\Delta_{\max} > 0$. We finally place the (center of) selected particle randomly on the sphere of radius a centered at \mathbf{p} . We use the L -periodical boundary condition in each direction, so that all the ions remain in the region Ω of electrolyte. The parameter Δ_{\max} can change during the MC moves. The acceptance or rejection of the move is determined by the Metropolis criterion. We calculate the difference $\Delta U = U_{\text{new}} - U_{\text{old}}$ of the energies of the previous (old) and current (new) configurations. If $\Delta U \leq 0$, the move is accepted. Otherwise, it is accepted if $\exp(-\beta \Delta U)$ is greater than a randomly generated number in $[0, 1]$.

The entire sequence of our MC moves are divided into three parts: acceleration, equilibration, and statistics. Typically, our simulation system consists of $M = 3$ or 4 ionic species; and the number of ions in each of these species can vary from 25 to 50 and to 200. With these parameters, we usually perform $12 \times 10^5 N$ MC moves in total, with the first $10^5 N$ moves for acceleration, the next $10^5 N$ moves for equilibrating the system, and the last $10^6 N$ moves for statistics, where N is the total number of mobile ions.

We introduce a parameter \tilde{l}_B to replace l_B in the definition of interaction (3.1) and dynamically change \tilde{l}_B in the first part of the moves, a total of $10^5 N$ of them, to speed up the thermal equilibration of the crowded system of particles. We generate a geometrical sequence of $10^5 N$ terms with the first and last terms being 1 and $l_B = 7 \text{ \AA}$, respectively. In the m th MC move with $m \leq 10^5 N$, the parameter \tilde{l}_B is taken to be the m th term in the geometrical sequence. After the first $10^5 N$ moves, we fix $\tilde{l}_B = l_B$ for all of the rest MC moves. We run another $10^5 N$ moves so that the system can reach an equilibrium.

Throughout the entire simulation, we keep the percentage of acceptance of MC moves between 20% and 50% by adaptively adjusting the value of the maximum length Δ_{\max} . Initially, we set $\Delta_{\max} = 2 \text{ \AA}$. We then change it after every 100 moves. If the acceptance rate is larger than 50% in current 100 moves, we increase Δ_{\max} by multiplying it by 1.05 but always keep

the new value of Δ_{\max} to be less than or equal to \tilde{l}_B . If the acceptance rate is smaller than 20% in current 100 moves, we decrease Δ_{\max} by multiplying it by 0.95, and we keep the new Δ_{\max} to be greater than $0.001 \tilde{l}_B$.

In the last part of the MC moves, a total of $10^6 N$ of them, we derive the production statistics and calculate the local radial particle density (RPD) for each ionic species. The RPD of the i th ionic species is defined by

$$\rho_i(r) = \frac{\langle N_i(r, r + \Delta r) \rangle}{\frac{4}{3}\pi[(r + \Delta r)^3 - r^3]}, \quad (3.2)$$

where $N_i(r, r + \Delta r)$ is the number of ions of the i th species whose centers are in the spherical shell between r and $r + \Delta r$, and the bracket $\langle \cdot \cdot \rangle$ represents an ensemble average over the shell. Notice that the denominator in the definition (3.2) is the volume of the shell. We choose Δr to be 1 Å. In our implementation, we approximate $\langle N_i(r, r + \Delta r) \rangle$ in (3.2) by the total number of ions of the i th species that move (in the last part of moves for statistics) into the shell between r and $r + \Delta r$, multiplied by the total number N_i of ions of i th species, divided by the total number of moves (in the last part of moves) in which an ion of i th species is displaced.

We remark that the use of periodic boundary condition effectively introduces a spatial cutoff of the underlying system region. In principle, this can affect the accuracy of the calculation of electrostatic interactions. However, we have tried simulations on boxes with different linear sizes and found almost no differences in the results. In fact, we find that on average only in one of 10000 moves does an ion have to “leave” through one side of the box and “come back” to the box through the opposite side. The reason for this is that most of the ions are crowded around the charged sphere, away from the boundary of simulation box.

In Fig. 1, we display our typical MC simulations results for two systems: one without salt and one with salt. Notice that

the counterions with smaller valence-to-volume ratios have less possibility to be adsorbed to the charged surface.

IV. RESULTS AND DISCUSSION

We set the linear size of our computational box $(-L/2, L/2)^3$ to be $L = 150$ Å and the radius of the spherical colloidal particle (the macroion) to be $R = 15$ Å. The Bjerrum length is set to be $l_B = 7$ Å. The surface charge density σ ranges from -0.05 to $-0.21 e/\text{Å}^2$. In our simulations, we investigate mixed solutions of three types of counterions, with their valences $(z_1, z_2, z_3) = (+1, +2, +3)$. We choose their radii to range from 1 Å to 4 Å. These are within the interval of physical interest. For example, the hydrated radii of monovalent hydrogen, sodium, and potassium, divalent magnesium and calcium, and trivalent aluminum ions are 4.5, 2.25, 1.5, 4.0, 3.0, and 4.5 Å, respectively [67]. For salt electrolytes, we choose monovalent or divalent coions with radius 2 Å.

One of the main objectives of our study is to understand the competitive adsorption of counterions with different valences and sizes. Such property has been already investigated previously (see Refs. [16,17,50,51,63,64,68] and the references therein). Most of these studies found that the valence of counterion determines the competition in adsorption to a charged surface with a low surface charge density and that smaller ions are stronger in such competition for a high surface charge density. These conclusions result naturally from the competition between electrostatic attraction and entropic repulsion expressed in the free-energy functional (2.3), where the electrostatics dominates the free energy for the low surface charge density, and the entropy dominates otherwise. It has been recently found in our previous work [37] using the mean-field model described in the last section that the competition between different ions in adsorption to a charged surface can be, in fact, characterized by the ionic valence-to-volume ratios. Here, we use MC simulations to further explore this characterization and compare our results




FIG. 1. (Color online) Typical MC simulations of ions surrounding a highly charged macroion. (a) A salt free system composed of monovalent (black), divalent (red, or dark gray in print version), and trivalent (green, or light gray in print version) counterions, with radii 3 Å, 2.5 Å, and 3.5 Å, respectively. (b) A system of salt solvent with coions (blue, or dark in print version), and counterions of valences and radii +1 and 2 Å (black), +2 and 3 Å (red, or dark gray in print version), and +3 and 4 Å (green, or light gray in print version), respectively.




FIG. 2. (Color online) The radial densities by MC simulations (a), (b), and (c) and by mean-field computations (d), (e), and (f) of the three species of counterions in Group 1: $(z_1, z_2, z_3) = (+1, +2, +3)$, $(R_1, R_2, R_3) = (3.0, 2.5, 3.5)$ in \AA , and $\alpha_{+1}/\alpha_{+2}/\alpha_{+3} = 1/3.5/1.9$. All three species have the same number of ions. This number is 100 in (a) and (d), 50 in (b) and (e), and 25 in (c) and (f). Hence, the constant surface charge density decreases from (a) and (d) to (b) and (e) and to (c) and (f).

with those from mean-field calculations. In particular, we study a system with a crowded ionic population near a highly charged surface, as shown in Fig. 1.

In what follows, for an ion of the i th species ($1 \leq i \leq M$), we denote by R_i its radius and by

$$\alpha_i = \frac{z_i}{v_i} = \frac{3z_i}{4\pi R_i^3}$$

its valence-to-volume ratio.

A. Crucial factors in the competition between counterions

We first study salt-free systems with monovalent, divalent, and trivalent counterions: $z_1 = +1$, $z_2 = +2$, and $z_3 = +3$. We investigate three different groups of such counterions with the following order of valence-to-volume ratios: $\alpha_{+2} > \alpha_{+3} > \alpha_{+1}$; $\alpha_{+3} > \alpha_{+1} > \alpha_{+2}$; and $\alpha_{+1} > \alpha_{+2} > \alpha_{+3}$. Here and below, we use α_{+i} to denote the valence-to-volume ratio of the counterion with valence $+i$. We use the following parameters:

Group 1 : $(R_1, R_2, R_3) = (3.0, 2.5, 3.5)$ in \AA ,

$$\alpha_{+1}/\alpha_{+2}/\alpha_{+3} = 1/3.5/1.9;$$

Group 2 : $(R_1, R_2, R_3) = (2.5, 3.5, 3.0)$ in \AA ,

$$\alpha_{+1}/\alpha_{+2}/\alpha_{+3} = 1.4/1/2.4;$$

Group 3 : $(R_1, R_2, R_3) = (2.0, 3.0, 4.0)$ in \AA ,

$$\alpha_{+1}/\alpha_{+2}/\alpha_{+3} = 2.7/1.6/1.$$

For each group, we choose the same number of ions for each of the three different species: $N_1 = N_2 = N_3$. Moreover,

we select three different surface charge densities by setting $N_1 = N_2 = N_3 = 100, 50$, and 25 , respectively, and by using the charge neutrality (2.1). The corresponding surface charge densities of the macroions are $-0.212, -0.106$, and $-0.053 e/\text{\AA}^2$, all in the regime of strong surface charge.

To report our MC simulations, we use bar plots, with each bar representing the radial density $\rho_i(r)$ for the i th ionic species ($1 \leq i \leq M$) as defined in (3.2). We choose the thickness of the spherical shell to be $\Delta r = 1 \text{\AA}$. We also convert the units number/volume to mol/L, which is abbreviated M . To show our results of mean-field computations, we plot smooth curves of radial densities $c_i(r)$ for the i th ionic species ($1 \leq i \leq M$) that are just the ionic concentrations in the radial direction. For each i the two quantities $\rho_i(r)$ and $c_i(r)$ should be close to each other, in particular, if the shell size Δr is very small. However, since the solvent molecules are not explicitly included in our MC simulations but the concentration of solvent molecules is included in our mean-field model, the two quantities $\rho_i(r)$ and $c_i(r)$ are not exactly the same.

The quantitative results of our MC simulations are illustrated in Figs. 2–4, where in Fig. 2 we also show our results of mean-field computations for comparison. Results in Fig. 2 are obtained using Group 1 parameters, while those in Fig. 3 and Fig. 4 are obtained using Group 2 and Group 3 parameters, respectively. We observe clearly that counterions are adsorbed tightly to the highly charged surface, and near the surface layers of counterions of different species form, leading to the remarkable structure of stratification. Moreover, we find that the order of layering depends on the valence-to-volume ratio,




FIG. 3. (Color online) The radial densities obtained by MC simulations of the three species of counterions in Group 2: $(z_1, z_2, z_3) = (+1, +2, +3)$, $(R_1, R_2, R_3) = (2.5, 3.5, 3.0)$ in Å, and $\alpha_{+1}/\alpha_{+2}/\alpha_{+3} = 1.4/1/2.4$. All three species have the same number of ions. This number is 100 (a), 50 (b), and 25 (c), respectively. Hence, the constant surface charge density decreases from (a) to (b) and to (c).

instead of the valence or the size independently. Counterions with the largest valence-to-volume ratio form the first layer closest to the charged surface, those with the second largest such ratio form the second layer, and so on. When the surface charge density σ becomes smaller, the role of valence is more important in determining which ionic species form a layer closest to the surface. These results demonstrate that the selective adsorption and layer ordering in the stratification depend on the competition between energetics and entropy, and that the valence-to-volume ratio is an important parameter in such adsorption and layering for a highly charged surface.

In Fig. 2, we find a qualitative agreement of our size-effect-included mean-field theory with the MC simulations. We note that the peaks of ionic densities close to the surface predicted by the MC simulations have a larger magnitude and are closer to the surface than those predicted by the mean-field theory.

We now investigate the sensitivity of the ionic sizes with respect to the ionic structure in the vicinity of a charged surface. We fix the surface charge density to be $\sigma = -0.22 e/\text{Å}^2$. We consider three species of counterions with valences $z_i = +i$ ($i = 1, 2, 3$) and number of ions $N_1 = N_2 = N_3 = 100$. In Fig. 5, we plot the ionic densities for various combinations of the ionic radii R_1 , R_2 , and R_3 . The radius of the trivalent

counterion is decreased from 4 Å in Fig. 5(a) to 3.5 Å in Fig. 5(b) so that the divalent and trivalent species have almost the same valence-to-volume ratio. The radius of the monovalent ion is decreased from 2 Å in Fig. 5(b) to 1.5 Å in Fig. 5(c). In both cases, the species with the highest valence-to-volume ratio, that is, the monovalent ionic species, remains the strongest in the competition to form the first layer closest to the charged surface. This indicates that the ionic competitive ability in adsorption is greatly improved by a slight decrease of its radius, which weakens the ionic entropic repulsion. From Fig. 5(b), we also find that when two species of counterions have close values of valence-to-volume ratios, the species with a higher valence will have a stronger ability of adsorption.

Figure 5(c) also illustrates an interesting phenomenon. With the decrease of the radius of monovalent ions, the concentration of the divalent ions, which have the second largest of the three valence-to-volume ratios, is increased. The trivalent ions which have the smallest valence-to-volume ratio are depleted in the vicinity of the surface. This can be interpreted that the more tightly binding of the monovalent ions to the surface decreases more the electrostatic energy contributed by divalent and trivalent ions. The effect of the valence-to-volume ratio is strengthened in the competition




FIG. 4. (Color online) The radial densities obtained by MC simulations of the three species of counterions in Group 3: $(z_1, z_2, z_3) = (+1, +2, +3)$, $(R_1, R_2, R_3) = (2.0, 3.0, 4.0)$ in Å, and $\alpha_{+1}/\alpha_{+2}/\alpha_{+3} = 2.7/1.6/1$. All three species have the same number of ions. This number is 100 in (a), 50 in (b), and 25 in (c). Hence, the constant surface charge density decreases from (a) to (b) and to (c).




FIG. 5. (Color online) MC simulations of the layering structure of ionic radial densities of three counterion species with the valences $z_1 = 1$, $z_2 = 2$, and $z_3 = 3$, and numbers of ions in each species $N_1 = N_2 = N_3 = 100$. The set of radii (R_1, R_2, R_3) is different in the three plots. The corresponding valence-to-volume ratios are (a) $\alpha_{+1}/\alpha_{+2}/\alpha_{+3} = 2.7/1.6/1$; (b) $\alpha_{+1}/\alpha_{+2}/\alpha_{+3} = 1.8/1.1/1$; (c) $\alpha_{+1}/\alpha_{+2}/\alpha_{+3} = 6.3/1.6/1$.

between the latter two species. This result demonstrates that the nonuniform ionic size effect plays a very important role in determining the properties of electrolyte solutions.

We now fix the numbers of ions $N_1 = N_2 = N_3 = 100$, the surface charge density $\sigma = -0.21 e/\text{\AA}^2$, the ratios of radii $R_1/R_2/R_3 = 2/3/4$, and the order of valence-to-volume ratios $\alpha_{+1} > \alpha_{+2} > \alpha_{+3}$. We vary simultaneously the ionic radii of the three species of counterions by changing a common multiplier. We use three different sets with the radii of monovalent ions being 1.2, 2.0, and 2.4 Å, respectively. We study how the different ionic sizes affect the layering structure of counterions and how the competition in ionic adsorption is changed with the change of entropy. The corresponding results are plotted in Fig. 6. It can be found that, with the increase of the ionic radii, the entropic contribution to the electrostatic free energy is increased, leading to the enhancement of the counterion repulsion. Moreover, the particle numbers of all the three ionic species in the layers closest to the surface are diminished. In the meantime, when ionic sizes are increased from small to large, the entropic contribution to the free energy becomes more significant. Hence, the valence-to-volume ratios give a clear characterization of stratification. In fact, the monovalent counterions, which have the smallest value of valence but the largest valence-to-volume ratio, always forms

the first layer closest to the surface. It is a further evidence that the competition between electrostatic energetics and entropy leads to the following limits: At the limit of the electrostatics domination the valence is the main indicator of the ordering of layers, while at the limit of the entropy domination the valence-to-volume ratio is the main indicator of layer ordering.

B. Systems with the presence of coions

We now add coions in the system and study the effect of coions to the competitive adsorption and order of packing of counterions, in comparison with the salt-free systems. We consider two cases. In the first case, we add monovalent coions to the system. We assume that the radius of such a coion is $R_4 = 2 \text{ \AA}$ and that the total number of coions is $N_4 = 204$. In the second case, we add divalent coions to the system. We assume that the radius of such a divalent coion is $R_4 = 2 \text{ \AA}$ and that the total number of such coions is $N_4 = 102$. In both cases, we still have the monovalent, divalent, and trivalent counterions, with now their radii 2, 3, and 4 Å, respectively, and their total numbers $N_1 = N_2 = N_3 = 134$. We also assume a high surface charge density $\sigma = -0.21 e/\text{\AA}^2$. The charge neutrality (2.1) is now satisfied with $M = 4$ species of




FIG. 6. (Color online) MC simulations of radial densities of three species of counterions with the valences $z_1 = 1$, $z_2 = 2$, and $z_3 = 3$, and numbers of ions in each species $N_1 = N_2 = N_3 = 100$. The radii R_1, R_2, R_3 in the three plots differ by a common factor. (a) Small ionic radii. (b) Medium ionic radii. (c) Large ionic radii.




FIG. 7. (Color online) MC simulations of counterion and coion radial densities. (a),(b) Systems with monovalent coions. (c),(d) Systems with divalent coions.

counterions and coions. The system will have an averaged 100 mM concentration of monovalent coions in the first case and 50 mM concentration of divalent coions in the second case.

For these two systems with coions, we plot the radial densities of counterions and coions obtained by our MC simulations in Fig. 7 and those obtained by our mean-field numerical computations in Fig. 8. In comparison with those salt free systems, we find that the addition of coions slightly enhances the layering effect. The densities of all three counterions are increased. This has a minor influence to their layering order. It is clear that a qualitative agreement between mean-field calculations and MC simulations is reached on the competition of counterion adsorption.

We observe from Fig. 7(b) and 7(d) that the coion density predicted by MC simulations is nonmonotonic, while from Figs. 8(b) and 8(d) that the coion density predicted by the mean-field theory is monotonic. In Fig. 9, we plot the total ionic charge density for each of the two systems obtained by our MC simulations. We find the overcharging of the system; that is, the total charge density is above zero, which corresponds to the charge neutrality [29,69]. Interestingly, the overcharging of the monovalent-coion system is stronger

than that in the divalent-coion system: The inverted charges of the monovalent-coion system and divalent-coion system are $1.86e$ and $0.15e$, respectively. This is mainly due to the fact that it is easier to form anion-cation binding pairs in the divalent-coion system than in the monovalent-coion system. Thus, the density of free counterions is decreased. In contrast, the mean-field theory can only produce a monotonic profile of the total charge density as proved mathematically in [26]. Therefore, the mean-field theory with the nonuniform size effect still fails in predicting the charge inversion.

V. CONCLUSIONS

In this work, we study the competition of multiple counterions of different valences and different sizes in binding to the surface of a spherical colloidal particle by both a mean-field theory and MC simulations. The parameters of the underlying system of electrolyte include the valences z_i , radii R_i (or volumes v_i), and numbers N_i of ions of i th species with $i = 1, \dots, M$, the radius R_0 of a solvent molecule, the constant dielectric coefficient ϵ of the electrolyte, and the surface charge density σ . The entire system is assumed to be in charge neutrality. In the mean-field approach, we




FIG. 8. (Color online) Counterion and coion radial densities from numerical computations based on the mean-field theory with the ionic size effect. (a) Radial densities of counterions for the system with monovalent coions. (b) Radial density of coions for the system with monovalent coions. (c) Radial densities of counterions for the system with divalent coions. (d) Radial density of coions for the system with divalent coions.

minimize a semiphenomenological electrostatic free-energy functional of ionic concentrations constrained by Poisson’s equation. The electrostatic potential is not an independent variable of the functional. The different ionic sizes are described through the entropic contributions of ions and solvent molecules. The constrained free-energy minimization

is realized numerically by an augmented Lagrange multiplier method. We also use an unrestricted primitive model and canonical ensemble MC simulations with the Metropolis criterion to predict the ionic distributions around the charged surface.

Through our extensive MC simulations and mean-field computations, we have found the following.

(1) For a low surface charge density, the adsorption of counterions with a higher valence is preferable. This agrees with previous studies in the existing literature. For a highly charged surface, both of the mean-field theory and MC simulations show that the counterions bind tightly around the charged surface, forming stratification or layering of counterions of different species.

(2) The ionic valence-to-volume ratios, instead of ionic valences alone, are the key parameters that determine the binding of counterions to the charged surface. Due to the ionic size effect, counterions with the largest valence-to-volume ratio form the first layer of stratification, while those with the second largest valence-to-volume ratio form the second layer, and so on. We call this the “criterion of valence-to-volume ratios” in ionic stratification. Our MC simulations confirm the validity of this criterion that was discovered in our previous mean-field calculations [37].




FIG. 9. (Color online) MC simulations of the total charge density for the system with monovalent coions (marked -1) and that with divalent coions (marked -2).

(3) Our MC simulations predict the charge inversion for ionic systems with salt. Moreover, we find that the overcharging is more significant for a system with monovalent cations than for a system with divalent cations. The mean-field theory, however, fails in predicting the charge inversion, since it does not include the ion-ion correlation.

In our mean-field computations, we have never found a case where our criterion of valence-to-volume ratios fails for the prediction of stratification of multiple counterions for highly charged surfaces. For MC simulations, we sometimes find the criterion does not work when those ratios are too close and the surface charge is too low. In fact, the MC simulation reported in Fig. 6(b) of Ref. [51] for a low surface charge density contradicts our criterion.

While our mean-field theory and MC simulations have both predicted the stratification of counterions near a highly charged surface and the crucial role of the ionic valence-to-volume ratios in such stratification, we have neglected several effects in our theory and methods.

First, in our MC simulations, we treat ions as hard spheres to describe the short-range repulsion in the van der Waals interactions between different kinds of ions of multiple valences and different sizes, and between the ions and the charged macroion. We have neglected the long-range attraction in such interactions that can contribute largely to the ion-ion correlations. For highly charged surfaces, counterions are crowded near the surface; and the van der Waals attraction may not be as strong as the corresponding repulsion. While we have taken a rather common approach in MC simulations, we understand that including the attraction part of the van der Waals interactions is practically quite possible. We include such interaction in our subsequent works.

Second, in both of our mean-field treatment and MC simulations, we use a uniform dielectric coefficient for the ionic solution. This is only an approximation in the description of the dielectric properties of solvent, as the water in the proximity of a highly charged surface is not expected to behave like bulk solvent. In fact, the dielectric coefficient can depend

on the ionic concentrations [13,70,71]. Such dependence is experimentally known to be continuous and linear (cf. Eq. (1) and Table I in Ref. [72]). Near the charged surface the dielectric coefficient is locally close to a constant; and the ion-ion interactions in such a region can be still modeled well by our interaction energy (3.1) but with a dielectric coefficient different from that in the bulk. We thus do not expect that this will significantly affect the competition of different counterions in the stratification. To further explore the detailed consequences of the concentration dependent dielectrics, we are currently extending our work to such dielectric systems.

Third, the size effect of solvent molecules is not directly included in our MC simulations. This makes our comparison between the mean-field theory and MC simulations only qualitative. There is clearly a need to develop models and algorithms to include the solvent molecular size effect in MC simulations of electrolyte systems.

We are currently working to improve our theory and methods to include some of these effects. In the future, it is desirable to apply our efficient theory and methods to large-scale modeling of biomolecular systems in which nonuniform ionic size effects can be sometimes very important. On the theoretical development, it is also necessary to derive from statistical mechanics theory our mean-field, electrostatic free-energy functional that includes the nonuniform ionic size effect.

ACKNOWLEDGMENTS

This work was supported by the US National Science Foundation (NSF) through Grant no. DMS-0811259 (B.L. and J.W.), the US NSF Center for Theoretical Biological Physics (CTBP) through Grant No. PHY-0822283 (B.L. and S.Z.), the National Institutes of Health through the Grant No. R01GM096188 (B.L.), the Natural Science Foundation of China through Grant No. NSFC-11101276 (Z.X.), the Chinese Ministry of Education through Grant No. NCET-09-0556 (Z.X.), and the China Scholarship Council (S.Z.).

-
- [1] D. Andelman, in *Handbook of Biological Physics*, edited by R. Lipowsky and E. Sackmann (Elsevier, Amsterdam, 1995), Vol. 1, pp. 603–642.
- [2] H. Boroudjerdi, Y.-W. Kim, A. Naji, R. R. Netz, X. Schlagberger, and A. Serr, *Phys. Rep.* **416**, 129 (2005).
- [3] M. E. Davis and J. A. McCammon, *Chem. Rev.* **90**, 509 (1990).
- [4] F. Fogolari, A. Brigo, and H. Molinari, *J. Mol. Recognit.* **15**, 377 (2002).
- [5] R. H. French, V. A. Parsegian, R. Podgornik, R. F. Rajter, A. Jagota, J. Luo, D. Asthagiri, M. K. Chaudhury, Y.-M. Chiang, S. Granick, S. Kalinin, M. Kardar, R. Kjellander, D. C. Langreth, J. Lewis, S. Lustig, D. Wesolowski, J. S. Wettlaufer, W.-Y. Ching, M. Finnis, F. Houlihan, O. A. von Lilienfeld, C. J. van Oss, and T. Zemb, *Rev. Mod. Phys.* **82**, 1887 (2010).
- [6] Y. Levin, *Rep. Prog. Phys.* **65**, 1577 (2002).
- [7] J. A. McCammon, *Proc. Natl. Acad. Sci. USA* **106**, 7683 (2009).
- [8] K. A. Sharp and B. Honig, *Annu. Rev. Biophys. Biophys. Chem.* **19**, 301 (1990).
- [9] D. Lambert, D. Leipply, R. Shiman, and D. E. Draper, *J. Mol. Biol.* **390**, 791 (2009).
- [10] Y. Bai, K. Travers, V. B. Chu, J. Lipfert, S. Doniach, and D. Herschlag, *J. Am. Chem. Soc.* **129**, 14981 (2007).
- [11] M. L. Bleam, C. F. Anderson, and M. T. J. Record, *Proc. Natl. Acad. Sci. USA* **77**, 3085 (1980).
- [12] B. Eisenberg, In *Advances in Chemical Physics*, edited by S. A. Rice and A. R. Dinner (Wiley, New York, 2011), Vol. 148, p. 77.
- [13] J. J. Howard, J. S. Perkyns, and B. M. Pettitt, *J. Phys. Chem. B* **114**, 6074 (2010).
- [14] A. A. Kornyshev, *J. Phys. Chem. B* **111**, 5545 (2007).
- [15] M. Quesada-Pérez, A. Martín-Molina, and R. Hidalgo-Álvarez, *J. Chem. Phys.* **121**, 8618 (2004).
- [16] M. Valiskó, D. Boda, and D. Gillespie, *J. Phys. Chem. C* **111**, 15575 (2007).

- [17] K. Wang, Y.-X. Yu, G.-H. Gao, and G.-S. Luo, *J. Chem. Phys.* **123**, 234904 (2005).
- [18] D. L. Chapman, *Philos. Mag.* **25**, 475 (1913).
- [19] P. Debye and E. Hückel, *Phys. Z.* **24**, 185 (1923).
- [20] F. Fixman, *J. Chem. Phys.* **70**, 4995 (1979).
- [21] G. Gouy, *J. Phys. Théor. Appl.* **9**, 457 (1910).
- [22] P. Grochowski and J. Trylska, *Biopolymers* **89**, 93 (2008).
- [23] B. Z. Lu, Y. C. Zhou, M. J. Holst, and J. A. McCammon, *Commun. Comput. Phys.* **3**, 973 (2008).
- [24] J. Che, J. Dzubiella, B. Li, and J. A. McCammon, *J. Phys. Chem. B* **112**, 3058 (2008).
- [25] F. Fogolari and J. M. Briggs, *Chem. Phys. Lett.* **281**, 135 (1997).
- [26] B. Li, *Nonlinearity* **22**, 811 (2009).
- [27] B. Li, *SIAM J. Math. Anal.* **40**, 2536 (2009).
- [28] E. S. Reiner and C. J. Radke, *J. Chem. Soc. Faraday Trans.* **86**, 3901 (1990).
- [29] A. Y. Grosberg, T. T. Nguyen, and B. I. Shklovskii, *Rev. Mod. Phys.* **74**, 329 (2002).
- [30] V. Vlachy, *Annu. Rev. Phys. Chem.* **50**, 145 (1999).
- [31] I. Borukhov, D. Andelman, and H. Orland, *Phys. Rev. Lett.* **79**, 435 (1997).
- [32] I. Borukhov, D. Andelman, and H. Orland, *Electrochim. Acta* **46**, 221 (2000).
- [33] A. H. Boschitsch and P. V. Danilov, *J. Comput. Chem.* **33**, 1152 (2012).
- [34] V. B. Chu, Y. Bai, J. Lipfert, D. Herschlag, and S. Doniach, *Biophys. J.* **93**, 3202 (2007).
- [35] V. Kralj-Iglič and A. Iglič, *J. Phys. II (France)* **6**, 477 (1996).
- [36] G. Tresset, *Phys. Rev. E* **78**, 061506 (2008).
- [37] S. Zhou, Z. Wang, and B. Li, *Phys. Rev. E* **84**, 021901 (2011).
- [38] D. Antypov, M. C. Barbosa, and C. Holm, *Phys. Rev. E* **71**, 061106 (2005).
- [39] M. Bazant, B.D. Storey, and A. A. Kornyshev, *Phys. Rev. Lett.* **106**, 046102 (2011).
- [40] M. Z. Bazant, M. S. Kilic, B. D. Storey, and A. Ajdari, *Adv. Colloid Interface Sci.* **152**, 48 (2009).
- [41] J. J. Bikerman, *Philos. Mag.* **33**, 384 (1942).
- [42] B. Eisenberg, Y.-K. Hyon, and C. Liu, *J. Chem. Phys.* **133**, 104104 (2010).
- [43] M. S. Kilic, M. Z. Bazant, and A. Ajdari, *Phys. Rev. E* **75**, 021502 (2007).
- [44] M. S. Kilic, M. Z. Bazant, and A. Ajdari, *Phys. Rev. E* **75**, 021503 (2007).
- [45] B. Lu and Y. Zhou, *Biophys. J.* **100**, 2475 (2011).
- [46] A. R. J. Silalahi, A. H. Boschitsch, R. C. Harris, and M. O. Fenley, *J. Chem. Theory Comput.* **6**, 3631 (2010).
- [47] M. Baptista, R. Schmitz, and B. Dunweg, *Phys. Rev. E* **80**, 016705 (2009).
- [48] A. C. Maggs and V. Rossetto, *Phys. Rev. Lett.* **88**, 196402 (2002).
- [49] A. Tulpar, P. R. Van Tassel, and J. Y. Walz, *Langmuir* **22**, 2876 (2006).
- [50] C. Avendaño and A. Gil-Villegas, *Mol. Phys.* **104**, 1475 (2006).
- [51] P. Taboada-Serrano, S. Yiacoumi, and C. Tsouris, *J. Chem. Phys.* **123**, 054703 (2005).
- [52] D. Frenkel and B. Smit, *Understanding Molecular Simulation: From Algorithms to Applications* (Academic Press, New York, 2002).
- [53] P. Linse, *Adv. Polym. Sci.* **185**, 111 (2005).
- [54] T. Goel, C. N. Patra, S. K. Ghosh, and T. Mukherjee, *J. Phys. Chem. B* **115**, 10903 (2011).
- [55] Z. Y. Wang and Y. Q. Ma, *J. Chem. Phys.* **133**, 064704 (2010).
- [56] Z. Gan and Z. Xu, *Phys. Rev. E* **84**, 016705 (2011).
- [57] R. Kjellander and S. Marčelja, *Chem. Phys. Lett.* **112**, 49 (1984).
- [58] R. Messina, *J. Chem. Phys.* **117**, 11062 (2002).
- [59] G. M. Torrie, J. P. Valleau, and G. N. Patey, *J. Chem. Phys.* **76**, 4615 (1982).
- [60] Z. Y. Wang and Y. Q. Ma, *J. Chem. Phys.* **131**, 244715 (2009).
- [61] D. P. Bertsekas, *Constrained Optimization and Lagrange Multiplier Method* (Academic Press, New York, 1982).
- [62] J. Nocedal and S. Wright, *Numerical Optimization* (Springer-Verlag, New York, 1999).
- [63] Z. Abbas, E. Ahlberg, and S. Nordholm, *J. Phys. Chem. B* **113**, 5905 (2009).
- [64] M. Bešter and V. Vlachy, *J. Chem. Phys.* **96**, 7656 (1992).
- [65] N. Metropolis, A. W. Rosenbluth, M. N. Rosenbluth, A. H. Teller, and E. Teller, *J. Chem. Phys.* **21**, 1087 (1953).
- [66] J. P. Valleau and L. K. Cohen, *J. Chem. Phys.* **72**, 5935 (1980).
- [67] J. Kielland, *J. Am. Chem. Soc.* **59**, 1675 (1937).
- [68] G. I. Guerrero-Garcia, E. Gonzalez-Tovar, and M. O. de la Cruz, *J. Chem. Phys.* **135**, 054701 (2011).
- [69] M. Quesada-Pérez, E. González-Tovar, A. Martín-Molina, M. Lozada-Cassou, and R. Hidalgo-Álvarez, *Chem. Phys. Chem.* **4**, 234 (2003).
- [70] M. Eigen and E. Wicke, *J. Phys. Chem.* **58**, 702 (1954).
- [71] I. Kalcher, J. C. F. Schulz, and J. Dzubiella, *J. Chem. Phys.* **133**, 164511 (2011).
- [72] D. Ben-Yaakov, D. Andelman, and R. Podgornik, *J. Chem. Phys.* **134**, 074705 (2011).