

**Math 270C: Numerical Mathematics (Part C)**  
**LECTURE NOTES**

**Bo Li**  
**Department of Mathematics**  
**University of California, San Diego**

**June 10, 2009**

**Warning!**

- While being expanded with the addition of new material and being carefully polished continuously, these notes may still contain many typos and mistakes.
- These notes are mainly for the graduate course Math 270C, Spring quarter, 2009, at UC San Diego. Any individuals are however welcome to use these notes for personal studies and classes. As always, any comments are very much appreciated.
- Please do not copy these notes partially or entirely for publication without permission of the author. Please do not distribute these notes.

**Acknowledgments**

Many thanks to Yi Ling Ng for typing out the first version of the latex file of these notes, and to the graduate students of Math 270C, Spring quarter, 2007, 2008, and 2009, for their valuable comments on these notes.



# Contents

<b>1</b>	<b>Polynomial Approximation</b>	<b>5</b>
1.1	The Weierstrass Theorem . . . . .	5
1.2	Best Uniform Approximation . . . . .	9
1.3	Chebyshev Polynomials . . . . .	14
1.4	Uniform Approximation by Trigonometric Polynomials . . . . .	17
1.5	Modulus of Continuity and Jackson's Theorems . . . . .	17
1.6	Least-Squares Approximation . . . . .	17
1.7	Orthogonal Polynomials . . . . .	21
1.8	More Properties of Orthogonal Polynomials . . . . .	25
1.9	Legendre Polynomials . . . . .	30
	Exercises . . . . .	33
<b>2</b>	<b>Polynomial Interpolation</b>	<b>41</b>
2.1	Lagrange Interpolation . . . . .	41
2.2	Newton's Formula and Divided Differences . . . . .	46
2.3	Peano Kernel and the Remainder Theorem . . . . .	52
2.4	Hermite Interpolation and Divided Differences with Repeated Points . . . . .	56
2.5	Convergence of Interpolation Polynomials . . . . .	58
2.6	Piecewise Polynomial Interpolation . . . . .	62
2.7	Cubic Splines . . . . .	62
2.8	Trigonometric Interpolation and Fast Fourier Transforms . . . . .	62
	Exercises . . . . .	62
<b>3</b>	<b>Numerical Integration</b>	<b>69</b>
3.1	The Basics . . . . .	69
3.2	Interpolatory Quadrature . . . . .	74
3.3	Peano Kernel and Error Representation . . . . .	81
3.4	Euler–Maclaurin Formula, Richardson Extrapolation, and Romberg Algorithm . . . . .	82
3.5	Weighted Gaussian Quadrature . . . . .	82
3.6	Convergence of Sequences of Numerical Quadrature . . . . .	88
	Exercises . . . . .	90



# Chapter 1

## Polynomial Approximation

### 1.1 The Weierstrass Theorem

Let  $a, b \in \mathbb{R}$  with  $a < b$ . Let  $C[a, b]$  denote the set of all continuous, real-valued functions on the closed interval  $[a, b]$ . Let  $\mathcal{P}$  denote the set of all real polynomials. For each integer  $n \geq 0$ , let  $\mathcal{P}_n$  denote the set of all real polynomials of degree less than or equal to  $n$ .

**Theorem 1.1 (The First Weierstrass Approximation Theorem).** *Let  $f \in C[a, b]$ . For any  $\varepsilon > 0$ , there exists  $p \in \mathcal{P}$  such that*

$$|f(x) - p(x)| < \varepsilon \quad \forall x \in [a, b]. \quad (1.1)$$

To prove Theorem 1.1, we define for any given  $f \in C[0, 1]$

$$(B_0 f)(x) = f(0) \quad \forall x \in [0, 1],$$
$$(B_n f)(x) = \sum_{k=0}^n f\left(\frac{k}{n}\right) \binom{n}{k} x^k (1-x)^{n-k} \quad \forall x \in [0, 1] \quad \forall n \geq 1,$$

where

$$\binom{n}{k} = \frac{n!}{k!(n-k)!}.$$

For each  $n \geq 0$ , we call  $B_n f$  the  $n$ th Bernstein polynomial of  $f \in C[0, 1]$ . Clearly,  $B_n f \in \mathcal{P}_n$  for each  $n$ .

**Lemma 1.2 (Properties of Bernstein polynomials).**

(1) For each  $n \geq 0$ ,  $B_n : C[0, 1] \rightarrow \mathcal{P}_n$  is linear, i.e.,

$$B_n(f + g) = B_n f + B_n g \quad \forall f, g \in C[0, 1],$$
$$B_n(\alpha f) = \alpha B_n f \quad \forall \alpha \in \mathbb{R} \quad \forall f \in C[0, 1].$$

(2) For each  $n \geq 0$ ,  $B_n : C[0, 1] \rightarrow \mathcal{P}_n$  is non-negative, i.e.,  $(B_n f)(x) \geq 0$  for any  $x \in [0, 1]$  provided that  $f(x) \geq 0$  for any  $x \in [0, 1]$ .

(3) Let  $p_i(x) = x^i$  ( $i = 1, 2, 3$ ). Then,

$$(B_n p_0)(x) = p_0(x) \quad \forall x \in [0, 1] \quad \forall n \geq 0, \quad (1.2)$$

$$(B_n p_1)(x) = p_1(x) \quad \forall x \in [0, 1] \quad \forall n \geq 1, \quad (1.3)$$

$$(B_n p_2)(x) = p_2(x) + \frac{1}{n}x(1-x) \quad \forall x \in [0, 1] \quad \forall n \geq 2. \quad (1.4)$$

*Proof.* Part (1) and Part (2) are obvious. To prove Part (3), we use the binomial formula:

$$(\alpha + \beta)^n = \sum_{k=0}^n \binom{n}{k} \alpha^k \beta^{n-k} \quad \forall \alpha, \beta \in \mathbb{R}. \quad (1.5)$$

Let  $x \in [0, 1]$ . We have by (1.5) that for  $n \geq 1$

$$(B_n p_0)(x) = \sum_{k=0}^n \binom{n}{k} x^k (1-x)^{n-k} = (x + (1-x))^n = 1 = p_0(x).$$

By definition,  $(B_0 p_0)(x) = p_0(x) = 1$  for all  $x \in [0, 1]$ . Thus, (1.2) is true.

Let  $n \geq 1$ . Using (1.5), we obtain for any  $x \in [0, 1]$  that

$$\begin{aligned} (B_n p_1)(x) &= \sum_{k=0}^n p_1 \left( \frac{k}{n} \right) \binom{n}{k} x^k (1-x)^{n-k} \\ &= \sum_{k=0}^n \frac{k}{n} \frac{n!}{k!(n-k)!} x^k (1-x)^{n-k} \\ &= x \sum_{k=1}^n \frac{(n-1)!}{(k-1)!((n-1)-(k-1))!} x^{k-1} (1-x)^{(n-1)-(k-1)} \\ &\stackrel{j=k-1}{=} x \sum_{j=0}^{n-1} \binom{n-1}{j} x^{n-1-j} (1-x)^{n-1-j} \\ &= x(x + (1-x))^{n-1} \\ &= p_1(x). \end{aligned}$$

This is (1.3). The proof of (1.4) is left as an exercise. □

**Lemma 1.3 (Convergence of Bernstein polynomials).** *We have*

$$\lim_{n \rightarrow \infty} \max_{0 \leq x \leq 1} |(B_n f)(x) - f(x)| = 0 \quad \forall f \in C[0, 1].$$

*Proof.* Let  $f \in C[0, 1]$ . Let  $\varepsilon > 0$ . We show in three steps that

$$\max_{0 \leq x \leq 1} |(B_n f)(x) - f(x)| < \varepsilon, \quad (1.6)$$

if  $n$  is sufficiently large.

*Step 1.* Let  $M = \max_{0 \leq x \leq 1} |f(x)|$ . Then,

$$-2M \leq f(x) - f(y) \leq 2M \quad \forall x, y \in [0, 1]. \quad (1.7)$$

Since  $f \in C[0, 1]$ ,  $f$  is uniformly continuous on  $[0, 1]$ . Thus, there exists  $\delta > 0$  such that

$$-\frac{\varepsilon}{2} < f(x) - f(y) < \frac{\varepsilon}{2} \quad \forall x, y \in [0, 1] \text{ with } |x - y| < \delta. \quad (1.8)$$

We claim that

$$-\frac{\varepsilon}{2} - \frac{2M}{\delta^2}(x - y)^2 \leq f(x) - f(y) \leq \frac{\varepsilon}{2} + \frac{2M}{\delta^2}(x - y)^2 \quad \forall x, y \in [0, 1]. \quad (1.9)$$

In fact, if  $|x - y| < \delta$ , then (1.8) implies (1.9). If  $|x - y| \geq \delta$ , then  $((x - y)/\delta)^2 \geq 1$ , and hence, (1.7) implies (1.9).

*Step 2.* Fix  $y$  in (1.9). Apply  $B_n$  ( $n \geq 1$ ) to each side of (1.9) as a continuous function of  $x$ . Using the properties of  $B_n$  (cf. Lemma 1.2), we obtain

$$-\frac{\varepsilon}{2} - \frac{2M}{\delta^2}(B_n(x - y)^2)(x) \leq (B_n f)(x) - f(y) \leq \frac{\varepsilon}{2} + \frac{2M}{\delta^2}(B_n(x - y)^2)(x). \quad (1.10)$$

Since

$$(x - y)^2 = x^2 - 2xy + y^2 = p_2(x) - 2yp_1(x) + y^2p_0(x),$$

we have again by Lemma 1.2 that

$$\begin{aligned} (B_n(x - y)^2)(x) &= (B_n p_2)(x) - 2y(B_n p_1)(x) + y^2(B_n p_0)(x) \\ &= x^2 + \frac{1}{n}x(1 - x) - 2yx + y^2 \\ &= (x - y)^2 + \frac{1}{n}x(1 - x). \end{aligned}$$

This and (1.10) lead to

$$-\frac{\varepsilon}{2} - \frac{2M}{\delta^2} \left[ (x - y)^2 + \frac{x(1 - x)}{n} \right] \leq (B_n f)(x) - f(y) \leq \frac{\varepsilon}{2} + \frac{2M}{\delta^2} \left[ (x - y)^2 + \frac{x(1 - x)}{n} \right] \quad \forall x, y \in [0, 1]. \quad (1.11)$$

*Step 3.* Setting  $y = x$  in (1.11), we obtain

$$|(B_n f)(x) - f(x)| \leq \frac{\varepsilon}{2} + \frac{2M}{\delta^2} \frac{x(1 - x)}{n} \leq \frac{\varepsilon}{2} + \frac{2M}{\delta^2 n} \quad \forall x \in [0, 1].$$

Therefore, (1.6) holds true for all  $n > 4M/(\delta^2\varepsilon)$ .  $\square$

*Proof of the First Weierstrass Approximation Theorem.* Define  $g(t) = f((b-a)t + a)$  ( $0 \leq t \leq 1$ ). Then,  $g \in C[0, 1]$ . Hence, by Lemma 1.3, there exists  $q \in \mathcal{P}$  such that

$$|g(t) - q(t)| < \varepsilon \quad \forall t \in [0, 1].$$

Let  $p(x) = q((x-a)/(b-a))$ . Then,  $p \in \mathcal{P}$ , and

$$|f(x) - p(x)| = |g(t) - q(t)| < \varepsilon \quad \forall x \in [a, b].$$

This is (1.1). □

We define for any  $f \in C[a, b]$ ,

$$\|f\|_{C[a,b]} = \max_{a \leq x \leq b} |f(x)|.$$

When no confusion arises, we also write  $\|f\|$  instead of  $\|f\|_{C[a,b]}$ . One easily verifies that  $\|\cdot\|$  is a norm of the vector space  $C[a, b]$ , i.e., the following hold:

- (1)  $\|f\| \geq 0 \quad \forall f \in C[a, b]$ .  
 $\|f\| = 0$  if and only if  $f(x) = 0 \quad \forall x \in [a, b]$ ;
- (2)  $\|\alpha f\| = |\alpha| \|f\| \quad \forall \alpha \in \mathbb{R} \quad \forall f \in C[a, b]$ ;
- (3)  $\|f + g\| \leq \|f\| + \|g\| \quad \forall f, g \in C[a, b]$ .

The last inequality is called the *triangle inequality*. It implies that

$$|\|f\| - \|g\|| \leq \|f - g\| \quad \forall f, g \in C[a, b]. \quad (1.12)$$

This norm is called the *maximum norm* or  $C[a, b]$ -*norm*.

With the notion of maximum norm, the Weierstrass Theorem states exactly that, for any  $f \in C[a, b]$  and any  $\varepsilon > 0$ , there exists  $p \in \mathcal{P}$  such that  $\|f - p\| < \varepsilon$ . Equivalently, for any  $f \in C[a, b]$ , there exists a sequence of polynomials  $\{p^{(k)}\}_{k=1}^{\infty}$  such that  $p^{(k)} \rightarrow f$  in  $C[a, b]$ , i.e.,  $\|f - p^{(k)}\| \rightarrow 0$  as  $k \rightarrow \infty$ .

The proof of Lemma 1.3 can be extended to the proof of the following Bohman–Korovkin Theorem for more general sequences of linear, non-negative operators; and clearly the Bohman–Korovkin Theorem and the properties of Bernstein polynomials imply the First Weierstrass Approximation Theorem:

**Theorem 1.4 (Bohman–Korovkin).** *For each integer  $n \geq 1$ , let  $L_n : C[a, b] \rightarrow \mathcal{P}_n$  be an operator. Assume:*

- (1) *Each  $L_n : C[a, b] \rightarrow \mathcal{P}_n$  is linear;*
- (2) *Each  $L_n : C[a, b] \rightarrow \mathcal{P}_n$  is non-negative;*
- (3)  *$L_n p_k \rightarrow p_k$  in  $C[a, b]$  for  $k = 0, 1, 2$ , where  $p_k(x) = x^k$ .*

*Then  $\|L_n f - f\| \rightarrow 0$  for any  $f \in C[a, b]$ .* □

## 1.2 Best Uniform Approximation

We denote for any  $f \in C[a, b]$

$$E_n(f) = \inf_{p \in \mathcal{P}_n} \|f - p\| \quad \forall n \geq 0. \quad (1.13)$$

Here, the norm  $\|\cdot\|$  is the  $C[a, b]$ -norm. Clearly,  $0 \leq E_n(f) \leq \|f\|$  for all  $n \geq 0$ . Moreover, since  $\mathcal{P}_0 \subset \cdots \subset \mathcal{P}_n \subset \cdots$ , we have  $E_0(f) \geq \cdots \geq E_n(f) \geq \cdots$ .

**Proposition 1.5.** *The First Weierstrass Approximation Theorem is equivalent to*

$$\lim_{n \rightarrow \infty} E_n(f) = 0 \quad \forall f \in C[a, b]. \quad \square$$

**Definition 1.6 (Best uniform approximation).** *A best uniform approximation of a given function  $f \in C[a, b]$  in  $\mathcal{P}_n$  is a polynomial  $p_n \in \mathcal{P}_n$  that satisfies*

$$\|f - p_n\| = \min_{q_n \in \mathcal{P}_n} \|f - q_n\|. \quad (1.14)$$

Since (1.14) can be written as

$$\max_{a \leq x \leq b} |f(x) - p_n(x)| = \min_{q_n \in \mathcal{P}_n} \max_{a \leq x \leq b} |f(x) - q_n(x)|,$$

a best uniform approximation is also called a *minimax approximation*.

**Theorem 1.7 (Existence of best uniform approximation).** *For any  $f \in C[a, b]$  and any integer  $n \geq 0$ , there exists a best uniform approximation of  $f$  in  $\mathcal{P}_n$ .*

*Proof.* Let  $f \in C[a, b]$  and fix an integer  $n \geq 0$ . For any  $c = (c_0, \dots, c_n) \in \mathbb{R}^{n+1}$ , we associate with a unique polynomial  $p_c \in \mathcal{P}_n$  by  $p_c(x) = \sum_{k=0}^n c_k x^k$ . Define  $F : \mathbb{R}^{n+1} \rightarrow \mathbb{R}$  by

$$F(c) = \|f - p_c\| = \max_{a \leq x \leq b} \left| f(x) - \sum_{k=0}^n c_k x^k \right|.$$

Clearly, the assertion of the theorem is equivalent to the existence of  $c \in \mathbb{R}^{n+1}$  such that

$$F(c) = \min_{d \in \mathbb{R}^{n+1}} F(d). \quad (1.15)$$

*Step 1.* We show that the function  $F : \mathbb{R}^{n+1} \rightarrow \mathbb{R}$  satisfies the following properties:

- (1)  $F : \mathbb{R}^{n+1} \rightarrow \mathbb{R}$  is continuous;
- (2)  $F(c) \rightarrow +\infty$  as  $\|c\| \rightarrow \infty$ .

By (1.12), we have

$$\begin{aligned}
|F(c) - F(d)| &= \left| \|f - p_c\| - \|f - p_d\| \right| \leq \|p_d - p_c\| \\
&= \max_{a \leq x \leq b} \left| \sum_{k=0}^n (d_k - c_k) x^k \right| \leq \sum_{k=0}^n |d_k - c_k| \max\{|a|^k, |b|^k\} \\
&\rightarrow 0 \quad \text{as } d \rightarrow c \text{ in } \mathbb{R}^{n+1}.
\end{aligned}$$

This proves Part (1).

To prove Part (2), we define  $G : \mathbb{R}^{n+1} \rightarrow \mathbb{R}$  by  $G(c) = \|p_c\|$  for any  $c \in \mathbb{R}^{n+1}$ . As proved in Part (1),  $G : \mathbb{R}^{n+1} \rightarrow \mathbb{R}$  is continuous. Let

$$\mathbb{S}^n = \{c \in \mathbb{R}^{n+1} : \|c\| = 1\},$$

where

$$\|c\| = \sqrt{\sum_{k=0}^n c_k^2} \quad \forall c = (c_0, \dots, c_n) \in \mathbb{R}^{n+1}.$$

Clearly,  $\mathbb{S}^n$  is a closed and bounded subset, and hence a compact subset, of  $\mathbb{R}^{n+1}$ . Moreover,  $G(c) > 0$  for any  $c \in \mathbb{S}^n$ . The continuity of  $G : \mathbb{R}^{n+1} \rightarrow \mathbb{R}$  then implies that

$$\mu := \inf_{c \in \mathbb{S}^n} G(c) > 0. \quad (1.16)$$

Now, let  $c \in \mathbb{R}^{n+1}$  with  $c \neq 0$ . Set  $\hat{c} := (1/\|c\|)c \in \mathbb{S}^n$ . We have

$$p_c(x) = \sum_{k=0}^n c_k x^k = \|c\| \sum_{k=0}^n \frac{c_k}{\|c\|} x^k = \|c\| p_{\hat{c}}(x) \quad \forall x \in [a, b],$$

i.e.,  $p_c = \|c\| p_{\hat{c}}$ . Therefore, by (1.12) and (1.16), we have

$$\begin{aligned}
F(c) &\geq \|p_c\| - \|f\| = \| \|c\| p_{\hat{c}} \| - \|f\| \\
&= \|c\| G(p_{\hat{c}}) - \|f\| \geq \|c\| \mu - \|f\| \rightarrow +\infty \quad \text{as } \|c\| \rightarrow \infty,
\end{aligned}$$

proving Part (2).

*Step 2.* Clearly  $F(c) \geq 0$  for any  $c \in \mathbb{R}^{n+1}$ . Let  $m = \inf_{c \in \mathbb{R}^{n+1}} F(c) \geq 0$ . Since  $\lim_{c \rightarrow \infty} F(c) = +\infty$ , there exists  $R > 0$  such that  $F(c) \geq m + 1$  if  $\|c\| > R$ . Therefore,  $m = \inf_{\|c\| \leq R} F(c)$ . Since  $F$  is continuous on the bounded and closed subset  $B_R := \{c \in \mathbb{R}^{n+1} : \|c\| \leq R\}$ , it attains its minimum over  $B_R$ , and hence over the entire  $\mathbb{R}^{n+1}$ .  $\square$

**Theorem 1.8 (The Chebyshev Alternation Theorem).** *Let  $f \in C[a, b]$  and  $f \notin \mathcal{P}_n$ . Then,  $p \in \mathcal{P}_n$  is a best uniform approximation if and only if  $f - p$  achieves its maximum magnitude at  $n + 2$  points with alternating signs, i.e., there exist  $n + 2$  points  $x_k$  ( $1 \leq k \leq n + 2$ ) with  $a \leq x_1 < \dots < x_{n+2} \leq b$  such that*

$$\begin{aligned}
|f(x_k) - p(x_k)| &= \|f - p\|, \quad k = 1, \dots, n + 2, \\
[f(x_k) - p(x_k)][f(x_{k+1}) - p(x_{k+1})] &< 0, \quad k = 1, \dots, n + 1.
\end{aligned}$$

**Example.** Consider  $f(x) = x^2$  with  $x \in [0, 1]$  and  $n = 1$ . Let  $p_1 \in \mathcal{P}_1$  be the best uniform approximation of  $f$  over the interval  $[0, 1]$ . Assume  $p_1(x) = \alpha x + \beta$  with  $\alpha, \beta$  two constants. Let  $g(x) = f(x) - p_1(x)$  for any  $x \in [0, 1]$ . By the Chebyshev Alternation Theorem, there exist  $x_1, x_2, x_3 \in [0, 1]$  with  $0 \leq x_1 < x_2 < x_3 \leq 1$  such that  $|g(x_i)| = \|g\|_{C[0,1]}$ ,  $i = 1, 2, 3$ , and the sign of  $g(x_1), g(x_2), g(x_3)$  alternates.

Notice that  $g'(x) = 2x - \alpha$  which has only one root inside  $(0, 1)$ . Since  $x_2$  is an interior extrem point of  $g(x)$ , we must have  $g'(x_2) = 0$ , leading to  $x_2 = \alpha/2$ . Moreover, the other maximal or minimal points  $x_1$  and  $x_3$  must be the boundary points:  $x_1 = 0$  and  $x_3 = 1$ .

Now by the fact that  $g(x_1) = g(x_3)$ , i.e.,  $g(0) = g(1)$ , we obtain that  $-\beta = 1 - \alpha - \beta$ . Hence  $\alpha = 1$  and  $x_2 = 1/2$ . By the fact that  $g(x_1) = -g(x_2)$ , i.e.,  $g(0) = -g(1/2)$ , we obtain  $-\beta = -1/4 + (1/2 + \beta)$ . Hence  $\beta = -1/8$ . Therefore  $p_1(x) = x - 1/8$ . We also have that  $\|g\|_{C[0,1]} = g(0) = 1/8$ .

To summarize, we have

$$\min_{p \in \mathcal{P}_1} \|f - p\|_{C[0,1]} = \min_{\gamma, \delta \in \mathbb{R}} \max_{0 \leq x \leq 1} |x^2 - (\gamma x + \delta)| = \max_{0 \leq x \leq 1} \left| x^2 - \left( x - \frac{1}{8} \right) \right| = \frac{1}{8}.$$

*Proof of the Chebyshev Alternation Theorem.* Notice that  $p_n \in \mathcal{P}_n$  is the best uniform approximation of  $f \in C[a, b]$  in  $\mathcal{P}_n$  if and only if  $0 \in \mathcal{P}_n$  is the best uniform approximation of  $f - p_n \in C[a, b]$  in  $\mathcal{P}_n$ . Therefore, the Chebyshev Alternation Theorem is equivalent to the following statement:

*Let  $f \in C[a, b]$  but  $f \notin \mathcal{P}_n$ . Then, the zero polynomial  $0 \in \mathcal{P}_n$  is a best uniform approximation of  $f$  in  $\mathcal{P}_n$  if and only if  $f$  achieves its maximum magnitude at  $n + 2$  points in  $[a, b]$  with alternating signs.*

Therefore, we need only to prove this statement. We divide our proof into two parts.

*Part 1. The “if” part.* If  $0$  were not the best uniform approximation in  $\mathcal{P}_n$  of  $f$ , then there would exist  $p \in \mathcal{P}_n$  such that

$$0 < \|f - p\| < \|f - 0\| = \|f\|. \quad (1.17)$$

Let  $a \leq x_1 < \dots < x_{n+2} \leq b$  be such that

$$\begin{aligned} |f(x_k)| &= \|f\|, & k &= 1, \dots, n+2, \\ f(x_k)f(x_{k+1}) &< 0, & k &= 1, \dots, n+1. \end{aligned}$$

It follows from (1.17) that

$$|p(x_k) - f(x_k)| \leq \|p - f\| < \|f\| = |f(x_k)|, \quad k = 1, \dots, n+2.$$

Consequently,

$$\text{sign } p(x_k) = \text{sign } (p(x_k) - f(x_k) + f(x_k)) = \text{sign } f(x_k), \quad k = 1, \dots, n+2,$$

where  $\text{sign } z = z/|z|$  for any nonzero  $z \in \mathbb{R}$  and  $\text{sign } 0 = 0$ . Thus,  $p$  changes its sign at  $n + 2$  points. Hence,  $p$  has at least  $n + 1$  roots. But  $p \in \mathcal{P}_n$ . So,  $p = 0$ . By (1.17),  $\|0 - f\| < \|f\|$ . This is a contradiction.

*Part 2. The “only if” part.* Assume the assertion were not true. Then, there would exist  $m + 1$  points with  $0 \leq m \leq n$  such that  $f$  achieves its maximum magnitude at these points with alternating signs. If  $m = 0$ , then  $f$  would have the same sign at all its points of maximum magnitude. In this case, there would exist a nonzero constant  $c$  such that  $\|f - c\| < \|f\|$ . This contradicts the fact that 0 is the best uniform approximation of  $f$  in  $\mathcal{P}_n$ . Thus,  $1 \leq m \leq n$ . We shall construct a polynomial in  $\mathcal{P}_n$  that would be closer to  $f$  than 0 with respect to the norm  $\|\cdot\|$ .

*Step 1.* Let  $Z = \{x \in [a, b] : |f(x)| = \|f\|\}$ . Without loss of generality, we may assume that  $f > 0$  at the smallest number of  $Z$ . Such smallest number exists, since  $Z$  is closed in  $[a, b]$ , a consequence of the continuity of  $f$ . Since  $\text{sign } f$  alternates at  $m + 1$  points in  $Z$ , there exist  $\xi_k$  ( $1 \leq k \leq m$ ) with  $a \leq \xi_1 < \dots < \xi_m \leq b$  such that  $\text{sign } f$  alternates in  $Z \cap [a, \xi_1], Z \cap [\xi_1, \xi_2], \dots, Z \cap [\xi_{m-1}, \xi_m], Z \cap [\xi_m, b]$ . Define

$$p(x) = (\xi_1 - x) \cdots (\xi_m - x).$$

Clearly,  $p \in \mathcal{P}_n$ , and

$$\text{sign } p(x) = \text{sign } f(x) \quad \forall x \in Z. \quad (1.18)$$

Let  $x \in Z$ . Since  $|f(x)| = \|f\| > 0$ , (1.18) implies that  $p(x) \neq 0$ . Since  $Z$  is closed in  $[a, b]$ ,

$$\min_{x \in Z} |p(x)| > 0. \quad (1.19)$$

In particular, this implies that  $\|p\| > 0$ .

*Step 2.* We show that

$$\left\| f - \frac{\varepsilon}{\|p\|} p \right\| < \|f\| \quad (1.20)$$

for small  $\varepsilon > 0$ .

Let

$$\delta = \min_{x \in Z} \frac{p(x)f(x)}{\|p\|} = \min_{x \in Z} \frac{|p(x)||f(x)|}{\|p\|} = \frac{\|f\|}{\|p\|} \min_{x \in Z} |p(x)|.$$

By (1.19),  $\delta > 0$ . Define

$$A = \left\{ x \in [a, b] : \frac{p(x)f(x)}{\|p\|} > \frac{\delta}{2} \right\} \quad \text{and} \quad B = \left\{ x \in [a, b] : \frac{p(x)f(x)}{\|p\|} \leq \frac{\delta}{2} \right\}.$$

Clearly,  $[a, b] = A \cup B$ ,  $A$  and  $B$  are disjoint, and  $B$  is closed in  $[a, b]$ . If  $x \in A$ , then

$$\left| f(x) - \frac{\varepsilon}{\|p\|} p(x) \right|^2 = |f(x)|^2 - 2\varepsilon \frac{p(x)f(x)}{\|p\|} + \varepsilon^2 \left( \frac{p(x)}{\|p\|} \right)^2$$

$$\begin{aligned}
&\leq \|f\|^2 - 2\varepsilon \frac{p(x)f(x)}{\|p\|} + \varepsilon^2 \\
&\leq \|f\|^2 - \varepsilon\delta + \varepsilon^2 = \|f\|^2 - \frac{\delta^2}{4} \quad \text{for } 0 < \varepsilon < \frac{\delta}{2}.
\end{aligned} \tag{1.21}$$

If  $x \in B$ , then by the definition of  $\delta$  and  $B$ ,

$$\frac{p(x)f(x)}{\|p\|} \leq \frac{\delta}{2} < \delta \leq \frac{p(y)f(y)}{\|p\|} \quad \forall y \in Z.$$

This implies that  $x \notin Z$ , and hence  $|f(x)| < \|f\|$ . Consequently, since  $f$  is continuous and  $B$  is closed in  $[a, b]$ , we obtain that

$$M := \max_{x \in B} |f(x)| < \|f\|.$$

Therefore, for  $0 < \varepsilon < \|f\| - M$ ,

$$\left| f(x) - \frac{\varepsilon}{\|p\|} p(x) \right| \leq |f(x)| + \left| \frac{\varepsilon}{\|p\|} p(x) \right| \leq M + \varepsilon < \|f\| \quad \forall x \in B. \tag{1.22}$$

It therefore follows from (1.21) and (1.22) that

$$\left\| f - \frac{\varepsilon}{\|p\|} p \right\| < \|f\| \quad \text{if } 0 < \varepsilon < \min\left(\frac{\delta}{2}, \|f\| - M\right).$$

This contradicts the fact that 0 is the best uniform approximation in  $\mathcal{P}_n$  of  $f$ .  $\square$

**Theorem 1.9 (Uniqueness of best uniform approximation).** *For any  $f \in C[a, b]$  and integer  $n \geq 0$ , the best approximation of  $f$  in  $\mathcal{P}_n$  is unique.*

*Proof.* Without loss of generality, we may assume  $f \notin \mathcal{P}_n$ . Assume that  $p, q \in \mathcal{P}_n$  are best uniform approximations of  $f$  in  $\mathcal{P}_n$ , i.e.,

$$\|f - p\| = \|f - q\| = E_n(f),$$

where  $E_n(f)$  is defined in (1.13). Let  $r = (p + q)/2 \in \mathcal{P}_n$ . Then,

$$E_n(f) \leq \|f - r\| = \left\| \frac{1}{2}(f - p) + \frac{1}{2}(f - q) \right\| \leq \frac{1}{2}\|f - p\| + \frac{1}{2}\|f - q\| = E_n(f).$$

Hence,  $r \in \mathcal{P}_n$  is also a best uniform approximation of  $f$  in  $\mathcal{P}_n$ .

By the Chebyshev Alternation Theorem, there exist  $n + 2$  points  $x_k$  ( $k = 1, \dots, n + 2$ ) with  $a \leq x_1 < \dots < x_{n+2} \leq b$  such that

$$|f(x_k) - r(x_k)| = \|r - f\| = E_n(f), \quad k = 1, \dots, n + 2.$$

Fix  $k$  ( $1 \leq k \leq n+2$ ). Assume first  $f(x_k) - r(x_k) = E_n(f)$ . Then, we have

$$f(x_k) - \frac{1}{2}[p(x_k) + q(x_k)] = f(x_k) - r(x_k) = E_n(f) = \|f - p\| \geq f(x_k) - p(x_k),$$

leading to  $q(x_k) \leq p(x_k)$ . Similarly, we have  $p(x_k) \leq q(x_k)$ . Thus,  $q(x_k) = p(x_k)$ . Assume now  $f(x_k) - r(x_k) = -E_n(f)$ . We have

$$f(x_k) - \frac{1}{2}[p(x_k) + q(x_k)] = f(x_k) - r(x_k) = -E_n(f) = -\|f - p\| \leq f(x_k) - p(x_k),$$

leading to  $q(x_k) \geq p(x_k)$ . Similarly,  $p(x_k) \geq q(x_k)$ . Hence,  $q(x_k) = p(x_k)$ . Therefore,  $p(x_k) = q(x_k)$  for all the  $n+2$  distinct points  $x_1, \dots, x_{n+2}$ . Hence,  $p = q$  in  $\mathcal{P}_n$ .  $\square$

### 1.3 Chebyshev Polynomials

Consider  $n \geq 1$  and  $f(x) = x^n$  for  $x \in [-1, 1]$ . By Theorem 1.7, There exists a unique  $p_{n-1} \in \mathcal{P}_{n-1}$  such that

$$\|f - p_{n-1}\|_{C[-1,1]} \leq \|f - q_{n-1}\|_{C[-1,1]} \quad \forall q_{n-1} \in \mathcal{P}_{n-1}.$$

Denote for any integer  $k \geq 0$

$$\tilde{\mathcal{P}}_k = \{p \in \mathcal{P}_k : \text{the leading coefficient of } p \text{ is } 1\}.$$

Then  $\tilde{T}_n := f - p_{n-1} \in \tilde{\mathcal{P}}_n$  is the unique polynomial in  $\tilde{\mathcal{P}}_n$  that satisfies

$$\|\tilde{T}_n\|_{C[-1,1]} = \min_{\tilde{p} \in \tilde{\mathcal{P}}_n} \|\tilde{p}\|_{C[-1,1]}. \quad (1.23)$$

By the Chebyshev Alternation Theorem,  $\tilde{T}_n$  is characterized by achieving its maximum magnitude at  $n+1$  points in  $[-1, 1]$  with alternating signs.

To find  $\tilde{T}_n$ , we consider for each  $n \geq 0$ <sup>1</sup>

$$T_n(x) = \cos(n \arccos x) \quad \forall x \in [-1, 1]. \quad (1.24)$$

Introducing  $x = \cos \theta$  for all  $\theta \in [0, \pi]$ , we can write

$$T_n(x) = \cos n\theta = \cos(n \arccos x) \quad \forall x \in [-1, 1].$$

Since

$$\cos(n+1)\theta + \cos(n-1)\theta = 2 \cos \theta \cos n\theta, \quad (1.25)$$

---

<sup>1</sup>See [8] for more on the derivation of  $\tilde{T}_n$ .

we obtain

$$T_{n+1}(x) = 2xT_n(x) - T_{n-1}(x) \quad x \in [-1, 1]. \quad (1.26)$$

Clearly,  $T_0(x) = 1$  and  $T_1(x) = x$ . Therefore, by induction, we conclude that  $T_n$  is a polynomial of degree  $n$ , and for  $n \geq 1$  the leading coefficient of  $T_n$  is  $2^{n-1}$ .

Let  $\theta_k = k\pi/n \in [0, \pi]$  and  $x_k = \cos \theta_k$ ,  $k = 0, \dots, n$ . Then,

$$T_n(x_k) = \cos n\theta_k = \cos k\pi = (-1)^k = (-1)^k \|T_n\|_{C[-1,1]}, \quad k = 0, \dots, n.$$

This means that  $T_n \in \mathcal{P}_n$  achieves its maximum magnitude at  $n + 1$  points in  $[-1, 1]$  with alternating signs. Therefore, for  $n \geq 1$ ,

$$\tilde{T}_n(x) = 2^{1-n}T_n(x) = 2^{1-n} \cos(n \arccos x), \quad x \in [-1, 1]. \quad (1.27)$$

We call  $T_n$  ( $n \geq 0$ ) the *Chebyshev polynomials of the first kind*. The first few of these polynomials are

$$\begin{aligned} T_0(x) &= 1, \\ T_1(x) &= x, \\ T_2(x) &= 2x^2 - 1, \\ T_3(x) &= 4x^3 - 3x, \\ T_4(x) &= 8x^4 - 8x^2 + 1. \end{aligned}$$

The following theorem summarizes the properties of such polynomials:

**Theorem 1.10 (Properties of Chebyshev polynomials of the first kind).**

(1) Orthogonality.

$$\int_{-1}^1 \frac{1}{\sqrt{1-x^2}} T_m(x) T_n(x) dx = \begin{cases} 0 & \text{if } m \neq n, \\ \pi & \text{if } m = n = 0, \\ \pi/2 & \text{if } m = n > 0. \end{cases} \quad (1.28)$$

(2) Recurrence.

$$\begin{aligned} T_0(x) &= 1, \\ T_1(x) &= x, \\ T_{n+1}(x) &= 2xT_n(x) - T_{n-1}(x), \quad n = 1, \dots \end{aligned}$$

*In particular, each  $T_n$  is a polynomial of degree  $n$ . Moreover, if  $n$  is an even (odd) number, then  $T_n$  is an even (odd) function.*

(3) Extreme points and zeros.

$$\begin{aligned} \|T_n\|_{C[-1,1]} &= 1, \quad n = 0, \dots, \\ T_n \left( \cos \frac{k\pi}{n} \right) &= (-1)^k = (-1)^k, \quad k = 0, \dots, n, \quad n = 1, \dots, \\ T_n \left( \cos \frac{(2k-1)\pi}{2n} \right) &= 0, \quad k = 1, \dots, n, \quad n = 1, \dots \end{aligned}$$

(4) Best uniform approximation and least-squares approximation. For each  $n \geq 1$ ,  $\tilde{T}_n = 2^{1-n}T_n \in \tilde{\mathcal{P}}_n$  is the unique polynomial in  $\tilde{\mathcal{P}}_n$  such that

$$\begin{aligned} \|\tilde{T}_n\|_{C[-1,1]} &= \min_{\tilde{p} \in \tilde{\mathcal{P}}_n} \|\tilde{p}\|_{C[-1,1]} = 2^{1-n}, \\ \int_{-1}^1 \frac{1}{\sqrt{1-x^2}} [\tilde{T}_n(x)]^2 dx &= \min_{\tilde{p}_n \in \tilde{\mathcal{P}}_n} \int_{-1}^1 \frac{1}{\sqrt{1-x^2}} [\tilde{p}_n(x)]^2 dx = 2^{1-2n}\pi. \end{aligned}$$

(5) Differential equation.

$$(1-x^2)T_n''(x) - xT_n'(x) + n^2T_n(x) = 0, \quad n = 0, \dots$$

(6) The generating function.

$$\frac{1-tx}{1-2tx+t^2} = \sum_{n=0}^{\infty} T_n(x)t^n \quad \forall t \in (-1, 1) \quad \forall x \in [-1, 1].$$

(7) Rodrigues' formula.

$$T_n(x) = \frac{(-1)^n}{(2n-1)!!} (1-x^2)^{\frac{1}{2}} \frac{d^n}{dx^n} (1-x^2)^{n-\frac{1}{2}}, \quad n = 0, \dots$$

*Proof.* We only prove Parts (1)–(6). The proof of Part (7) can be found in [9].

(1) By the change of variable  $x = \cos \theta$ , we obtain

$$\begin{aligned} \int_{-1}^1 \frac{1}{\sqrt{1-x^2}} T_m(x)T_n(x) dx &= \int_0^\pi \cos m\theta \cos n\theta d\theta \\ &= \frac{1}{2} \int_0^\pi [\cos(m+n)\theta + \cos(m-n)\theta] d\theta. \end{aligned}$$

Since

$$\int_0^\pi \cos k\theta d\theta = 0 \quad \text{for any integer } k \geq 1,$$

simple calculations then lead to (1.28).

(2) This is proved above, cf. (1.24)–(1.26).

(3) This follows from the definition (1.24) and direct calculations.

(4) The first minimization property is proved above, cf. (1.23) and (1.27). The second minimization property follows from Theorem 1.21 in Section 1.8 on properties of orthogonal polynomials and (1.28).

(5) By the definition (1.24), we have

$$T'_n(x) = \frac{n \sin(n \arccos x)}{\sqrt{1-x^2}},$$

$$T''_n(x) = \frac{-n^2 \cos(n \arccos x)}{1-x^2} + \frac{nx \sin(n \arccos x)}{(1-x^2)^{3/2}}.$$

These, together with (1.24), imply the desired differential equation.

(6) Let  $i$  be the complex unit, i.e.,  $i^2 = -1$ . Denote  $\mathcal{R}(z)$  the real part of a complex number  $z$ . We have using  $x = \cos \theta$  that

$$\sum_{n=0}^{\infty} T_n(x)t^n = \sum_{n=0}^{\infty} \cos(n\theta)t^n = \sum_{n=0}^{\infty} \mathcal{R}(t^n e^{in\theta}) = \mathcal{R}\left(\sum_{n=0}^{\infty} (te^{i\theta})^n\right).$$

Since

$$\sum_{n=0}^{\infty} (te^{i\theta})^n = \frac{1}{1-te^{i\theta}} = \frac{1}{1-t\cos\theta-it\sin\theta} = \frac{1-t\cos\theta+it\sin\theta}{(1-t\cos\theta)^2+t^2\sin^2\theta} = \frac{1-tx+it\sin\theta}{1-2tx+t^2},$$

we have

$$\sum_{n=0}^{\infty} T_n(x)t^n = \mathcal{R}\left(\sum_{n=0}^{\infty} (te^{i\theta})^n\right) = \frac{1-tx}{1-2tx+t^2},$$

completing the proof. □

## 1.4 Uniform Approximation by Trigonometric Polynomials

## 1.5 Modulus of Continuity and Jackson's Theorems

## 1.6 Least-Squares Approximation

**Definition 1.11 (Weight functions).** A weight function on a finite interval  $[a, b]$  is a non-negative, integrable function  $\rho : (a, b) \rightarrow \mathbb{R}$  that satisfies

$$\int_c^d \rho(x)dx > 0 \quad \text{for any sub-interval } (c, d) \subseteq (a, b). \quad (1.29)$$

For a measurable function  $\rho : (a, b) \rightarrow \mathbb{R}$ , the above condition (1.29) is equivalent to the condition that  $\rho > 0$  almost everywhere in  $(a, b)$ .

**Examples.**

- (1) For any  $[a, b]$ ,  $\rho(x) \equiv 1$  defines a weight function.
- (2) For  $[a, b] = [-1, 1]$ ,  $\rho(x) = 1/\sqrt{1-x^2}$  defines a weight function.
- (3) Let  $\rho : [a, b] \rightarrow \mathbb{R}$  satisfy the following: (a)  $\rho$  is integrable on  $[a, b]$ ; (b)  $\rho$  is continuous in  $(a, b)$ ; (c)  $\rho(x) \geq 0$  for any  $x \in (a, b)$  and  $\rho$  has at most finitely many zeros in  $(a, b)$ . Then,  $\rho$  is a weight function on  $[a, b]$ .

We assume in the rest of this section that  $\rho$  is a weight function on  $[a, b]$ . We denote by  $L^2_\rho(a, b)$  the set of all measurable functions  $f : [a, b] \rightarrow \mathbb{R}$  such that

$$\int_a^b \rho(x)[f(x)]^2 dx < \infty.$$

If  $\rho(x) = 1$  for all  $x \in (a, b)$ , then we simply write  $L^2(a, b)$  instead of  $L^2_\rho(a, b)$ . Under the usual addition and scalar multiplication,  $L^2_\rho(a, b)$  is a vector space. Clearly,  $\mathcal{P} \subset C[a, b] \subseteq L^2_\rho(a, b)$ . If  $f, g \in L^2_\rho(a, b)$ , then

$$\int_a^b \rho(x)|f(x)g(x)| dx \leq \int_a^b \rho(x) \frac{[f(x)]^2 + [g(x)]^2}{2} dx < \infty. \quad (1.30)$$

Thus,  $\rho fg$  is integrable. In particular, setting  $g(x) = 1$ , we see that  $\rho f$  is integrable.

**Proposition 1.12 (The Cauchy–Schwarz inequality).** *We have*

$$\left| \int_a^b \rho(x)f(x)g(x) dx \right| \leq \sqrt{\int_a^b \rho(x)[f(x)]^2 dx} \sqrt{\int_a^b \rho(x)[g(x)]^2 dx} \quad \forall f, g \in L^2_\rho(a, b). \quad (1.31)$$

*Proof.* Consider the non-negative quadratic function  $\phi : \mathbb{R} \rightarrow \mathbb{R}$  defined by

$$\begin{aligned} \phi(\lambda) &= \int_a^b \rho(x) [f(x) + \lambda g(x)]^2 dx \\ &= \lambda^2 \int_a^b \rho(x)[g(x)]^2 dx + 2\lambda \int_a^b \rho(x)f(x)g(x) dx + \int_a^b \rho(x)[f(x)]^2 dx \quad \forall \lambda \in \mathbb{R}. \end{aligned}$$

If the leading coefficient of  $\phi$  is 0, then the inequality (1.31) holds true trivially. If it is nonzero, then the discriminant of  $\phi$  is non-positive, i.e.,

$$4 \left[ \int_a^b \rho(x)f(x)g(x) dx \right]^2 - 4 \int_a^b \rho(x)[g(x)]^2 dx \int_a^b \rho(x)[f(x)]^2 dx \leq 0.$$

This leads to the inequality in (1.31). □

A different proof of the Cauchy–Schwarz inequality (1.31) is as follows: Applying Fubini’s Theorem, we have for any  $f, g \in L^2_\rho(a, b)$  that

$$\begin{aligned}
0 &\leq \int_0^1 \int_0^1 \rho(x)\rho(y) [f(x)g(y) - f(y)g(x)]^2 dx dy \\
&= \int_0^1 \int_0^1 \rho(x)\rho(y)[f(x)]^2[g(y)]^2 dx dy + \int_0^1 \int_0^1 \rho(x)\rho(y)[f(y)]^2[g(x)]^2 dx dy \\
&\quad - 2 \int_0^1 \int_0^1 \rho(x)\rho(y)f(x)g(x)f(y)g(y) dx dy \\
&= 2 \left( \int_0^1 \rho(x)[f(x)]^2 dx \right) \left( \int_0^1 \rho(x)[g(x)]^2 dx \right) - 2 \left( \int_0^1 f(x)g(x) dx \right)^2,
\end{aligned}$$

leading to (1.31).

**Definition 1.13 (Least-squares approximation).** A least-squares approximation of a given function  $f \in L^2_\rho(a, b)$  in  $\mathcal{P}_n$  is a polynomial  $p_n \in \mathcal{P}_n$  that satisfies

$$\int_a^b \rho(x)[f(x) - p_n(x)]^2 dx \leq \int_a^b \rho(x)[f(x) - q_n(x)]^2 dx \quad \forall q_n \in \mathcal{P}_n. \quad (1.32)$$

**Theorem 1.14 (Existence, uniqueness, and characterization of least-squares approximations).** Let  $f \in L^2_\rho(a, b)$ . Let  $n \geq 0$  be an integer.

- (1) There exists a unique least-squares approximation of  $f$  in  $\mathcal{P}_n$ .
- (2) Let  $p_n \in \mathcal{P}_n$ . Then  $p_n$  is the least-squares approximation of  $f$  in  $\mathcal{P}_n$  if and only if

$$\int_a^b \rho(x)[f(x) - p_n(x)]q_n(x) dx = 0 \quad \forall q_n \in \mathcal{P}_n. \quad (1.33)$$

- (3) If  $p_n \in \mathcal{P}_n$  is the least-squares approximation of  $f$  in  $\mathcal{P}_n$ , then

$$\int_a^b \rho(x)[f(x) - p_n(x)]^2 dx = \int_a^b \rho(x)[f(x)]^2 dx - \int_a^b \rho(x)[p_n(x)]^2 dx. \quad (1.34)$$

To prove this theorem, we introduce the  $(n + 1) \times (n + 1)$  matrix

$$G_{n+1} := \left[ \int_a^b \rho(x)x^{i+j} dx \right]_{i,j=0}^n. \quad (1.35)$$

In the special case that  $[a, b] = [0, 1]$  and  $\rho(x) \equiv 1$ , this is a Hilbert matrix.

**Lemma 1.15.** The matrix  $G_{n+1}$  is symmetric positive definite.

*Proof.* Define

$$Q(\xi) = \frac{1}{2} \int_a^b \rho(x) \left( \sum_{i=0}^n \xi_i x^i \right)^2 dx \quad \forall \xi = (\xi_0, \dots, \xi_n) \in \mathbb{R}^{n+1}.$$

Clearly,  $Q : \mathbb{R}^{n+1} \rightarrow \mathbb{R}$  is a non-negative quadratic form. If  $Q(\xi) = 0$  for some  $\xi \in \mathbb{R}^{n+1}$ , then, by Lemma 1.16,  $\sum_{i=0}^n \xi_i x^i = 0$  for all  $x \in [a, b]$ . Hence,  $\xi = 0$  in  $\mathbb{R}^{n+1}$ . Therefore,  $Q : \mathbb{R}^{n+1} \rightarrow \mathbb{R}$  is a positive quadratic form.

It is clear that

$$Q(\xi) = \frac{1}{2} \sum_{i,j=0}^n \left( \int_a^b \rho(x) x^{i+j} dx \right) \xi_i \xi_j \quad \forall \xi = (\xi_0, \dots, \xi_n) \in \mathbb{R}^{n+1}.$$

Therefore,  $G_{n+1}$  is the matrix associated with the positive quadratic form  $Q : \mathbb{R}^{n+1} \rightarrow \mathbb{R}$ . Hence,  $G_{n+1}$  is symmetric positive definite.  $\square$

*Proof of Theorem 1.14.* (1) Define

$$F(c) = \int_a^b \rho(x) \left[ f(x) - \sum_{i=0}^n c_i x^i \right]^2 dx \quad \forall c = (c_0, \dots, c_n) \in \mathbb{R}^{n+1}. \quad (1.36)$$

Clearly,  $F : \mathbb{R}^{n+1} \rightarrow \mathbb{R}$  is a non-negative quadratic function. Direct calculations lead to

$$F(c) = \sum_{i,j=0}^n \left[ \int_a^b \rho(x) x^{i+j} dx \right] c_i c_j - 2 \sum_{i=0}^n \left[ \int_a^b \rho(x) f(x) x^i dx \right] c_i - \int_a^b \rho(x) [f(x)]^2 dx.$$

Thus, the Hessian matrix of  $F$  is

$$\left[ \frac{\partial^2 F}{\partial c_i \partial c_j} \right]_{i,j=0}^n = 2 \left[ \int_a^b \rho(x) x^{i+j} dx \right]_{i,j=0}^n = 2G_{n+1},$$

where  $G_{n+1}$  is the matrix defined in (1.35). By Lemma 1.15, the Hessian matrix of  $F$  is therefore symmetric positive definite. Consequently,  $F : \mathbb{R}^{n+1} \rightarrow \mathbb{R}$  is a convex quadratic form. Since it is non-negative, it admits a unique minimizer  $\hat{c} = (\hat{c}_0, \dots, \hat{c}_n) \in \mathbb{R}^{n+1}$ . Define

$$p_n(x) = \sum_{i=0}^n \hat{c}_i x^i \quad \forall x \in \mathbb{R}. \quad (1.37)$$

Then,  $p_n \in \mathcal{P}_n$  is the unique polynomial in  $\mathcal{P}_n$  that satisfies (1.32).

(2) From Part (1), the unique least-squares approximation  $p_n \in \mathcal{P}_n$  is given by (1.37), where  $\hat{c} = (\hat{c}_0, \dots, \hat{c}_n) \in \mathbb{R}^{n+1}$  is the unique minimizer of the convex quadratic function  $F : \mathbb{R}^{n+1} \rightarrow \mathbb{R}$ . Clearly,  $\hat{c}$  is the unique critical point of  $F$  determined by

$$\partial_j F(\hat{c}) = 0, \quad j = 0, \dots, n. \quad (1.38)$$

By the definition of  $F : \mathbb{R}^{n+1} \rightarrow \mathbb{R}$  (cf. (1.36)) and the Chain Rule, we obtain for any  $c = (c_0, \dots, c_n) \in \mathbb{R}^{n+1}$  that

$$\partial_j F(c) = 2 \int_a^b \rho(x) \left[ f(x) - \sum_{i=0}^n c_i x^i \right] (-x^j) dx, \quad j = 0, \dots, n. \quad (1.39)$$

Therefore, by the definition of  $p_n$  in (1.37), the system of equations (1.38) is equivalent to

$$\int_a^b \rho(x) [f(x) - p_n(x)] x^j dx, \quad j = 0, \dots, n,$$

which in turn is equivalent to (1.33), since  $\mathcal{P}_n = \text{Span}\{1, x, \dots, x^n\}$ .

(3) Suppose  $p_n \in \mathcal{P}_n$  is the least-squares approximation of  $f$ . By (1.33) with  $q_n = p_n$ , we have

$$\int_a^b \rho(x) [f(x) - p_n(x)] p_n(x) dx = 0.$$

Therefore,

$$\begin{aligned} \int_a^b \rho(x) [f(x)]^2 dx &= \int_a^b \rho(x) [f(x) - p_n(x) + p_n(x)]^2 dx \\ &= \int_a^b \rho(x) [f(x) - p_n(x)]^2 dx + \int_a^b \rho(x) [p_n(x)]^2 dx \\ &\quad + 2 \int_a^b \rho(x) [f(x) - p_n(x)] p_n(x) dx \\ &= \int_a^b \rho(x) [f(x) - p_n(x)]^2 dx + \int_a^b \rho(x) [p_n(x)]^2 dx. \end{aligned}$$

This implies (1.34). □

From the proof, cf. (1.38) and (1.39), we see that this least-squares approximation  $p_n(x) = \sum_{i=0}^n \hat{c}_i x^i$  can be obtained by solving the system of linear equations

$$\sum_{i=0}^n \left[ \int_a^b \rho(x) x^{i+j} dx \right] \hat{c}_i = \int_a^b \rho(x) f(x) x^j dx \quad j = 0, \dots, n.$$

The coefficient matrix of this system of linear equations is exactly  $G_{n+1}$ , defined in (1.35).

## 1.7 Orthogonal Polynomials

Fix a weight function  $\rho$  on  $[a, b]$ . For convenience, we denote

$$\langle f, g \rangle = \int_a^b \rho(x) f(x) g(x) dx \quad \forall f, g \in L_\rho^2(a, b).$$

By (1.30),  $\langle f, g \rangle$  is finite for any  $f, g \in L^2_\rho(a, b)$ . The mapping  $\langle \cdot, \cdot \rangle : L^2_\rho(a, b) \times L^2_\rho(a, b) \rightarrow \mathbb{R}$  clearly satisfies the following properties:

(1) Symmetry.

$$\langle f, g \rangle = \langle g, f \rangle \quad \forall f, g \in L^2_\rho(a, b);$$

(2) Bilinearity.

$$\langle \alpha f + \beta g, h \rangle = \alpha \langle f, h \rangle + \beta \langle g, h \rangle \quad \forall f, g, h \in L^2_\rho(a, b) \quad \forall \alpha, \beta \in \mathbb{R};$$

(3) Non-negativity.

$$\langle f, f \rangle \geq 0 \quad \forall f \in L^2_\rho(a, b).$$

**Lemma 1.16.** *Let  $f \in C[a, b]$ . If*

$$\int_a^b \rho(x)[f(x)]^2 dx = 0,$$

*then  $f(x) = 0$  for all  $x \in [a, b]$ .*

*Proof.* Suppose there existed a point  $x_0 \in [a, b]$  such that  $f(x_0) \neq 0$ . By the continuity of  $f$ , there would have existed a sub-interval  $[c, d] \subseteq [a, b]$  with  $c < d$  such that  $|f(x)| \geq \varepsilon_0$  for all  $x \in [c, d]$  for some constant  $\varepsilon_0 > 0$ . Now, by the definition of a weight function,  $\int_c^d \rho(x) dx > 0$ . Therefore,

$$\int_a^b \rho(x)[f(x)]^2 dx \geq \int_c^d \rho(x)[f(x)]^2 dx \geq \varepsilon_0^2 \int_c^d \rho(x) dx > 0.$$

This contradicts the assumption. □

Consider the subspace  $C[a, b]$  of  $L^2_\rho(a, b)$ . The mapping  $\langle \cdot, \cdot \rangle : C[a, b] \times C[a, b] \rightarrow \mathbb{R}$  is in fact an inner product of  $C[a, b]$ , i.e., it is symmetric, bilinear, and positive. Here, the positivity means

$$\begin{aligned} \langle f, f \rangle &\geq 0 && \forall f \in C[a, b]. \\ \langle f, f \rangle &= 0 && \text{if and only if } f = 0 \text{ in } C[a, b]. \end{aligned}$$

The “only if” part in the last equation follows from Lemma 1.16. The norm of  $C[a, b]$  induced by this inner product is

$$\|f\|_{L^2_\rho(a, b)} = \sqrt{\langle f, f \rangle} = \sqrt{\int_a^b \rho(x)[f(x)]^2 dx} \quad (1.40)$$

for any  $f \in C[a, b]$ . This is called the  $L^2_\rho(a, b)$ -norm. We will also use (1.40) for any  $f \in L^2_\rho(a, b)$ . It is often written as  $\|\cdot\|$  instead of  $\|\cdot\|_{L^2_\rho(a, b)}$ , when no confusion arises.

**Definition 1.17 (Orthogonal polynomials).** A sequence of polynomials  $Q_n$  ( $n = 0, \dots$ ) are called orthogonal polynomials in  $L^2_\rho(a, b)$ , if the following hold true:

- (1) For each integer  $n \geq 0$ ,  $Q_n$  is a polynomial of degree exactly  $n$ ;
- (2)  $\langle Q_m, Q_n \rangle = 0$  whenever  $m \neq n$ .

A sequence of orthogonal polynomials  $Q_n$  ( $n = 0, \dots$ ) in  $L^2_\rho(a, b)$  are called orthonormal polynomials in  $L^2_\rho(a, b)$ , if they satisfy

- (3)  $\langle Q_n, Q_n \rangle = 1$  for all  $n \geq 0$ .

A set of polynomials are called orthogonal (or orthonormal) in  $L^2_\rho(a, b)$  if it is a subset of a sequence of orthogonal (or orthonormal) polynomials.

If  $Q_n$  ( $n = 0, \dots$ ) are orthogonal polynomials, then  $\langle Q_n, Q_n \rangle \neq 0$  for each  $n \geq 0$ . Moreover, the polynomials  $Q_n/\sqrt{\langle Q_n, Q_n \rangle}$  ( $n = 0, \dots$ ) are orthonormal. In general, orthonormal polynomials  $Q_n$  ( $n = 0, \dots$ ) are characterized by

$$\langle Q_i, Q_j \rangle = \delta_{i,j} \quad \forall i, j \geq 0, \quad (1.41)$$

where  $\delta_{ij}$  is defined to be 1 if  $i = j$  and 0 if  $i \neq j$ .

The Chebyshev polynomials  $T_n$  ( $n = 0, \dots$ ) are orthogonal polynomials in  $L^2_\rho(-1, 1)$  with  $\rho(x) = 1/\sqrt{1-x^2}$ . Another important class of orthogonal polynomials are Legendre polynomials  $P_n$  ( $n = 0, \dots$ ) in  $L^2(-1, 1)$  that will be discussed in Section 1.9.

**Lemma 1.18.** If  $Q_0, \dots, Q_n$  be orthogonal polynomials in  $L^2_\rho(a, b)$ , then

- (1) The  $n + 1$  polynomials  $Q_0, \dots, Q_n$  are linearly independent in  $L^2_\rho(a, b)$ ;
- (2)  $\mathcal{P}_n = \text{Span}\{Q_0, \dots, Q_n\}$ . Moreover,

$$p_n = \sum_{k=0}^n \frac{\langle p_n, Q_k \rangle}{\langle Q_k, Q_k \rangle} Q_k \quad \forall p_n \in \mathcal{P}_n. \quad (1.42)$$

*Proof.* (1) Suppose there exist  $\alpha_0, \dots, \alpha_n \in \mathbb{R}$  such that  $\sum_{k=0}^n \alpha_k Q_k = 0$ . Multiplying both sides of this equation by  $\rho Q_j$  for an arbitrary but fixed  $j$  with  $0 \leq j \leq n$  and then integrate over  $[a, b]$ , we obtain by (1.41) that  $\alpha_j \langle Q_j, Q_j \rangle = 0$ , implying  $\alpha_j = 0$ , since  $\langle Q_j, Q_j \rangle \neq 0$  by Lemma 1.16. Hence,  $Q_0, \dots, Q_n$  are linearly independent.

(2) Since the dimension of  $\mathcal{P}_n$  is  $n + 1$ , the  $n + 1$  linearly independent polynomials  $Q_0, \dots, Q_n$  in  $\mathcal{P}_n$  form a basis of  $\mathcal{P}_n$ . Hence, they span  $\mathcal{P}_n$ .

Fix  $p_n \in \mathcal{P}_n = \text{Span}\{Q_0, \dots, Q_n\}$ . There exist constants  $c_i \in \mathbb{R}$  ( $i = 0, \dots, n$ ) such that  $p_n = \sum_{k=0}^n c_k Q_k$ . Multiplying both sides of this equation by  $\rho Q_j$  for an arbitrary but fixed  $j$  with  $0 \leq j \leq n$  and then integrate over  $[a, b]$ , we obtain by (1.41) that  $\langle p_n, Q_j \rangle = c_j \langle Q_j, Q_j \rangle$ . This implies (1.42).  $\square$

**Theorem 1.19 (Characterization of least-squares approximations using orthonormal polynomials).** Let  $Q_0, \dots, Q_n$  be orthonormal polynomials in  $L^2_\rho(a, b)$ . Then the least-squares approximation of a given  $f \in L^2_\rho(a, b)$  in  $\mathcal{P}_n$  is given by

$$p_n = \sum_{k=0}^n \langle f, Q_k \rangle Q_k. \quad (1.43)$$

Moreover,

$$\|f - q_n\|^2 = \|f\|^2 - \sum_{k=0}^n \langle f, Q_k \rangle^2. \quad (1.44)$$

*Proof.* Clearly, the polynomial  $p_n$  defined in (1.43) is in  $\mathcal{P}_n$ . Fix an arbitrary integer  $j$  with  $0 \leq j \leq n$ . Since  $Q_0, \dots, Q_n$  are orthonormal, we have

$$\langle f - p_n, Q_j \rangle = \langle f, Q_j \rangle - \sum_{k=0}^n \langle f, Q_k \rangle \langle Q_k, Q_j \rangle = \langle f, Q_j \rangle - \langle f, Q_j \rangle = 0.$$

Hence, by Lemma 1.18,

$$\langle f - p_n, q_n \rangle = 0 \quad \forall q_n \in \mathcal{P}_n.$$

This is exactly (1.33). Therefore, by Theorem 1.14,  $p_n \in \mathcal{P}_n$  is the unique least-squares approximation of  $f$  in  $\mathcal{P}_n$ .

It follows from (1.43), the symmetry and bilinearity of  $\langle \cdot, \cdot \rangle$ , and the fact that  $Q_0, \dots, Q_n$  are orthonormal that

$$\langle p_n, p_n \rangle = \left\langle \sum_{i=0}^n \langle f, Q_i \rangle Q_i, \sum_{i=0}^n \langle f, Q_i \rangle Q_i \right\rangle = \sum_{i,j=0}^n \langle f, Q_i \rangle \langle f, Q_j \rangle \langle Q_i, Q_j \rangle = \sum_{i=0}^n \langle f, Q_i \rangle^2.$$

This, together with Part (3) of Theorem 1.14, implies (1.44).  $\square$

**Theorem 1.20 (The Gram–Schmidt orthogonalization).** Consider  $L^2_\rho(a, b)$ . Let  $f_0(x) = 1$  and  $f_k(x) = x^k$  for any integer  $k \geq 1$ . Define

$$g_0(x) = \frac{1}{\sqrt{\langle f_0, f_0 \rangle}} f_0(x), \quad (1.45)$$

$$\begin{cases} \hat{g}_k(x) = f_k(x) - \sum_{j=0}^{k-1} \langle f_k, g_j \rangle g_j(x), \\ g_k(x) = \frac{1}{\sqrt{\langle \hat{g}_k, \hat{g}_k \rangle}} \hat{g}_k(x), \end{cases} \quad k = 1, 2, \dots \quad (1.46)$$

Then  $g_k$  ( $k = 0, 1, \dots$ ) are orthonormal polynomials.

*Proof.* Clearly,  $g_0$  and  $g_1$  are polynomial of degrees 0 and 1, respectively. Moreover, direct calculations using (1.45) and (1.46) lead to  $\langle g_0, g_0 \rangle = \langle g_1, g_1 \rangle = 1$  and  $\langle g_0, g_1 \rangle = 0$ . Let  $k \geq 1$  be an integer. Assume that  $g_i$  is a polynomial of degree  $i$  for each  $i$  with  $0 \leq i \leq k$  and that

$$\langle g_i, g_j \rangle = \delta_{ij}, \quad 0 \leq i, j \leq k. \quad (1.47)$$

Since  $f_{k+1}$  is a polynomial of degree  $k + 1$ , we see from (1.46) with  $k$  replaced by  $k + 1$  that  $g_{k+1}$  is a polynomial of degree  $k + 1$ . Clearly,  $\langle g_{k+1}, g_{k+1} \rangle = 1$ . Moreover, for each  $i$  with  $0 \leq i \leq k$ , we have by (1.46) and (1.47) that

$$\langle \hat{g}_{k+1}, g_i \rangle = \langle f_{k+1}, g_i \rangle - \sum_{j=0}^k \langle f_{k+1}, g_j \rangle \langle g_j, g_i \rangle = \langle f_{k+1}, g_i \rangle - \langle f_{k+1}, g_i \rangle = 0.$$

Since  $g_{k+1}$  and  $\hat{g}_{k+1}$  differ by a nonzero constant multiplier, we have

$$\langle g_{k+1}, g_i \rangle = 0, \quad i = 0, \dots, k.$$

Therefore, (1.47) is true with  $k$  replaced by  $k + 1$ . Consequently,  $g_k$  ( $k = 0, 1, \dots$ ) are orthonormal polynomials.  $\square$

**Example.** Consider  $L^2[-1, 1]$ .

$$\begin{aligned} g_0(x) &= \frac{1}{\sqrt{\langle f_0, f_0 \rangle}} f_0(x) = \frac{1}{\sqrt{\int_{-1}^1 1 \cdot 1 dx}} = \frac{1}{\sqrt{2}}. \\ \hat{g}_1(x) &= f_1(x) - \langle f_1, g_0 \rangle g_0(x) = x - \left( \int_{-1}^1 \frac{x}{\sqrt{2}} dx \right) \frac{1}{\sqrt{2}} = x, \\ g_1(x) &= \frac{1}{\sqrt{\langle \hat{g}_1, \hat{g}_1 \rangle}} \hat{g}_1(x) = \frac{x}{\sqrt{\int_{-1}^1 x^2 dx}} = \sqrt{\frac{3}{2}} x. \\ \hat{g}_2(x) &= f_2(x) - \langle f_2, g_0 \rangle g_0(x) - \langle f_2, g_1 \rangle g_1(x) \\ &= x^2 - \left( \int_{-1}^1 \frac{x^2}{\sqrt{2}} dx \right) \frac{1}{\sqrt{2}} - \left( \int_{-1}^1 x^2 \sqrt{\frac{3}{2}} x dx \right) \sqrt{\frac{3}{2}} x = x^2 - \frac{1}{3}, \\ g_2(x) &= \frac{1}{\sqrt{\langle \hat{g}_2, \hat{g}_2 \rangle}} \hat{g}_2(x) = \frac{1}{\sqrt{\int_{-1}^1 (x^2 - 1/3)^2 dx}} \left( x^2 - \frac{1}{3} \right) = \frac{\sqrt{15}}{4} \left( x^2 - \frac{1}{3} \right). \end{aligned}$$

Therefore,  $g_0, g_1, g_2$  are orthonormal polynomials in  $L^2[-1, 1]$ .

## 1.8 More Properties of Orthogonal Polynomials

**Theorem 1.21 (Minimization).** *Let  $Q_n$  ( $n = 0, \dots$ ) be orthogonal polynomials in  $L^2_\rho(a, b)$ . Suppose  $n \geq 1$  and  $Q_n \in \tilde{\mathcal{P}}_n$ , then  $Q_n$  is the unique polynomial in  $\tilde{\mathcal{P}}_n$  that satisfies*

$$\|Q_n\| = \min_{q_n \in \tilde{\mathcal{P}}_n} \|q_n\|.$$

*Proof.* Since  $Q_n \in \tilde{\mathcal{P}}_n$ , we have  $Q_n(x) = x^n - q_{n-1}(x)$  for some  $q_{n-1} \in \mathcal{P}_{n-1}$ . Thus, by the orthogonality,

$$0 = \langle Q_n, q \rangle = \langle x^n - q_{n-1}, q \rangle \quad \forall q \in \mathcal{P}_{n-1}.$$

Theorem 1.14 then implies that  $q_{n-1}$  is the unique least-squares approximation of  $x^n$  in  $\mathcal{P}_{n-1}$ . This is equivalent to the assertion of the theorem.  $\square$

**Colloary 1.22 (Uniqueness of orthogonal polynomials).** *If  $\{P_n\}_{n=0}^\infty$  and  $\{Q_n\}_{n=0}^\infty$  are two systems of orthogonal polynomials in  $L_\rho^2(a, b)$ , then, for each  $n \geq 0$ , there exists  $c_n \in \mathbb{R}$  with  $c_n \neq 0$  such that  $P_n = c_n Q_n$ .*

*Proof.* Let  $\alpha_n$  and  $\beta_n$  be the leading coefficients of  $P_n$  and  $Q_n$ , respectively. Then, by Theorem 1.21,  $(1/\alpha_n)P_n = (1/\beta_n)Q_n$  for each  $n \geq 1$ . This is clearly true also for  $n = 0$ . Thus,  $P_n = c_n Q_n$  with  $c_n = \alpha_n/\beta_n$  for each  $n \geq 0$ .  $\square$

**Theorem 1.23 (The three-term recurrence).** *Consider  $L_\rho^2(a, b)$ .*

(1) *Define*

$$\begin{aligned} \tilde{Q}_0(x) &= 1, \\ \tilde{Q}_1(x) &= x - a_1, \\ \tilde{Q}_n(x) &= (x - a_n)\tilde{Q}_{n-1}(x) - b_n\tilde{Q}_{n-2}(x), \quad n = 2, 3, \dots, \\ a_n &= \frac{\langle x\tilde{Q}_{n-1}, \tilde{Q}_{n-1} \rangle}{\langle \tilde{Q}_{n-1}, \tilde{Q}_{n-1} \rangle}, \quad n = 1, 2, \dots, \\ b_n &= \frac{\langle \tilde{Q}_{n-1}, \tilde{Q}_{n-1} \rangle}{\langle \tilde{Q}_{n-2}, \tilde{Q}_{n-2} \rangle}, \quad n = 2, 3, \dots \end{aligned}$$

*Then  $\tilde{Q}_n$  ( $n = 0, 1, \dots$ ) are the orthogonal polynomials with each  $\tilde{Q}_n \in \tilde{\mathcal{P}}_n$ .*

(2) *Let  $Q_n$  ( $n = 0, 1, \dots$ ) be orthogonal polynomials. Let  $\alpha_n$  be the leading coefficient of  $Q_n$  ( $n \geq 0$ ). Then*

$$Q_n(x) = (A_n x + B_n)Q_{n-1}(x) - C_n Q_{n-2}(x) \quad n = 2, 3, \dots \quad (1.48)$$

*where*

$$A_n = \frac{\alpha_n}{\alpha_{n-1}}, \quad B_n = -\frac{\alpha_n}{\alpha_{n-1}} \frac{\langle xQ_{n-1}, Q_{n-1} \rangle}{\langle Q_{n-1}, Q_{n-1} \rangle}, \quad C_n = \frac{\alpha_n \alpha_{n-2}}{\alpha_{n-1}^2} \frac{\langle Q_{n-1}, Q_{n-1} \rangle}{\langle Q_{n-2}, Q_{n-2} \rangle}.$$

*Proof.* (1) Clearly,  $\tilde{Q}_0 \in \tilde{\mathcal{P}}_0$  and  $\tilde{Q}_1 \in \tilde{\mathcal{P}}_1$ . Since

$$a_1 = \frac{\langle x\tilde{Q}_0, \tilde{Q}_0 \rangle}{\langle \tilde{Q}_0, \tilde{Q}_0 \rangle} = \frac{\langle x, 1 \rangle}{\langle 1, 1 \rangle},$$

we have

$$\langle \tilde{Q}_0, \tilde{Q}_1 \rangle = \langle 1, x - a_1 \rangle = \langle 1, x \rangle - \langle 1, 1 \rangle a_1 = 0.$$

Let  $n \geq 2$ . Assume  $\tilde{Q}_k \in \tilde{\mathcal{P}}_k$  for all  $k \in \{0, \dots, n-1\}$  and

$$\langle \tilde{Q}_j, \tilde{Q}_k \rangle = 0, \quad \text{if } 0 \leq j, k \leq n-1 \text{ and } j \neq k. \quad (1.49)$$

Clearly,  $\tilde{Q}_n \in \tilde{\mathcal{P}}_n$ . We have by the definition of  $\tilde{Q}_n$  and (1.49) that

$$\begin{aligned} \langle \tilde{Q}_n, \tilde{Q}_{n-1} \rangle &= \langle (x - a_n)\tilde{Q}_{n-1} - b_n\tilde{Q}_{n-2}, \tilde{Q}_{n-1} \rangle \\ &= \langle x\tilde{Q}_{n-1}, \tilde{Q}_{n-1} \rangle - a_n\langle \tilde{Q}_{n-1}, \tilde{Q}_{n-1} \rangle - b_n\langle \tilde{Q}_{n-2}, \tilde{Q}_{n-1} \rangle \\ &= \langle x\tilde{Q}_{n-1}, \tilde{Q}_{n-1} \rangle - \frac{\langle x\tilde{Q}_{n-1}, \tilde{Q}_{n-1} \rangle}{\langle \tilde{Q}_{n-1}, \tilde{Q}_{n-1} \rangle} \langle \tilde{Q}_{n-1}, \tilde{Q}_{n-1} \rangle \\ &= 0, \\ \langle \tilde{Q}_n, \tilde{Q}_{n-2} \rangle &= \langle (x - a_n)\tilde{Q}_{n-1} - b_n\tilde{Q}_{n-2}, \tilde{Q}_{n-2} \rangle \\ &= \langle x\tilde{Q}_{n-1}, \tilde{Q}_{n-2} \rangle - a_n\langle \tilde{Q}_{n-1}, \tilde{Q}_{n-2} \rangle - b_n\langle \tilde{Q}_{n-2}, \tilde{Q}_{n-2} \rangle \\ &= \langle x\tilde{Q}_{n-1}, \tilde{Q}_{n-2} \rangle - \frac{\langle \tilde{Q}_{n-1}, \tilde{Q}_{n-1} \rangle}{\langle \tilde{Q}_{n-2}, \tilde{Q}_{n-2} \rangle} \langle \tilde{Q}_{n-2}, \tilde{Q}_{n-2} \rangle \\ &= \langle \tilde{Q}_{n-1}, x\tilde{Q}_{n-2} \rangle - \langle \tilde{Q}_{n-1}, \tilde{Q}_{n-1} \rangle \\ &= \langle \tilde{Q}_{n-1}, x\tilde{Q}_{n-2} - \tilde{Q}_{n-1} \rangle \\ &= 0, \end{aligned}$$

where the last equation follows from that fact that  $x\tilde{Q}_{n-2} - \tilde{Q}_{n-1} \in \tilde{\mathcal{P}}_{n-2}$ . Let  $0 \leq k \leq n-3$  with  $n \geq 3$ . We have again by the definition of  $\tilde{Q}_n$  and (1.49) that

$$\begin{aligned} \langle \tilde{Q}_n, \tilde{Q}_k \rangle &= \langle (x - a_n)\tilde{Q}_{n-1} - b_n\tilde{Q}_{n-2}, \tilde{Q}_k \rangle \\ &= \langle x\tilde{Q}_{n-1}, \tilde{Q}_k \rangle - a_n\langle \tilde{Q}_{n-1}, \tilde{Q}_k \rangle - b_n\langle \tilde{Q}_{n-2}, \tilde{Q}_k \rangle \\ &= \langle \tilde{Q}_{n-1}, x\tilde{Q}_k \rangle \\ &= 0, \end{aligned}$$

since  $x\tilde{Q}_k \in \tilde{\mathcal{P}}_{k+1} \subseteq \tilde{\mathcal{P}}_{n-2}$ . Therefore, (1.49) holds true with  $n-1$  replaced by  $n$ . Part (1) is thus proved.

(2) Note that  $\tilde{Q}_n := (1/\alpha_n)Q_n \in \tilde{\mathcal{P}}_n$ . Thus, it follows from Part (1) and the definition of all  $a_n, b_n$ , and  $A_n, B_n, C_n$  that

$$\begin{aligned} Q_n(x) &= \alpha_n\tilde{Q}_n(x) \\ &= \alpha_n[(x - a_n)\tilde{Q}_{n-1}(x) - b_n\tilde{Q}_{n-2}(x)] \end{aligned}$$

$$\begin{aligned}
&= \alpha_n \left[ (x - a_n) \frac{1}{\alpha_{n-1}} Q_{n-1}(x) - b_n \frac{1}{\alpha_{n-2}} Q_{n-2}(x) \right] \\
&= \left( \frac{\alpha_n}{\alpha_{n-1}} x - \frac{\alpha_n a_n}{\alpha_{n-1}} \right) Q_{n-1}(x) - \frac{\alpha_n b_n}{\alpha_{n-2}} Q_{n-2}(x) \\
&= (A_n x + B_n) Q_{n-1}(x) - C_n Q_{n-2}(x) \quad \forall n \geq 2.
\end{aligned}$$

This is (1.48). □

**Example.** Consider  $L^2[-1, 1]$ .

$$\begin{aligned}
\tilde{Q}_0(x) &= 1. \\
a_1 &= \frac{\langle x \tilde{Q}_0, \tilde{Q}_0 \rangle}{\langle \tilde{Q}_0, \tilde{Q}_0 \rangle} = 0, \\
\tilde{Q}_1(x) &= x - a_1 = x. \\
a_2 &= \frac{\langle x \tilde{Q}_1, \tilde{Q}_1 \rangle}{\langle \tilde{Q}_1, \tilde{Q}_1 \rangle} = \frac{\int_{-1}^1 x^3 dx}{\int_{-1}^1 x^2 dx} = 0, \\
b_2 &= \frac{\langle \tilde{Q}_1, \tilde{Q}_1 \rangle}{\langle \tilde{Q}_0, \tilde{Q}_0 \rangle} = \frac{\int_{-1}^1 x^2 dx}{\int_{-1}^1 1 dx} = \frac{1}{3}, \\
\tilde{Q}_2(x) &= (x - a_2) \tilde{Q}_1(x) - b_2 \tilde{Q}_0(x) = x^2 - \frac{1}{3}.
\end{aligned}$$

Thus,  $\tilde{Q}_0, \tilde{Q}_1, \tilde{Q}_2$  are the orthogonal polynomials in  $L^2[-1, 1]$  that have leading coefficient 1. Each of them differs by a constant multiplier from the corresponding one obtained in the previous example using the Gram–Schmidt orthogonalization.

**Theorem 1.24 (Zeros of orthogonal polynomials).** *Let  $Q_n$  ( $n = 0, \dots$ ) be orthogonal polynomials in  $L^2_\rho(a, b)$ . Then, for each  $n \geq 1$ ,  $Q_n$  has exactly  $n$  simple roots in  $(a, b)$ .*

*Proof.* Fix an integer  $n \geq 1$ . By the orthogonality,

$$\int_a^b \rho(x) Q_n(x) dx = \langle Q_n, 1 \rangle = 0.$$

Hence,  $Q_n$  changes its sign in  $(a, b)$  at least once. If  $n = 1$ , this implies that  $Q_1$  has exactly one root. Consider  $n \geq 2$ . Suppose  $Q_n$  changes its sign in  $(a, b)$  only  $k$  times with  $1 \leq k \leq n - 1$  at  $x_1, \dots, x_k$  with  $a < x_1 < \dots < x_k < b$ . Define

$$p(x) = (x - x_1) \cdots (x - x_k).$$

Clearly,  $p \in \mathcal{P}_k \subseteq \mathcal{P}_{n-1}$ . Moreover, both  $Q_n$  and  $p$  change their signs only at  $x_1, \dots, x_k$ . Thus,

$$\langle Q_n, p \rangle = \int_a^b \rho(x) Q_n(x) p(x) dx \neq 0.$$

This contradicts the fact that  $\langle Q_n, q \rangle = 0$  for any  $q \in \mathcal{P}_{n-1}$ . Hence,  $k \geq n$ . But  $Q_n \in \mathcal{P}_n$  can have at most  $n$  roots. Thus,  $k = n$ , and  $Q_n$  has exactly  $n$  simple roots in  $(a, b)$ .  $\square$

Let  $Q_n$  ( $n = 0, \dots$ ) be orthonormal polynomials in  $L^2_\rho(a, b)$  and define for each  $n \geq 0$

$$K_n(x, t) = \sum_{k=0}^n Q_k(x)Q_k(t). \quad (1.50)$$

We call  $K_n : \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}$  the *Dirichlet kernel* associated with first  $n + 1$  orthonormal polynomials  $Q_0, \dots, Q_n$  in  $L^2_\rho(a, b)$ . We have for any  $f \in L^2_\rho(a, b)$  that

$$\begin{aligned} \sum_{k=0}^n \langle f, Q_k \rangle Q_k(x) &= \sum_{k=0}^n \int_a^b \rho(t) f(t) Q_k(t) dt Q_k(x) \\ &= \sum_{k=0}^n \int_a^b \rho(t) f(t) Q_k(t) Q_k(x) dt \\ &= \int_a^b \rho(t) K_n(x, t) f(t) dt \\ &= \langle K_n(x, \cdot), f(\cdot) \rangle. \end{aligned}$$

It then follows from (1.43) in Theorem 1.19 that the least-squares approximation of  $f \in L^2_\rho(a, b)$  in  $\mathcal{P}_n$  is given by

$$p_n(x) = \langle K_n(x, \cdot), f(\cdot) \rangle.$$

**Theorem 1.25 (The Christoffel–Darboux identity).** *Let  $Q_n$  ( $n = 0, 1, \dots$ ) be orthonormal polynomials in  $L^2_\rho[a, b]$  and  $K_n : \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}$  the Dirichlet kernel associated with  $Q_0, \dots, Q_n$  for each  $n \geq 0$ . Then,*

$$K_n(x, t) = \frac{\alpha_n}{\alpha_{n+1}} \frac{Q_{n+1}(x)Q_n(t) - Q_{n+1}(t)Q_n(x)}{x - t} \quad \forall x, t \in \mathbb{R} \text{ with } x \neq t, \quad (1.51)$$

where  $\alpha_k$  is the leading coefficient of  $Q_k$  ( $k = 0, 1, \dots$ ).

The formula (1.51) is called the *Christoffel–Darboux identity*.

*Proof of Theorem 1.25.* By Theorem 1.23, the orthonormal polynomials  $Q_n$  ( $n = 0, 1, \dots$ ) satisfy (1.48) with  $A_n = \alpha_n/\alpha_{n-1}$  and  $C_n = A_n/A_{n-1}$ , where  $\alpha_n$  is the leading coefficient of  $Q_n$ . Therefore, for a fixed  $n \geq 1$ ,

$$\begin{aligned} &A_{k+1}^{-1} [Q_{k+1}(x)Q_k(t) - Q_{k+1}(t)Q_k(x)] \\ &= A_{k+1}^{-1} [(A_{k+1}x + B_{k+1})Q_k(x) - C_{k+1}Q_{k-1}(x)] Q_k(t) \\ &\quad - A_{k+1}^{-1} [(A_{k+1}t + B_{k+1})Q_k(t) - C_{k+1}Q_{k-1}(t)] Q_k(x) \\ &= (x - t)Q_k(x)Q_k(t) + A_k^{-1} [Q_k(x)Q_{k-1}(t) - Q_k(t)Q_{k-1}(x)], \quad k = 1, \dots, n. \end{aligned}$$

Summing over these, we obtain that

$$\begin{aligned}
& A_{n+1}^{-1} [Q_{n+1}(x)Q_n(t) - Q_{n+1}(t)Q_n(x)] \\
&= (x-t) \sum_{k=1}^n Q_k(x)Q_k(t) + A_1^{-1} [Q_1(x)Q_0(t) - Q_1(t)Q_0(x)] \\
&= (x-t) \sum_{k=1}^n Q_k(x)Q_k(t) + (x-t)Q_0(x)Q_0(t),
\end{aligned}$$

since  $Q_0(x) = Q_0(t) = \alpha_0$ . This leads to (1.51) for  $n \geq 2$ . For the case  $n = 0$  or  $1$ , one can directly verify that (1.51) is true.  $\square$

## 1.9 Legendre Polynomials

The Legendre polynomials  $P_n \in \mathcal{P}_n$  ( $n = 0, 1, \dots$ ) are the unique orthogonal polynomials in  $L^2(-1, 1)$  that are normalized by

$$P_n(1) = 1 \quad \forall n \geq 0.$$

A convenient way to define these polynomials is to use *Rodrigues' formula*:

$$P_n(x) = \frac{1}{2^n n!} \frac{d^n}{dx^n} [(x^2 - 1)^n], \quad n = 0, 1, \dots$$

Clearly, for each integer  $n \geq 0$ ,  $P_n$  is a polynomial of degree exactly  $n$ . Moreover, if  $n$  is even (odd), then  $P_n$  is an even (odd) polynomial, i.e., it only consists of even (odd) powers of  $x$ . The first few of these polynomials are

$$\begin{aligned}
P_0(x) &= 1, \\
P_1(x) &= x, \\
P_2(x) &= \frac{3}{2}x^2 - \frac{1}{2}, \\
P_3(x) &= \frac{5}{2}x^3 - \frac{3}{2}x, \\
P_4(x) &= \frac{35}{8}x^4 - \frac{15}{4}x^2 + \frac{3}{8}.
\end{aligned}$$

We summarize in the following theorem some of the important properties of Legendre polynomials:

### Theorem 1.26 (Properties of Legendre polynomials).

(1) Orthogonality.

$$\int_{-1}^1 P_m(x)P_n(x) dx = \begin{cases} 0 & \text{if } m \neq n, \\ 2/(2n+1) & \text{if } m = n. \end{cases} \quad (1.52)$$

(2) Recurrence.

$$\begin{aligned} P_0(x) &= 1, \\ P_1(x) &= x, \\ (n+1)P_{n+1}(x) - (2n+1)xP_n(x) + nP_{n-1}(x) &= 0, \quad n = 1, 2, \dots \end{aligned} \quad (1.53)$$

(3) Zeros. For each  $n \geq 1$ ,  $P_n$  has  $n$  simple roots in  $(-1, 1)$ .

(4) The least-squares approximation. For each  $n \geq 1$ ,  $\tilde{P}_n := [2^n(n!)^2/(2n)!]P_n \in \mathcal{P}_n$  is the unique polynomial in  $\tilde{\mathcal{P}}_n$  such that

$$\|\tilde{P}_n\|_{L^2(-1,1)} = \frac{2^n(n!)^2}{(2n)!} \sqrt{\frac{2}{2n+1}} = \min_{\tilde{p}_n \in \tilde{\mathcal{P}}_n} \|\tilde{p}_n\|_{L^2(-1,1)}.$$

(5) Normalization. For each  $n \geq 0$ ,

$$P_n(1) = (-1)^n P_n(-1) = 1.$$

(6) Differential equation. For all  $n \geq 0$ ,

$$(1-x^2)P_n''(x) - 2xP_n'(x) + n(n+1)P_n(x) = 0. \quad (1.54)$$

(7) The generating function. For any  $x \in [-1, 1]$ , the values  $P_n(x)$  ( $n = 0, 1, \dots$ ) are the coefficients of the Maclaurin series

$$\frac{1}{\sqrt{1-2tx+t^2}} = \sum_{n=0}^{\infty} P_n(x)t^n \quad \forall t \in (-1, 1).$$

*Proof.* We only prove Parts (1)–(6). The proof of Part (7) can be found in [9].

(1) Let  $m, n$  be non-negative integers. If one of them is 0, then it is clear that (1.52) holds true. Assume  $1 \leq m \leq n$ . Let  $\phi_k(x) = (x^2 - 1)^k$  for any integer  $k \geq 0$ . Then,  $P_k = [1/(2^k k!)]\phi_k^{(k)}$ . By integration by parts, we thus have

$$\begin{aligned} \int_{-1}^1 \phi_m^{(m)}(x)\phi_n^{(n)}(x) dx &= \phi_m^{(m)}(x)\phi_n^{(n-1)}(x)\Big|_{x=-1}^{x=1} - \int_{-1}^1 \phi_m^{(m+1)}(x)\phi_n^{(n-1)}(x) dx \\ &= - \int_{-1}^1 \phi_m^{(m+1)}(x)\phi_n^{(n-1)}(x) dx \\ &= -\phi_m^{(m+1)}(x)\phi_n^{(n-2)}(x)\Big|_{x=-1}^{x=1} + (-1)^2 \int_{-1}^1 \phi_m^{(m+2)}(x)\phi_n^{(n-2)}(x) dx \\ &= \dots \\ &= (-1)^m \int_{-1}^1 \phi_m^{(2m)}(x)\phi_n^{(n-m)}(x) dx \end{aligned}$$

$$= (-1)^m (2m)! \int_{-1}^1 \phi_n^{(n-m)}(x) dx.$$

This is 0 if  $m < n$ . For  $m = n \geq 1$ , we have by the change of variable  $x = \cos \theta$  that

$$\begin{aligned} \int_{-1}^1 [\phi_n^{(n)}(x)]^2 dx &= (-1)^n (2n)! \int_{-1}^1 (x^2 - 1)^n dx \\ &= 2(2n)! \int_0^1 (1 - x^2)^n dx \\ &= 2(2n)! \int_0^{\pi/2} (\sin \theta)^{2n+1} d\theta \\ &= \frac{2(2n)!(2n)!!}{(2n+1)!!}. \end{aligned}$$

This leads to

$$\begin{aligned} \int_{-1}^1 [P_n(x)]^2 dx &= \frac{1}{2^{2n}(n!)^2} \int_{-1}^1 [\phi_n^{(n)}(x)]^2 dx \\ &= \frac{2(2n)!(2n)!!}{2^{2n}(n!)^2(2n+1)!!} \\ &= \frac{2}{2n+1}. \end{aligned}$$

(2) Let  $\tilde{P}_n \in \tilde{\mathcal{P}}_n$  ( $n = 0, 1, \dots$ ) be the unique orthogonal polynomials each with leading coefficient 1 in  $L^2(-1, 1)$ . By Theorem 1.23, we have

$$\tilde{P}_n(x) = (x - a_n)\tilde{P}_{n-1}(x) - b_n\tilde{P}_{n-2}(x), \quad n = 2, 3, \dots, \quad (1.55)$$

where

$$\begin{aligned} a_n &= \frac{\int_{-1}^1 x[\tilde{P}_{n-1}(x)]^2 dx}{\int_{-1}^1 [\tilde{P}_{n-1}(x)]^2 dx}, \quad n = 1, 2, \dots, \\ b_n &= \frac{\int_{-1}^1 [\tilde{P}_{n-1}(x)]^2 dx}{\int_{-1}^1 [\tilde{P}_{n-2}(x)]^2 dx} \quad n = 2, 3, \dots \end{aligned}$$

By Corollary 1.22,  $\tilde{P}_n$  and  $P_n$  differ only by a constant. Comparing their leading coefficients, we have

$$\tilde{P}_n = \frac{2^n(n!)^2}{(2n)!} P_n, \quad n = 0, 1, \dots \quad (1.56)$$

In particular,  $\tilde{P}_n \in \tilde{\mathcal{P}}_n$  is even (odd) as  $P_n$  if  $n$  is even (odd). Hence, the function  $x[\tilde{P}_{n-1}]^2$  is an odd function, and its integral over  $(-1, 1)$  vanishes. Therefore, all  $a_n = 0$ . By (1.56)

and (1.52),

$$b_n = \frac{(n-1)^2 \int_{-1}^1 [P_{n-1}(x)]^2 dx}{(2n-3)^2 \int_{-1}^1 [P_{n-2}(x)]^2 dx} = \frac{(n-1)^2}{(2n-1)(2n-3)}, \quad n = 2, 3, \dots$$

Consequently, we have by (1.55) with  $n$  replaced by  $n+1$  that

$$\tilde{P}_{n+1}(x) = x\tilde{P}_n(x) - \frac{n^2}{4n^2-1}\tilde{P}_{n-1}(x), \quad n = 2, 3, \dots$$

This, together with (1.56), leads to (1.53).

(3) This follows from Theorem 1.24.

(4) This follows from Theorem 1.21 and (1.52).

(5) The fact that  $P_n(1) = 1$  follows from the three-term recurrence (1.53) with argument of induction. Since  $P_n$  is an even (odd) function if  $n$  is even (odd),  $P_n(-1) = (-1)^n P_n(1) = (-1)^n$ .

(6) For  $n = 1$ , (1.54) is clearly true. We thus assume that  $n \geq 2$ . Let  $q \in \mathcal{P}_{n-1}$ . By integration by parts and the fact that  $P_n$  is orthogonal to all polynomials in  $\mathcal{P}_{n-1}$ , we have

$$\begin{aligned} \int_{-1}^1 [(1-x^2)P_n'(x)]' q(x) dx &= - \int_{-1}^1 (1-x^2)P_n'(x)q'(x) dx \\ &= \int_{-1}^1 [(1-x^2)q'(x)]' P_n(x) dx \\ &= 0. \end{aligned}$$

Therefore,  $((1-x^2)P_n'(x))'$  is orthogonal to all polynomials in  $\mathcal{P}_{n-1}$ . It thus follows from Corollary 1.22 that

$$\alpha_n P_n(x) = ((1-x^2)P_n'(x))' = (1-x^2)P_n''(x) - 2xP_n'(x).$$

Let  $c_n$  be the leading coefficient of  $P_n$ . We have

$$\begin{aligned} ((1-x^2)P_n'(x))' &= c_n [(1-x^2)P_n''(x) - 2xP_n'(x)] \\ &= c_n \{(1-x^2)[n(n-1)x^{n-2} + \dots] - 2x(nx^{n-1} + \dots)\} \\ &= c_n n(n+1)(x^n + \dots). \end{aligned}$$

These two equations imply (1.54) for  $n \geq 2$ . □

## Exercises

1. Let  $B_n f \in \mathcal{P}_n$  ( $n = 0, 1, \dots$ ) be the Bernstein polynomials of  $f \in C[0, 1]$ .

(a) Let  $f_0(x) = 1$ ,  $f_1(x) = x$ , and  $f_2(x) = x^2$ . Show that

$$B_n f_0(x) = 1, \quad B_n f_1(x) = x, \quad B_n f_2(x) = \frac{n-1}{n}x^2 + \frac{1}{n}x, \quad \forall x \in [0, 1].$$

- (b) In general, is  $B_n f \in \mathcal{P}_n$  the best uniform approximation of  $f \in C[0, 1]$  in  $\mathcal{P}_n$  on  $[0, 1]$ ?
2. Let  $B_n f \in \mathcal{P}_n$  ( $n = 0, 1, \dots$ ) be the Bernstein polynomials of  $f \in C[0, 1]$ . Prove the following:
- (1)  $\|(B_n f)' - f'\|_{C[a,b]} \rightarrow 0$  as  $n \rightarrow \infty$  for any  $f \in C^1[0, 1]$ ;
  - (2) Let  $k \geq 1$  be any integer. Then  $\|(B_n f)^{(j)} - f^{(j)}\|_{C[0,1]} \rightarrow 0$  as  $n \rightarrow \infty$  for any  $f \in C^k[0, 1]$  and  $j \in \{1, \dots, k\}$ .
3. Let  $0 < a < b < 1$  and  $f \in C[a, b]$ . Show that there exist a sequence of polynomials with integer coefficients that converge to  $f$  in the  $C[a, b]$ -norm.
4. Denote  $\|f\| = \|f\|_{C[a,b]}$ . Show that for any  $f, g \in C[a, b]$

$$\|f + g\| \leq \|f\| + \|g\| \quad \text{and} \quad \left| \|f\| - \|g\| \right| \leq \|f - g\|.$$

5. Prove Theorem 1.4.
6. Prove Proposition 1.5.
7. Let  $n \geq 0$  be an integer and  $a, b$  two real numbers with  $a < b$ . Define

$$F(c) = \max_{a \leq x \leq b} \left| \sum_{k=0}^n c_k x^k \right| \quad \forall c = (c_0, \dots, c_n) \in \mathbb{R}^{n+1}.$$

Show that  $F : \mathbb{R}^{n+1} \rightarrow \mathbb{R}$  is a continuous function.

8. Let  $k \geq 1$  be an integer and  $f \in C^k[a, b]$ . Show that, for any  $\epsilon > 0$ , there exists  $p \in \mathcal{P}$  such that

$$\|f - p\|_{C[a,b]} < \epsilon, \quad \|f' - p'\|_{C[a,b]} < \epsilon, \quad \dots, \quad \|f^{(k)} - p^{(k)}\|_{C[a,b]} < \epsilon.$$

9. Let  $f \in C[a, b]$  but  $f \notin \mathcal{P}$ . Show that there exists no polynomial  $p \in \mathcal{P}$  such that

$$\|f - p\|_{C[a,b]} \leq \|f - q\|_{C[a,b]} \quad \forall q \in \mathcal{P}.$$

10. Let  $x_0, \dots, x_n$  be  $n + 1$  distinct points in  $[a, b]$ . Let  $f \in C[a, b]$ . Does there exist a unique  $p \in \mathcal{P}_m$  such that

$$\max_{0 \leq j \leq n} |f(x_j) - p(x_j)| \leq \max_{0 \leq j \leq n} |f(x_j) - q(x_j)| \quad \forall q \in \mathcal{P}_m?$$

Discuss the cases  $0 \leq m < n$ ,  $m = n$ , and  $m > n$ .

11. Let  $f_1, \dots, f_n$  be  $n$  linearly independent functions in  $C[a, b]$ . Let  $f \in C[a, b]$ . Show that there exists  $p \in \text{Span}\{f_1, \dots, f_n\}$  such that

$$\|f - p\|_{C[a,b]} \leq \|f - q\|_{C[a,b]} \quad \forall q \in \text{Span}\{f_1, \dots, f_n\}.$$

Discuss the uniqueness of such an approximation.

12. Let  $f \in C[a, b]$  and  $q_n \in \mathcal{P}_n$  for some integer  $n \geq 0$ . Suppose  $p_n \in \mathcal{P}_n$  is the best uniform approximation of  $f$  in  $\mathcal{P}_n$ . Prove that  $p_n + q_n$  is the best uniform approximation of  $f + q_n$  in  $\mathcal{P}_n$ .
13. Let  $c > 0$ . Let  $f \in C[-c, c]$  be an even (odd) function. Show that the best uniform approximation of  $f$  in  $\mathcal{P}_n$  for an integer  $n \geq 0$  is also an even (odd) function.
14. Show that  $p_1(x) = x - 1/8$  is the best uniform approximation of  $f(x) = x^2$  in  $\mathcal{P}_1$  on  $[0, 1]$ .
15. Let  $f(x) = x^4$  ( $0 \leq x \leq 1$ ). Find the best uniform approximation of  $f$  in  $\mathcal{P}_1$  on  $[0, 1]$ .
16. Find the best uniform approximation of  $x^{n+2}$  in  $\mathcal{P}_n$  with respect to the  $C[-1, 1]$ -norm.
17. Let  $f \in C[a, b]$  but  $f \notin \mathcal{P}_n$  for some integer  $n \geq 0$ . Let  $p \in \mathcal{P}_n$  be the best uniform approximation of  $f$  in  $\mathcal{P}_n$  on  $[a, b]$ . Can there exist a sequence of strictly increasing points  $\{x_k\}_{k=1}^\infty$  in  $[a, b]$  such that

$$|f(x_k) - p(x_k)| = \|f - p\|_{C[a,b]} \quad \forall k \geq 1?$$

or such that

$$f(x_k) - p(x_k) = (-1)^k \|f - p\|_{C[a,b]} \quad \forall k \geq 1?$$

18. Let  $n \geq 1$  be an integer. Denote by  $\hat{\mathcal{T}}_n$  the set of all functions

$$\hat{T}(x) = a_0 + \sum_{k=1}^n (a_k \cos^k x + b_k \sin^k x)$$

with  $a_0, a_1, \dots, a_n$  and  $b_1, \dots, b_n$  all real numbers.

- (1) Show that  $\mathcal{T}_n \supseteq \hat{\mathcal{T}}_n$ .
- (2) Show that  $\mathcal{T}_2 \neq \hat{\mathcal{T}}_2$ .
19. Show that any nonzero trigonometric polynomial of degree less than or equal to  $n$  can have at most  $2n$  zeros in  $[0, 2\pi)$ .
20. Prove the Second Weierstrass Approximation Theorem by the First Weierstrass Approximation Theorem.
21. Show that the best uniform approximation of an even (odd) function  $f \in C_{2\pi}$  in  $\mathcal{T}_n$  is also an even (odd) function.
22. Given any function  $g$  on  $[a, b]$ , define

$$g^*(\theta) = g\left(\frac{(b-a)\cos\theta + (a+b)}{2}\right) \quad \forall \theta \in (-\infty, \infty).$$

Let  $f \in C[a, b]$  and  $n \geq 0$  be an integer. Let  $p \in \mathcal{P}_n$  and  $T \in \mathcal{T}_n$  satisfy, respectively,

$$\|f - p\|_{C[a,b]} = E_n(f) := \min_{q \in \mathcal{P}_n} \|f - q\|_{C[a,b]}$$

and

$$\|f^* - T\|_{C_{2\pi}} = E_n^*(f^*) := \min_{S \in \mathcal{T}_n} \|f^* - S\|_{C_{2\pi}}.$$

Show that  $E_n(f) = E_n^*(f^*)$  and that  $T = p^*$ .

23. Let  $f \in C[a, b]$ . Let  $\omega_f$  be the modulus of continuity of  $f$  over  $[a, b]$ . Show for any integer  $k \geq 1$  that

$$\omega_f(k\delta) \leq k\omega_f(\delta)$$

and for any  $\lambda > 0$  that

$$\omega_f(\lambda\delta) \leq (\lambda + 1)\omega_f(\delta).$$

24. Let  $f \in C(-\infty, \infty)$ . Define for any  $\delta > 0$

$$R_f(\delta) = \frac{1}{\delta} \int_{x-\frac{\delta}{2}}^{x+\frac{\delta}{2}} f(t) dt.$$

Show that

$$|f(x) - R_f(\delta)(x)| \leq \omega_f\left(\frac{\delta}{2}\right) \quad \forall x \in (-\infty, \infty), \delta > 0,$$

where  $\omega_f(\delta)$  is the modulus of continuity of  $f$ .

25. Show that there exists a constant  $K = K(a, b) > 0$  independent of  $f$  such that

$$E_n(f) \leq \frac{K}{n} E_{n-1}(f') \quad \forall f \in C^1[a, b], \forall n \geq 1,$$

where for any integer  $k \geq 0$

$$E_k(f) = \min_{q \in \mathcal{P}_k} \|f - q\|_{C[a, b]}.$$

Let  $f \in C^p[a, b]$  for some integer  $p \geq 1$ . Show that there exists a constant  $K = K(a, b, p) > 0$  independent of  $f$  such that

$$E_n(f) \leq \frac{K \|f^{(p)}\|_{C[a, b]}}{n^p} \quad \forall n \geq p.$$

26. Show that

$$\frac{2}{\pi} < \frac{\sin t}{t} < 1 \quad \forall t \in \left(0, \frac{\pi}{2}\right).$$

27. Let  $f \in C_{2\pi}$ . Let

$$a_k = \frac{1}{\pi} \int_{-\pi}^{\pi} f(x) \cos kx dx \quad \text{and} \quad b_k = \frac{1}{\pi} \int_{-\pi}^{\pi} f(x) \sin kx dx$$

be the Fourier coefficients of  $f$ . Let

$$S_n f(x) = \frac{a_0}{2} + \sum_{k=1}^n (a_k \cos kx + b_k \sin kx)$$

be the partial sum of the Fourier series of  $f$  for any integer  $n \geq 1$ .

(a) Show that

$$S_n f(x) = \frac{1}{\pi} \int_{-\pi}^{\pi} f(t+x) \frac{\sin\left(n + \frac{1}{2}\right)t}{2 \sin \frac{t}{2}} dt$$

and that

$$\|S_n f\|_{C_{2\pi}} \leq \lambda_n \|f\|_{C_{2\pi}},$$

where

$$\lambda_n = \frac{1}{\pi} \int_{-\pi}^{\pi} \left| \frac{\sin\left(n + \frac{1}{2}\right)t}{2 \sin \frac{t}{2}} \right| dt.$$

(b) Show that

$$\|f - S_n f\|_{C_{2\pi}} \leq (1 + \lambda_n) E_n^*(f),$$

where

$$E_n^*(f) = \min_{T \in \mathcal{T}_n} \|f - T\|_{C_{2\pi}}.$$

Show also that

$$\frac{4 \log n}{\pi^2} < \lambda_n < 2 + \log n.$$

(c) Show that

$$\|f - S_n f\|_{C_{2\pi}} \leq (3 + \log n) E_n^*(f).$$

28. Prove that

$$T_m(T_n(x)) = T_{mn}(x) \quad \forall m, n \geq 0.$$

29. Prove that

$$\int_{-1}^1 [T_n(x)]^2 dx = 1 - \frac{1}{4n^2 - 1} \quad \forall n \geq 0.$$

30. Calculate  $T'_n(\pm 1)$  for any integer  $n \geq 0$ .

31. (Chebyshev) Let  $n \geq 0$  be an integer and  $T_n$  the  $n$ th Chebyshev polynomial of first kind. Let  $P \in \mathcal{P}_n$  satisfy that  $|P(x)| \leq 1$  for all  $x \in [-1, 1]$ . Show that

$$|P(y)| \leq |T_n(y)| \quad \forall y \notin [-1, 1].$$

32. Prove the following properties of the Chebyshev Polynomials of second kind

$$U_n(x) = \frac{\sin(n+1)\theta}{\sin \theta}, \quad n = 0, 1, \dots,$$

where  $x = \cos \theta$  ( $\theta \in [0, \pi]$ ):

(a) *Recursion formula.*

$$\begin{aligned} U_0(x) &= 1, & U_1(x) &= 2x, \\ U_{n+1}(x) &= 2xU_n(x) - U_{n-1}(x), & n &= 1, 2, \dots; \end{aligned}$$

(b) *Orthogonality.*

$$\int_{-1}^1 \sqrt{1-x^2} U_m(x) U_n(x) dx = \begin{cases} 0 & \text{if } m \neq n, \\ \pi/2 & \text{if } m = n; \end{cases}$$

(c) *Differential equations.*

$$(1-x^2) U_n''(x) - 3x U_n'(x) + n(n+2) U_n(x) = 0 \quad n = 0, 1, \dots;$$

(d) *Relations with the Chebyshev polynomials of first kind.*

$$\begin{aligned} n U_{n-1}(x) &= T_n'(x), & n &= 1, 2, \dots, \\ U_n(x) &= x U_{n-1}(x) + T_n(x), & n &= 1, 2, \dots; \end{aligned}$$

(e) For each  $n \geq 0$ ,  $U_n$  is a polynomial of degree  $n$  with leading coefficient  $2^n$ .

Moreover, if  $n$  is even (odd), then  $U_n$  is an even (odd) polynomial.

33. Define  $\chi(x) = -1$  if  $-1 \leq x < 0$  and  $\chi(x) = 1$  if  $0 \leq x \leq 1$ .

(a) Show that

$$\inf_{f \in C[-1,1]} \sup_{-1 \leq x \leq 1} |f(x) - \chi(x)| = 1,$$

and that there exist infinitely many  $f \in C[-1, 1]$  such that

$$\sup_{-1 \leq x \leq 1} |f(x) - \chi(x)| = 1.$$

(b) Show that

$$\inf_{f \in C[-1,1]} \int_{-1}^1 |f(x) - \chi(x)|^2 dx = 0,$$

and that there exists no  $f \in C[-1, 1]$  such that

$$\int_{-1}^1 |f(x) - \chi(x)|^2 dx = 0.$$

34. Let  $\mathfrak{S}$  be an inner product space and define  $\|f\| = \sqrt{\langle f, f \rangle}$  for any  $f \in \mathfrak{S}$ . Prove the following.

(a) *Triangle inequality.*

$$\|f + g\| \leq \|f\| + \|g\| \quad \forall f, g \in \mathfrak{S}.$$

(b) *The Pythagoras Law.* For any  $f, g \in \mathfrak{S}$ ,

$$\|f + g\|^2 = \|f\|^2 + \|g\|^2 \quad \text{if and only if} \quad \langle f, g \rangle = 0.$$

(c) *Parallelogram Law.*

$$\|f + g\|^2 + \|f - g\|^2 = 2\|f\|^2 + 2\|g\|^2 \quad \forall f, g \in \mathfrak{S}.$$

35. Show that

$$\int_0^1 |f(x)| dx \leq \sqrt{\int_0^1 |f(x)|^2 dx} \quad \forall f \in C[0, 1].$$

36. Let  $f_1, \dots, f_n$  be  $n$  elements in an inner product space  $\mathcal{S}$ . Prove that  $f_1, \dots, f_n$  are linearly independent if and only if the Gram matrix

$$G(f_1, \dots, f_n) := (\langle f_i, f_j \rangle) \in \mathbb{R}^{n \times n}$$

is symmetric positive definite.

37. Let  $\mathcal{S}$  be an inner product space. Let  $f_1, \dots, f_n$  be  $n$  linearly independent vectors in  $\mathcal{S}$  and  $\mathcal{S}_n = \text{Span}\{f_1, \dots, f_n\}$ . Let  $f \in \mathcal{S}$ . Prove the following:

(a) There exists a unique  $p \in \mathcal{S}_n$  such that

$$\|f - p\| = \min_{q \in \mathcal{S}_n} \|f - q\|.$$

This  $p$  is called the *least-squares approximation* of  $f$  in  $\mathcal{S}_n$ ;

(b) The least-squares approximation  $p \in \mathcal{S}_n$  of  $f$  is characterized by

$$\langle f - p, q \rangle = 0 \quad \forall q \in \mathcal{S}_n;$$

(c) The error of the least-squares approximation is given by

$$\|f - p\|^2 = \|f\|^2 - \|p\|^2.$$

38. Let  $\{f_1, \dots, f_n\}$  be an orthonormal system in an inner product space  $\mathcal{S}$ . Let  $\mathcal{S}_n = \text{Span}\{f_1, \dots, f_n\}$ . Prove the following.

(a) The vectors  $f_1, \dots, f_n$  are linearly independent.

(b) Any  $q \in \mathcal{S}_n$  has the unique expression

$$q = \sum_{k=1}^n \langle q, f_k \rangle f_k.$$

Moreover,

$$\|q\|^2 = \sum_{k=1}^n \langle q, f_k \rangle^2.$$

(c) The least squares approximation of a given  $f \in \mathcal{S}$  in  $\mathcal{S}_n$  is given by

$$p = \sum_{k=1}^n \langle f, f_k \rangle f_k.$$

Moreover,

$$\|f - p\|^2 = \|f\|^2 - \sum_{k=1}^n \langle f, f_k \rangle^2.$$

39. Let  $\{f_n\}_{n=1}^{\infty}$  be an orthonormal system of an inner product space  $\mathcal{S}$ . Prove the Bessel inequality

$$\sum_{n=1}^{\infty} \langle f, f_n \rangle^2 \leq \|f\|^2 \quad \forall f \in \mathcal{S}.$$

40. Find the least-squares approximation of  $f(x) = x^3$  in  $\mathcal{P}_1$  over  $[-1, 1]$ .  
 41. Find the least-squares approximation of  $f(x) = x^4$  in  $\mathcal{P}_1$  over  $[0, 1]$ .  
 42. Let  $p(x) = \sum_{k=0}^n a_k x^k \in \mathcal{P}_n$  be the least squares approximation of a given  $f \in C[0, 1]$  over  $[0, 1]$ . Find the coefficient matrix of the linear system that determines  $a_0, \dots, a_n$ .  
 43. Let  $f \in C[a, b]$  and define

$$\mu_n(f) = \int_a^b x^n f(x) dx, \quad n = 0, 1, \dots$$

Show that  $f(x) = 0$  for all  $x \in [a, b]$  if and only  $\mu_n(f) = 0$  for all  $n = 0, 1, \dots$

44. Let

$$P_n(x) = \frac{1}{2^n n!} \frac{d^n}{dx^n} [(x^2 - 1)^n], \quad n = 0, 1, \dots$$

be the Legendre polynomials.

- (a) Let  $n \geq 1$ . Prove directly by Rolle's Theorem that  $P_n$  has  $n$  simple roots in  $(-1, 1)$ .  
 (b) Let  $r \geq 1$  be an integer. Show that the sequence of corresponding derivatives  $\{P_n^{(r)}\}_{n=r}^{\infty}$  is orthogonal with respect to the weight function  $(1 - x^2)^r$ , i.e.,

$$\int_{-1}^1 P_m^{(r)}(x) P_n^{(r)}(x) (1 - x^2)^r dx = 0 \quad \text{if } m \neq n.$$

45. Let  $n \geq 0$  be an integer and  $P_n$  be defined as in the previous problem.

- (a) Show that  $P'_{n+1}$  has  $n$  distinct roots  $\xi_1, \dots, \xi_n$  in  $(-1, 1)$ .  
 (b) Show that

$$\frac{d}{dx} \left[ x^{n+1} - \sum_{k=0}^n \left( \int_{-1}^1 t^{n+1} P_k(t) dt \right) P_k(x) \right]$$

vanishes at these points  $\xi_1, \dots, \xi_n$ .

46. Let  $\{Q_n\}_{n=0}^{\infty}$  be an orthonormal system of polynomials in  $L^2_{\rho}(a, b)$ . Prove for any  $n \geq 0$  the identity

$$\sum_{k=0}^n [Q_n(x)]^2 = \frac{\alpha_n}{\alpha_{n+1}} [Q'_{n+1}(x) Q_n(x) - Q'_n(x) Q_{n+1}(x)],$$

where  $\alpha_k$  is the leading coefficient of  $Q_k$  ( $k = 0, \dots$ ).

47. Let  $\{Q_n\}_{n=0}^{\infty}$  be an orthogonal system of polynomials in  $L^2_{\rho}(a, b)$ . Let  $n \geq 1$ . Prove that the zeros of  $Q_n$  and that of  $Q_{n+1}$  alternate.

# Chapter 2

## Polynomial Interpolation

### 2.1 Lagrange Interpolation

Let  $n \geq 0$  be an integer,  $x_0, \dots, x_n$  distinct points in a finite interval  $[a, b]$ , and  $y_0, \dots, y_n \in \mathbb{R}$ .

**Definition 2.1 (Larange interpolation).** A Lagrange interpolation polynomial, or Lagrange interpolant, that interpolates  $y_0, \dots, y_n$  at  $x_0, \dots, x_n$  is a polynomial  $p_n \in \mathcal{P}_n$  such that

$$p_n(x_i) = y_i, \quad i = 0, \dots, n. \quad (2.1)$$

The points  $x_0, \dots, x_n$  are called the interpolation points. In the case  $y_i = f(x_i)$  ( $i = 0, \dots, n$ ) for some function  $f : [a, b] \rightarrow \mathbb{R}$ ,  $p_n$  is also called a Lagrange interpolation polynomial, or Lagrange interpolant, of  $f$  at  $x_0, \dots, x_n$ .

**Theorem 2.2 (Existence and uniqueness of Lagrange interpolation).** There exists a unique Lagrange interpolation polynomial that interpolates  $y_0, \dots, y_n$  at  $x_0, \dots, x_n$ .

*Proof.* If  $n = 0$ , then the unique  $p_0 \in \mathcal{P}_0$  is given by  $p_0(x) = y_0$ . Let  $n \geq 1$ . Consider a general polynomial in  $\mathcal{P}_n$

$$p_n(x) = a_0 + a_1x + \dots + a_nx^n,$$

where  $a_i \in \mathbb{R}$  ( $i = 0, \dots, n$ ). Eq. (2.1) is equivalent to the system of linear equations of the unknowns  $a_0, \dots, a_n$

$$a_0 + a_1x_i + \dots + a_nx_i^n = y_i, \quad i = 0, \dots, n. \quad (2.2)$$

The determinant of the coefficient matrix of this linear system is the Vandermonde determinant

$$V(x_0, x_1, \dots, x_n) = \begin{vmatrix} 1 & x_0 & x_0^2 & \dots & x_0^n \\ 1 & x_1 & x_1^2 & \dots & x_1^n \\ \dots & \dots & \dots & \dots & \dots \\ 1 & x_n & x_n^2 & \dots & x_n^n \end{vmatrix} = \prod_{0 \leq i < j \leq n} (x_j - x_i).$$

This is nonzero, since  $x_0, \dots, x_n$  are distinct. Therefore, (2.2) has a unique solution. This implies the desired existence and uniqueness.  $\square$

For  $n \geq 1$ , we define

$$\begin{aligned} l_i(x) &= \frac{(x - x_0) \cdots (x - x_{i-1})(x - x_{i+1}) \cdots (x - x_n)}{(x_i - x_0) \cdots (x_i - x_{i-1})(x_i - x_{i+1}) \cdots (x_i - x_n)} \\ &= \prod_{\substack{j=0 \\ j \neq i}}^n \frac{x - x_j}{x_i - x_j}, \quad i = 0, \dots, n. \end{aligned} \quad (2.3)$$

Clearly, each  $l_i \in \mathcal{P}_n$  has the degree exactly  $n$ . Moreover,

$$l_i(x_j) = \delta_{ij}, \quad i, j = 0, \dots, n. \quad (2.4)$$

If  $n = 0$ , we define  $l_0(x) = 1$ .

**Theorem 2.3 (Lagrange's formula of Lagrange interpolation).** *The unique Lagrange interpolation polynomial  $p_n \in \mathcal{P}_n$  that interpolates  $y_0, \dots, y_n$  at  $x_0, \dots, x_n$  is given by*

$$p_n(x) = y_0 l_0(x) + y_1 l_1(x) + \cdots + y_n l_n(x) = \sum_{i=0}^n y_i l_i(x). \quad (2.5)$$

*Proof.* Let  $p_n$  be given by (2.5). Clearly,  $p_n \in \mathcal{P}_n$ . Moreover, by (2.4),

$$p_n(x_j) = \sum_{i=0}^n y_i l_i(x_j) = \sum_{i=0}^n y_i \delta_{ij} = y_j, \quad j = 0, \dots, n.$$

Thus,  $p_n$  is the Lagrange interpolation polynomial in  $\mathcal{P}_n$ .  $\square$

The formula (2.1) is called *Lagrange's formula* of the Lagrange interpolation.

**Example.** Find the Lagrange interpolation polynomial  $p_2 \in \mathcal{P}_2$  that interpolates  $y_0 = -1, y_1 = 3, y_2 = 2$  at  $x_0 = 0, x_1 = 1, x_2 = 2$ .

We first calculate the polynomials  $l_0, l_1$ , and  $l_2$  associated with the points  $x_0, x_1$ , and  $x_2$ .

$$\begin{aligned} l_0(x) &= \frac{(x - x_1)(x - x_2)}{(x_0 - x_1)(x_0 - x_2)} = \frac{(x - 1)(x - 2)}{(0 - 1)(0 - 2)} = \frac{1}{2}x^2 - \frac{3}{2}x + 1, \\ l_1(x) &= \frac{(x - x_0)(x - x_2)}{(x_1 - x_0)(x_1 - x_2)} = \frac{(x - 0)(x - 2)}{(1 - 0)(1 - 2)} = -x^2 + 2x, \\ l_2(x) &= \frac{(x - x_0)(x - x_1)}{(x_2 - x_0)(x_2 - x_1)} = \frac{(x - 0)(x - 1)}{(2 - 0)(2 - 1)} = \frac{1}{2}x^2 - \frac{1}{2}. \end{aligned}$$

We have now by Lagrange's formula (2.1) that

$$\begin{aligned} p_2(x) &= y_0 l_0(x) + y_1 l_1(x) + y_2 l_2(x) \\ &= (-1) \left( \frac{1}{2}x^2 - \frac{3}{2}x + 1 \right) + 3(-x^2 + 2x) + 2 \left( \frac{1}{2}x^2 - \frac{1}{2} \right) \\ &= -\frac{5}{2}x^2 + \frac{13}{2}x - 1. \end{aligned}$$

We can check that

$$\begin{aligned} p_2(0) &= -\frac{5}{2} \cdot 0^2 + \frac{13}{2} \cdot 0 - 1 = -1, \\ p_2(1) &= -\frac{5}{2} \cdot 1^2 + \frac{13}{2} \cdot 1 - 1 = 3, \\ p_2(2) &= -\frac{5}{2} \cdot 2^2 + \frac{13}{2} \cdot 2 - 1 = 2. \end{aligned}$$

The polynomials  $l_0, \dots, l_n$  defined in (2.3) are called the *Lagrange basis polynomials* associated with the  $n+1$  distinct points  $x_0, \dots, x_n$ . The word "basis" is justified in the first part of the following proposition:

**Proposition 2.4.** (1) *The polynomials  $l_0, \dots, l_n$  defined in (2.3) form a basis of  $\mathcal{P}_n$ .*  
(2) *For any  $p_n \in \mathcal{P}_n$ ,*

$$\sum_{i=0}^n p_n(x_i) l_i(x) = p_n(x) \quad \forall x \in \mathbb{R}, \quad (2.6)$$

$$\sum_{i=0}^n p_n(x - x_i) l_i(x) = p_n(0) \quad \forall x \in \mathbb{R}. \quad (2.7)$$

*Proof.* (1) Let  $c_0, \dots, c_n \in \mathbb{R}$  satisfy  $\sum_{i=0}^n c_i l_i = 0$  in  $\mathcal{P}_n$ , i.e.,  $\sum_{i=0}^n c_i l_i(x) = 0$  for all  $x \in \mathbb{R}$ . Setting  $x = x_j$  for an arbitrary  $j$  with  $0 \leq j \leq n$ , we obtain by (2.4) that  $c_j = 0$ . Thus,  $l_0, \dots, l_n$  are linearly independent in  $\mathcal{P}_n$ , and form a basis of  $\mathcal{P}_n$ , since  $\dim \mathcal{P}_n = n+1$ .

(2) Let  $q_n(x)$  denote the left-hand side of the identity in (2.6). Clearly,  $q_n \in \mathcal{P}_n$ . Moreover, it follows from (2.4) that  $q_n(x_j) = p_n(x_j)$  for all  $j = 0, \dots, n$ . Thus, the polynomial  $p_n - q_n \in \mathcal{P}_n$  vanishes at  $n+1$  distinct points. Since any nonzero polynomial in  $\mathcal{P}_n$  can have at most  $n$  zeros,  $q_n(x) - p_n(x)$  must be identically zero. This proves (2.6).

To prove (2.7), we fix an arbitrary  $x \in \mathbb{R}$ . Notice that  $p_n(x - \cdot) \in \mathcal{P}_n$ . Thus, by (2.6),

$$\sum_{i=0}^n p_n(x - x_i) l_i(t) = p_n(x - t) \quad \forall t \in \mathbb{R}.$$

Setting  $t = x$ , we obtain (2.7). □

For any  $f \in C[a, b]$ , we denote by  $L_n f \in \mathcal{P}_n$  the Lagrange interpolation polynomial that interpolates  $f$  at  $x_0, \dots, x_n$ . We call  $L_n : C[a, b] \rightarrow \mathcal{P}_n$  the *Lagrange interpolation operator*, or simply *Lagrange interpolator*, associated with the  $n + 1$  distinct points  $x_0, \dots, x_n \in [a, b]$ .

**Proposition 2.5.** (1) *Each  $L_n : C[a, b] \rightarrow \mathcal{P}_n$  is a linear operator.*

(2)  *$L_n f = f$  for any  $f \in \mathcal{P}_n$ .*

*Proof.* (1) This follows from Lagrange's formula (2.5).

(2) This follows from (2.5) and (2.6). □

For each  $k \geq 1$ , we denote by  $C^k[a, b]$  the set of functions  $f : [a, b] \rightarrow \mathbb{R}$  that have all the continuous derivatives  $f^{(j)}$  on  $[a, b]$  for  $1 \leq j \leq k$ . The derivatives at the end-points  $a$  and  $b$  are the one-sided derivatives, and the continuity at  $a$  and  $b$  is also one-sided. For convenience, we denote  $C^0[a, b] = C[a, b]$ .

**Theorem 2.6 (The remainder of Lagrange interpolation).** *Let  $x_0, \dots, x_n$  be  $n + 1$  distinct points in  $[a, b]$ . Let  $f \in C^{n+1}[a, b]$  and  $L_n f \in \mathcal{P}_n$  be the Lagrange interpolant of  $f$  at  $x_0, \dots, x_n$ . Then for any  $x \in [a, b]$  there exists  $\xi(x) \in [a, b]$  such that*

$$f(x) - (L_n f)(x) = \frac{f^{(n+1)}(\xi(x))}{(n+1)!} (x - x_0) \cdots (x - x_n) \quad \forall x \in [a, b]. \quad (2.8)$$

*Proof.* Let  $x \in [a, b]$ . If  $x = x_i$  for some  $i$  with  $0 \leq i \leq n$ , then (2.8) holds true for any  $\xi(x) \in [a, b]$ . Assume that  $x \neq x_i$  ( $0 \leq i \leq n$ ). Let  $\omega(t) = \prod_{i=0}^n (t - x_i)$  and define

$$\phi(t) = f(t) - (L_n f)(t) - \lambda \omega(t),$$

where  $\lambda \in \mathbb{R}$  is so chosen that  $\phi(x) = 0$ , i.e.,

$$\lambda = \frac{f(x) - (L_n f)(x)}{\omega(x)}. \quad (2.9)$$

Clearly  $\phi \in C^{n+1}[a, b]$ . Moreover,  $\phi = 0$  at the  $n + 2$  distinct points  $x, x_0, \dots, x_n$  in  $[a, b]$ . Thus, by Rolle's Theorem,

$$\phi' = 0 \text{ at } n + 1 \text{ distinct points in } [a, b],$$

$$\phi'' = 0 \text{ at } n \text{ distinct points in } [a, b],$$

...

$$\phi^{(n)} = 0 \text{ at } 2 \text{ distinct points in } [a, b].$$

Finally, there exists  $\xi(x) \in [a, b]$  such that  $\phi^{(n+1)}(\xi(x)) = 0$ . By the definition of  $\phi$ , we have

$$\phi^{(n+1)}(t) = f^{(n+1)}(t) - \lambda(n+1)!$$

Hence,

$$\phi^{(n+1)}(\xi(x)) = f^{(n+1)}(\xi(x)) - \lambda(n+1)! = 0$$

This, together with (2.9), implies (2.8). □

Consider now the special case  $[a, b] = [-1, 1]$ . Let  $f \in C^{n+1}[-1, 1]$ . By the above theorem, we have

$$|f(x) - (L_n f)(x)| \leq \frac{1}{(n+1)!} \left( \max_{a \leq x \leq b} |f^{(n+1)}(x)| \right) |\omega(x)| \quad \forall x \in [-1, 1], \quad (2.10)$$

where  $L_n : C[-1, 1] \rightarrow \mathcal{P}_n$  is the Lagrange interpolator associated with a given set of  $n+1$  distinct points  $x_0, \dots, x_n$  in  $[-1, 1]$  and  $\omega(x) = \prod_{k=0}^n (x - x_k)$ .

In order to minimize the error  $f - L_n f$  for all  $f \in C^{n+1}[-1, 1]$  with respect to the  $C[-1, 1]$ -norm, we choose  $x_0, \dots, x_n \in [-1, 1]$  to minimize the  $C[-1, 1]$ -norm of  $\omega$ . Since  $\omega \in \tilde{\mathcal{P}}_n$ , it follows from Theorem 1.10 on properties of Chebyshev polynomials that the optimal choice of  $\omega$  is the rescaled Chebyshev polynomial:

$$\omega(x) = \tilde{T}_{n+1}(x) = 2^{-n} T_{n+1}(x) \quad \forall x \in [-1, 1]. \quad (2.11)$$

In particular, the unique set of optimal interpolation points are the roots of Chebyshev polynomials  $T_{n+1}$ :

$$x_k = \cos \frac{(2k+1)\pi}{2(n+1)}, \quad k = 0, \dots, n.$$

Moreover,

$$\|\omega\|_{C[-1,1]} = \|\tilde{T}_{n+1}\|_{C[-1,1]} = \|2^{-n} T_{n+1}\|_{C[-1,1]} = 2^{-n}. \quad (2.12)$$

Therefore, with such a choice of interpolation points, we have the error estimate

$$\|f - L_n f\|_{C[-1,1]} \leq \frac{1}{2^n (n+1)!} \|f^{(n+1)}\| \quad \forall f \in C^{n+1}[-1, 1]. \quad (2.13)$$

If we set  $f = \tilde{T}_{n+1}$ , then the unique Lagrange interpolation polynomial in  $\mathcal{P}_n$  of  $f$  at all the roots of  $T_{n+1}$  is the zero polynomial. By (2.12), the equality in (2.13) holds true. Hence, the error estimate (2.13) is optimal.

Now consider the interpolation error in the  $L_\rho^2(a, b)$ -norm for some weight function  $\rho$  on  $[a, b]$ . Let  $f \in C^{n+1}[a, b]$ . By (2.10), we have

$$\|f - L_n f\|_{L_\rho^2(a,b)} \leq \frac{1}{(n+1)!} \|f^{(n+1)}\|_{C[a,b]} \|\omega\|_{L_\rho^2(a,b)}, \quad (2.14)$$

where again  $L_n : C[a, b] \rightarrow \mathcal{P}_n$  is the Lagrange interpolator associated with a given set of  $n+1$  distinct points  $x_0, \dots, x_n \in [a, b]$  and  $\omega(x) = \prod_{k=0}^n (x - x_k)$ . By Theorem 1.21, the unique set of optimal interpolation points  $x_0, \dots, x_n$  are the  $n+1$  distinct roots of the  $(n+1)$ st orthogonal polynomial  $Q_{n+1} \in \tilde{\mathcal{P}}_{n+1}$  in  $L_\rho^2(a, b)$  i.e.,  $\omega = Q_{n+1}$ . Thus, it follows from (2.14) that

$$\|f - L_n f\|_{L_\rho^2(a,b)} \leq \frac{1}{(n+1)!} \|f^{(n+1)}\|_{C[a,b]} \|Q_{n+1}\|_{L_\rho^2(a,b)} \quad \forall f \in C^{n+1}[a, b].$$

Taking  $f = Q_{n+1}$ , we have  $L_n f = 0$ . Hence, this estimate is optimal.

In the special case  $[a, b] = [-1, 1]$  and  $\rho(x) = 1/\sqrt{1-x^2}$ , we see from Theorem 1.10 that the unique set of optimal interpolation points in  $[-1, 1]$  are the zeros of Chebyshev polynomial  $T_{n+1}$  and that  $\omega = 2^{-n}T_{n+1}$ . By (1.28), we have

$$\|\omega\|_{L^2_\rho(-1,1)} = \|2^{n+1}T_{n+1}\|_{L^2_\rho(-1,1)} = \frac{\sqrt{\pi}}{2^{n+1/2}}.$$

Therefore, we have the optimal estimate

$$\|f - L_n f\|_{L^2_\rho(-1,1)} \leq \frac{\sqrt{\pi}}{2^{n+1/2}(n+1)!} \|f^{(n+1)}\|_{C[-1,1]} \quad \forall f \in C^{n+1}[-1, 1].$$

In the special case  $[a, b] = [-1, 1]$  and  $\rho(x) = 1$ , the unique set of optimal interpolation points in  $[-1, 1]$  are the zeros of Legendre polynomial  $P_{n+1}$  and that  $\omega = [2^n(n!)^2/(2n)!]P_{n+1}$ . By (1.52), we have

$$\|\omega\|_{L^2(-1,1)} = \frac{2^n(n!)^2}{(2n)!} \sqrt{\frac{2}{2n+1}}.$$

Therefore, we have the optimal estimate

$$\|f - L_n f\|_{L^2_\rho(-1,1)} \leq \frac{2^n n!}{(n+1)(2n)!} \sqrt{\frac{2}{2n+1}} \|f^{(n+1)}\|_{C[-1,1]} \quad \forall f \in C^{n+1}[-1, 1].$$

## 2.2 Newton's Formula and Divided Differences

Suppose  $p_k \in \mathcal{P}_k$  is the Lagrange interpolation polynomial that interpolates  $f_0, \dots, f_k$  at  $x_0, \dots, x_k$ . Consider adding one more interpolation point  $x_{k+1} \in \mathbb{R}$  that is different from all  $x_0, \dots, x_k$ , and adding one more value  $f_{k+1} \in \mathbb{R}$ . Let  $p_{k+1} \in \mathcal{P}_{k+1}$  be the Lagrange interpolation polynomial  $p_{k+1} \in \mathcal{P}_{k+1}$  that interpolates  $f_0, \dots, f_k$ , and  $f_{k+1}$  at  $x_0, \dots, x_k$ , and  $x_{k+1}$ . Since  $p_k(x_i) = p_{k+1}(x_i) = f_i$  for  $i = 0, \dots, k$ , we see that the polynomial  $p_{k+1} - p_k \in \mathcal{P}_{k+1}$  vanishes at  $x_0, \dots, x_k$ . Hence, it must have the form

$$p_{k+1}(x) - p_k(x) = d_{k+1}(x - x_0) \cdots (x - x_k)$$

for some  $d_{k+1} \in \mathbb{R}$ . The condition that  $p_{k+1}(x_{k+1}) = f_{k+1}$  determines uniquely that

$$d_{k+1} = \frac{f_{k+1} - p_k(x_{k+1})}{(x_{k+1} - x_0) \cdots (x_{k+1} - x_k)}. \quad (2.15)$$

Therefore, starting from the constant polynomial  $p_0(x) = d_0 \in \mathbb{R}$  that interpolates  $f_0$  at  $x_0$ , we have

$$p_0(x) = d_0,$$

$$\begin{aligned}
p_1(x) &= p_0(x) + d_1(x - x_0), \\
p_2(x) &= p_1(x) + d_2(x - x_0)(x - x_1), \\
&\dots \\
p_k(x) &= p_{k-1}(x) + d_k(x - x_0) \cdots (x - x_{k-1}),
\end{aligned}$$

where  $d_0, \dots, d_k \in \mathbb{R}$  are constants. Finally,

$$p_k(x) = d_0 + d_1(x - x_0) + \cdots + d_k(x - x_0) \cdots (x - x_{k-1}).$$

**Definition 2.7 (Divided differences).** *The divided differences of a given set of numbers  $f_0, \dots, f_n$  at  $n + 1$  distinct points  $x_0, \dots, x_n \in \mathbb{R}$  are*

$$\begin{aligned}
f[x_0] &= f_0, \\
f[x_0, \dots, x_k] &= \frac{f[x_1, \dots, x_k] - f[x_0, \dots, x_{k-1}]}{x_k - x_0}, \quad k = 2, \dots, n.
\end{aligned}$$

If  $f_i = f(x_i)$  ( $i = 0, \dots, n$ ) for some function  $f$  that is defined on a set of real numbers containing all  $x_0, \dots, x_n$ , then  $f[x_0], \dots, f[x_0, \dots, x_n]$  are called the divided differences of the function  $f$  at these points  $x_0, \dots, x_n$ .

**Theorem 2.8 (Newton's formula of Lagrange interpolation).** *Let  $x_0, \dots, x_n \in \mathbb{R}$  be  $n + 1$  distinct points and  $f_0, \dots, f_n \in \mathbb{R}$ . Then, for each integer  $k$  with  $0 \leq k \leq n$ ,*

$$p_k(x) = f[x_0] + f[x_0, x_1](x - x_0) + \cdots + f[x_0, \dots, x_k](x - x_0) \cdots (x - x_{k-1}) \quad (2.16)$$

is the Lagrange interpolation polynomial that interpolates  $f_0, \dots, f_k$  at  $x_0, \dots, x_k$ .

The formula (2.16) is called the *Newton's formula* of the Lagrange interpolation.

**Example.** Use Newton's formula to find the Lagrange interpolation polynomial  $p_2 \in \mathcal{P}_2$  that interpolates  $f_0 = -1, f_1 = 3, f_2 = 2$  at  $x_0 = 0, x_1 = 1, x_2 = 2$ .

We first calculate all the needed divided differences.

$$\begin{aligned}
f[x_0] &= f_0 = -1, \\
f[x_1] &= f_1 = 3, \\
f[x_2] &= f_2 = 2, \\
f[x_0, x_1] &= \frac{f[x_1] - f[x_0]}{x_1 - x_0} = \frac{3 - (-1)}{1 - 0} = 4, \\
f[x_1, x_2] &= \frac{f[x_2] - f[x_1]}{x_2 - x_1} = \frac{2 - 3}{2 - 1} = -1, \\
f[x_1, x_1, x_2] &= \frac{f[x_1, x_2] - f[x_0, x_1]}{x_2 - x_0} = \frac{-1 - 4}{2 - 0} = -\frac{5}{2}.
\end{aligned}$$

By Newton's formula (2.16), we have

$$\begin{aligned} p_2(x) &= f[x_0] + f[x_0, x_1](x - x_0) + f[x_0, x_1, x_2](x - x_0)(x - x_1) \\ &= -1 + 4(x - 0) + \left(-\frac{5}{2}\right)(x - 0)(x - 1) \\ &= -\frac{5}{2}x^2 + \frac{13}{2}x - 1. \end{aligned}$$

This is the same polynomial as obtained in the example in Section 2.1.

To prove Theorem 2.8, we first prove the following useful lemma:

**Lemma 2.9.** *Suppose  $p_k, q_k \in \mathcal{P}_k$  are the Lagrange interpolation polynomials that interpolate  $f_0, \dots, f_k$  at  $x_0, \dots, x_k$  and  $f_1, \dots, f_{k+1}$  at  $x_1, \dots, x_{k+1}$ , respectively. Then,*

$$r_{k+1}(x) = \frac{(x - x_0)q_k(x) - (x - x_{k+1})p_k(x)}{x_{k+1} - x_0} \quad (2.17)$$

is the Lagrange interpolation polynomial that interpolates  $f_0, \dots, f_k$ , and  $f_{k+1}$  at  $x_0, \dots, x_k$ , and  $x_{k+1}$ .

*Proof.* Since  $p_k(x_i) = q_k(x_i)$  for  $i = 1, \dots, k$ , we have by a direct calculation by (2.17) that  $r_{k+1}(x_i) = f_i$  for  $i = 1, \dots, k$ . Also by (2.17) we have  $r_{k+1}(x_0) = p_k(x_0) = f_0$  and  $r_{k+1}(x_{k+1}) = q_k(x_{k+1}) = f_{k+1}$ . Therefore,  $r_{k+1} \in \mathcal{P}_{k+1}$  is the Lagrange interpolation polynomial that interpolates  $f_0, \dots, f_{k+1}$  at  $x_0, \dots, x_{k+1}$ .  $\square$

*Proof of Theorem 2.8.* We prove this theorem by the induction on  $k$ , the number of interpolation points. For  $k = 0$ , clearly  $p_0(x) = d_0 = f_0$  is the Lagrange interpolation polynomial that interpolates  $f_0$  at  $x_0$ . Fix an integer  $k \geq 1$  and assume that

$$p_j(x) = f[x_0] + f[x_0, x_1](x - x_0) + \dots + f[x_0, \dots, x_j](x - x_0) \dots (x - x_{j-1})$$

interpolates  $f_0, \dots, f_j$  at  $x_0, \dots, x_j$  for each  $j = 0, \dots, k$ . We need to show that (2.16) holds true with  $k$  replaced by  $k + 1$ .

*Step 1.* By the assumption of induction, the polynomial

$$p_k(x) = f[x_0] + f[x_0, x_1](x - x_0) + \dots + f[x_0, \dots, x_k](x - x_0) \dots (x - x_{k-1}) \quad (2.18)$$

is the Lagrange interpolation polynomial that interpolates  $f_0, \dots, f_k$  at  $x_0, \dots, x_k$ . Let  $d_{k+1}$  be given as in (2.15) and

$$p_{k+1}(x) = p_k(x) + d_{k+1}(x - x_0) \dots (x - x_k).$$

Clearly,  $p_{k+1}(x_i) = p_k(x_i) = f_i$  for  $i = 0, \dots, k$ , and by (2.15),

$$p_{k+1}(x_{k+1}) = p_k(x_{k+1}) + d_{k+1}(x_{k+1} - x_0) \dots (x_{k+1} - x_k) = f_{k+1}.$$

Thus,  $p_{k+1} \in \mathcal{P}_{k+1}$  is the Lagrange interpolation polynomial that interpolates  $f_0, \dots, f_k$ , and  $f_{k+1}$  at  $x_0, \dots, x_k$ , and  $x_{k+1}$ .

*Step 2.* By the assumption of induction, the polynomial

$$q_k(x) = f[x_1] + f[x_1, x_2](x - x_1) + \cdots + f[x_1, \dots, x_{k+1}](x - x_1) \cdots (x - x_k)$$

is the Lagrange interpolation polynomial that interpolates  $f_1, \dots, f_{k+1}$  at  $x_1, \dots, x_{k+1}$ . Therefore, it follows from Lemma 2.9 that the polynomial  $r_{k+1} \in \mathcal{P}_{k+1}$  defined in (2.17), where  $p_k$  is given in (2.18), is the Lagrange interpolation polynomial that interpolates  $f_0, \dots, f_k$ , and  $f_{k+1}$  at  $x_0, \dots, x_k$ , and  $x_{k+1}$ . Hence, by the uniqueness of Lagrange interpolation,  $r_{k+1} = p_{k+1}$ .

*Step 3.* Comparing the leading coefficients of  $p_{k+1}$  and  $r_{k+1}$ , we obtain

$$d_{k+1} = \frac{f[x_1, \dots, x_{k+1}] - f[x_0, \dots, x_k]}{x_{k+1} - x_0} = f[x_0, \dots, x_{k+1}].$$

Therefore, (2.16) holds true with  $k$  replaced by  $k + 1$ . □

**Theorem 2.10.** *Let  $p_n \in \mathcal{P}_n$  be the Lagrange interpolation polynomial of a given function  $f : [a, b] \rightarrow \mathbb{R}$  at  $x_0, \dots, x_n \in [a, b]$ . Then for any  $x \in [a, b]$  with  $x \neq x_i$  ( $i = 0, \dots, n$ )*

$$f(x) - p_n(x) = f[x_0, \dots, x_n, x](x - x_0) \cdots (x - x_n).$$

*Proof.* Let  $p_{n+1} \in \mathcal{P}_{n+1}$  be the Lagrange interpolation polynomial of  $f$  at the  $n + 2$  points  $x_0, \dots, x_n, x$ . Then by Newton's formula

$$p_{n+1}(t) = p_n(t) + f[x_0, \dots, x_n, x](t - x_0) \cdots (t - x_n)$$

Setting  $t = x$ , we obtain

$$f(x) = p_{n+1}(x) = p_n(x) + f[x_0, \dots, x_n, x](x - x_0) \cdots (x - x_n),$$

completing the proof. □

**Proposition 2.11 (Properties of divided differences).** *Let  $x_0, \dots, x_n$  be  $n + 1$  distinct points in  $[a, b]$ .*

(1) *Linearity.* For any functions  $f, g : \{x_0, \dots, x_n\} \rightarrow \mathbb{R}$  and any  $\alpha, \beta \in \mathbb{R}$ ,

$$(\alpha f + \beta g)[x_0, \dots, x_n] = \alpha f[x_0, \dots, x_n] + \beta g[x_0, \dots, x_n].$$

(2) *Symmetry.* For any function  $f : \{x_0, \dots, x_n\} \rightarrow \mathbb{R}$  and any permutation  $(i_0 \dots i_n)$  of  $(0 \dots n)$ ,

$$f[x_0, \dots, x_n] = f[x_{i_0}, \dots, x_{i_n}]. \tag{2.19}$$

(3) For any function  $f : \{x_0, \dots, x_n\} \rightarrow \mathbb{R}$ ,

$$f[x_0, \dots, x_n] = \sum_{i=0}^n \frac{f_i}{(x_i - x_0) \cdots (x_i - x_{i-1})(x_i - x_{i+1}) \cdots (x_i - x_n)}. \quad (2.20)$$

(4) If  $f \in C^n[a, b]$ , then there exists  $\xi \in [a, b]$  such that

$$f[x_0, \dots, x_n] = \frac{f^{(n)}(\xi)}{n!}. \quad (2.21)$$

(5) If  $f(x) = x^m$  for some integer  $m \geq 0$ , then

$$f[x_0, \dots, x_n] = \begin{cases} 0 & \text{if } m < n, \\ 1 & \text{if } m = n, \\ \text{linear combinations of } x_0^{k_0} \cdots x_n^{k_n} \text{ with } \sum_{i=0}^n k_i = m - n & \text{if } m > n. \end{cases}$$

*Proof.* (1) This follows from the definition of divided differences and an argument by induction.

(2) By the uniqueness of the Lagrange interpolation, the Lagrange interpolation polynomial that interpolates  $f$  at  $x_{i_0}, \dots, x_{i_n}$  is the same as that interpolates  $f$  at  $x_0, \dots, x_n$ . By Newton's formula (2.16), the leading coefficients in these two polynomials are exactly the right-hand side and left-hand side of (2.19), respectively. Thus, they must be the same.

(3) The left-hand side and right-hand side of (2.20) are the leading coefficients in Newton's formula (2.16) and Lagrange's formula (2.5), respectively, of the unique Lagrange interpolation polynomial that interpolates  $f$  at  $x_0, \dots, x_n$ .

(4) This is obviously true for  $n = 0$ . Assume  $n \geq 1$ . Let  $p_{n-1} \in \mathcal{P}_{n-1}$  and  $p_n \in \mathcal{P}_n$  be the Lagrange interpolation polynomials that interpolate  $f_0, \dots, f_{n-1}$  at  $x_0, \dots, x_{n-1}$  and  $f_0, \dots, f_n$  at  $x_0, \dots, x_n$ , respectively. It follows from Newton's formula (2.16) that

$$p_n(x) = p_{n-1}(x) + f[x_0, \dots, x_n](x - x_0) \cdots (x - x_{n-1}).$$

Hence, replacing  $x$  by  $x_n$ , we obtain

$$f(x_n) = p_n(x_n) = p_{n-1}(x_n) + f[x_0, \dots, x_n](x_n - x_0) \cdots (x_n - x_{n-1}).$$

On the other hand, by Theorem 2.6 on the remainder of Lagrange interpolation, we have

$$f(x_n) - p_{n-1}(x_n) = \frac{f^{(n)}(\xi)}{n!}(x_n - x_0) \cdots (x_n - x_{n-1})$$

for some  $\xi \in [a, b]$ . The above two equations imply (2.21).

(5) By (2.21), we need only to consider the case that  $m > n$ . We use the argument by induction. For  $n = 0$ , the statement is clearly true. Assume that for any  $n \geq 0$  the statement is true, i.e.,

$$f[x_0, \dots, x_n] = \sum_{k_0 + \dots + k_n = m-n} \alpha_{k_0 \dots k_n} x_0^{k_0} \dots x_n^{k_n}, \quad (2.22)$$

where  $\alpha_{k_0, \dots, k_n}$  are constants independent of  $x_0, \dots, x_n$ . Assume  $m > n + 1$ . We have by the definition of divided differences, the symmetry property (2.19), and the assumption (2.22) that

$$\begin{aligned} & f[x_0, \dots, x_{n+1}] \\ &= \frac{f[x_1, \dots, x_{n+1}] - f[x_0, \dots, x_n]}{x_{n+1} - x_0} \\ &= \frac{f[x_{n+1}, x_1, \dots, x_n] - f[x_0, x_1, \dots, x_n]}{x_{n+1} - x_0} \\ &= \frac{\sum_{k_0 + \dots + k_n = m-n} \alpha_{k_0 \dots k_n} x_{n+1}^{k_0} x_1^{k_1} \dots x_n^{k_n} - \sum_{k_0 + \dots + k_n = m-n} \alpha_{k_0 \dots k_n} x_0^{k_0} x_1^{k_1} \dots x_n^{k_n}}{x_{n+1} - x_0} \\ &= \sum_{\substack{k_0 + \dots + k_n = m-n \\ k_0 \geq 1}} \alpha_{k_0 \dots k_n} x_1^{k_1} \dots x_n^{k_n} \left( \frac{x_{n+1}^{k_0} - x_0^{k_0}}{x_{n+1} - x_0} \right) \\ &= \sum_{\substack{k_0 + \dots + k_n = m-n \\ k_0 \geq 1}} \alpha_{k_0 \dots k_n} x_1^{k_1} \dots x_n^{k_n} \sum_{k_{n+1}=0}^{k_0-1} x_0^{k_0-1-k_{n+1}} x_{n+1}^{k_{n+1}} \\ &= \sum_{k'_0 + k_1 + \dots + k_{n+1} = m-(n+1)} \alpha_{k'_0 + k_{n+1} + 1, k_2, \dots, k_n} x_0^{k'_0} x_1^{k_1} \dots x_n^{k_n} x_{n+1}^{k_{n+1}}, \end{aligned}$$

where in the last step  $k'_0 = k_0 - 1 - k_{n+1}$ . This proves that the statement is true for  $n + 1$ .  $\square$

For each  $n \geq 1$ , we denote by  $\tau_n$  the unit simplex in  $\mathbb{R}^n$ :

$$\tau_n = \left\{ (t_1, \dots, t_n) \in \mathbb{R}^n : t_i \geq 0 \ (1 \leq i \leq n) \text{ and } \sum_{i=1}^n t_i \leq 1 \right\}.$$

**Theorem 2.12 (The Hermite–Genocchi formula).** *Let  $n \geq 1$  be an integer and  $x_0, \dots, x_n \in [0, 1]$  be distinct. We have for any  $f \in C^n[0, 1]$  that*

$$f[x_0, \dots, x_n] = \int_{\tau_n} f^{(n)} \left( x_0 + \sum_{j=1}^n t_j (x_j - x_0) \right) dt. \quad (2.23)$$

*Proof.* We use the argument of induction. The statement is true for  $n = 1$ , since

$$\int_0^1 f'(x_0 + t_1(x_1 - x_0)) dt_1 = \frac{f(x_1) - f(x_0)}{x_1 - x_0} = f[x_0, x_1].$$

Suppose (2.23) holds true for  $n \geq 1$ . Consider the case of  $n + 1$ . We have

$$\begin{aligned} & \int_{\tau_{n+1}} f^{(n+1)} \left( x_0 + \sum_{j=1}^{n+1} t_j(x_j - x_0) \right) dt_1 \cdots dt_{n+1} \\ &= \int_{\tau_n} \left[ \int_0^{1-\sum_{j=1}^n t_j} f^{(n+1)} \left( x_0 + \sum_{j=1}^{n+1} t_j(x_j - x_0) \right) dt_{n+1} \right] dt_1 \cdots dt_n \\ &= \frac{1}{x_{n+1} - x_0} \int_{\tau_n} f^{(n)} \left( x_{n+1} + \sum_{j=1}^n t_j(x_j - x_{n+1}) \right) dt_1 \cdots dt_n \\ &\quad - \frac{1}{x_{n+1} - x_0} \int_{\tau_n} f^{(n)} \left( x_0 + \sum_{j=1}^n t_j(x_j - x_0) \right) dt_1 \cdots dt_n \\ &= \frac{f[x_{n+1}, x_1, \dots, x_n] - f[x_0, \dots, x_n]}{x_{n+1} - x_0} \\ &= \frac{f[x_1, \dots, x_{n+1}] - f[x_0, \dots, x_n]}{x_{n+1} - x_0}. \end{aligned}$$

Thus, the statement is true for  $n + 1$ . This completes the proof.  $\square$

## 2.3 Peano Kernal and the Remainder Theorem

Let  $x_0, \dots, x_n$  be  $n + 1$  distinct points in  $[a, b]$ . Let  $L_n : C[a, b] \rightarrow \mathcal{P}_n$  be the Lagrange interpolator associated with  $x_0, \dots, x_n$ . We study the error of the Lagrange interpolation  $f - L_n f$  for  $f \in C[a, b]$  that is not necessary smooth enough, e.g.,  $f \notin C^{n+1}[a, b]$ .

To this end, let us introduce for any integer  $k \geq 1$  the function space  $W^{k,1}(a, b)$  that consists of all functions  $f \in C^{k-1}[a, b]$  such that  $f^{(k-1)}$  are absolutely continuous on  $[a, b]$ . If  $f \in W^{k,1}(a, b)$ , then  $f^{(k)}$  exists as an integrable function on  $[a, b]$ . Clearly,  $C^k[a, b] \subset W^{k,1}(a, b)$ .

Let  $f \in W^{m+1,1}(a, b)$  with  $0 \leq m \leq n$ . By the Taylor expansion,

$$\begin{aligned} f(x) &= f(a) + f'(a)(x - a) + \cdots + \frac{f^{(m)}(a)}{m!}(x - a)^m + \frac{1}{m!} \int_a^x (x - t)^m f^{(m+1)}(t) dt \\ &= Q_m(x) + R_m(x), \end{aligned}$$

where

$$Q_m(x) = f(a) + f'(a)(x - a) + \cdots + \frac{f^{(m)}(a)}{m!}(x - a)^m$$

is a polynomial in  $\mathcal{P}_m$  and

$$R_m(x) = \frac{1}{m!} \int_a^x (x-t)^m f^{(m+1)}(t) dt$$

is the remainder. Since  $Q_m \in \mathcal{P}_m \subseteq \mathcal{P}_n$ ,  $L_n Q_m = Q_m$ . Therefore,

$$f - (L_n f) = Q_m + R_m - [L_n Q_m + L_n R_m] = R_m - L_n R_m. \quad (2.24)$$

Let  $l_0, \dots, l_n \in \mathcal{P}_n$  be the Lagrange basis polynomials associated with  $x_0, \dots, x_n$ . By (2.24) and the Lagrange formula (2.5), we have

$$\begin{aligned} f(x) - (L_n f)(x) &= R_m(x) - (L_n R_m)(x) \\ &= \frac{1}{m!} \int_a^x (x-t)^m f^{(m+1)}(t) dt - \sum_{k=0}^n \left[ \frac{1}{m!} \int_a^{x_k} (x_k-t)^m f^{(m+1)}(t) dt \right] l_k(x) \\ &= \frac{1}{m!} \left[ \int_a^b (x-t)_+^m f^{(m+1)}(t) dt - \sum_{k=0}^n \left( \int_a^b (x_k-t)_+^m f^{(m+1)}(t) dt \right) l_k(x) \right] \\ &= \int_a^b \frac{1}{m!} \left[ (x-t)_+^m - \sum_{k=0}^n (x_k-t)_+^m l_k(x) \right] f^{(m+1)}(t) dt, \end{aligned} \quad (2.25)$$

where

$$c_+ = \begin{cases} c & \text{if } c \geq 0, \\ 0 & \text{if } c < 0. \end{cases}$$

We define

$$K_m(x, t) = \frac{1}{m!} \left[ (x-t)_+^m - \sum_{k=0}^n (x_k-t)_+^m l_k(x) \right] = \frac{1}{m!} E_n((\cdot - t)_+^m)(x), \quad (2.26)$$

where  $E_n(g) = g - L_n g$  is the error of the Lagrange interpolation for  $g \in C[a, b]$ . We shall call  $K_m : \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}$  the  $m$ th *Peano kernel* associated with the interpolation points  $x_0, \dots, x_n$ .

We have in fact proved the following:

**Theorem 2.13 (The Peano Remainder Theorem for the Lagrange interpolation).**

Let  $m$  be an integer with  $0 \leq m \leq n$ . Then for any  $f \in W^{m+1,1}(a, b)$

$$f(x) - (L_n f)(x) = \int_a^b K_m(x, t) f^{(m+1)}(t) dt \quad \forall x \in [a, b]. \quad (2.27)$$

To estimate the interpolation error using the Peano kernel representation (2.27), we introduce the function space  $W^{k,\infty}(a, b)$  that consists of all the functions  $f \in C^{k-1}[a, b]$

such that  $f^{(k-1)}$  are Lipschitz continuous on  $[a, b]$ . If  $f \in W^{k,\infty}(a, b)$ , then  $f^{(k)}$  exists as an integrable and bounded function on  $[a, b]$ . It can be proved that for any integer  $k \geq 1$

$$C^k[a, b] \subsetneq W^{k,\infty}(a, b) \subsetneq W^{k,1}(a, b) \subsetneq C[a, b].$$

We denote for a measurable and bounded function  $g : [a, b] \rightarrow \mathbb{R}$

$$\|g\|_{L^\infty(a,b)} = \sup_{a \leq x \leq b} |g(x)|.$$

**Theorem 2.14 (Error estimates for the Lagrange interpolation error).** *Let  $m$  be an integer with  $0 \leq m \leq n$ . Then for any  $f \in W^{m+1,\infty}(a, b)$ ,*

$$|f(x) - (L_n f)(x)| \leq \left[ \int_a^b |K_m(x, t)| dt \right] \|f^{(m+1)}\|_{L^\infty(a,b)} \quad \forall x \in [a, b]. \quad (2.28)$$

Moreover, for any  $x \in [a, b]$ , there exists  $f_0 \in W^{m+1,\infty}(a, b)$  such that

$$|f_0(x) - (L_n f_0)(x)| = \left[ \int_a^b |K_m(x, t)| dt \right] \|f_0^{(m+1)}\|_{L^\infty(a,b)}. \quad (2.29)$$

*Proof.* Let  $f \in W^{m+1,\infty}(a, b)$ . By (2.25) and (2.26), we have

$$|f(x) - (L_n f)(x)| \leq \int_a^b |K_m(x, t)| |f^{(m+1)}(t)| dt \leq \left[ \int_a^b |K_m(x, t)| dt \right] \|f^{(m+1)}\|_{L^\infty(a,b)},$$

implying (2.28). Fix  $x \in [a, b]$ . Define  $g_m(t) = \text{sign } K_m(x, t)$  and

$$f_0(x) = \underbrace{\int_a^t \dots \int_a^t}_{m+1 \text{ times}} g_m(t) \underbrace{dt \dots dt}_{m+1 \text{ times}}.$$

Then,  $f_0 \in W^{m+1,\infty}(a, b)$  and  $\|f_0^{(m+1)}\|_{L^\infty(a,b)} = \|g_m\|_{L^\infty(a,b)} = 1$ . By (2.25) and (2.26), we obtain

$$\begin{aligned} |f_0(x) - (L_n f_0)(x)| &= \left| \int_a^b K_m(x, t) f_0^{(m+1)}(t) dt \right| = \left| \int_a^b K_m(x, t) g_m(t) dt \right| \\ &= \int_a^b |K_m(x, t)| dt = \left[ \int_a^b |K_m(x, t)| dt \right] \|f_0^{(m+1)}\|_{L^\infty(a,b)}, \end{aligned}$$

leading to (2.29). □

**Theorem 2.15.** (1) *We have*

$$\int_a^b K_n(x, t) dt = \frac{1}{(n+1)!} \prod_{k=0}^n (x - x_k) \quad \forall x \in [a, b]. \quad (2.30)$$

(2) If  $a = \min_{0 \leq k \leq n} x_k$  and  $b = \max_{0 \leq k \leq n} x_k$ , then

$$\int_a^b |K_n(x, t)| dt = \frac{1}{(n+1)!} \prod_{k=0}^n |x - x_k| \quad \forall x \in [a, b]. \quad (2.31)$$

*Proof.* (1) Let  $Q_{n+1}(t) = t^{n+1}/(n+1)!$ . By Theorem 2.6, we have for any  $x \in [a, b]$  that

$$Q_{n+1}(x) - (L_n Q_{n+1})(x) = \frac{1}{(n+1)!} \prod_{k=0}^n (x - x_k).$$

On the other hand, we have by Theorem 2.13 that

$$Q_{n+1}(x) - (L_n Q_{n+1})(x) = \int_a^b K_n(x, t) Q_{n+1}^{(n+1)}(t) dt = \int_a^b K_n(x, t) dt.$$

Thus, (2.30) holds true.

(2) Let  $x \in [a, b]$ . By the lemma below,  $K_n(x, \cdot)$  does not change its sign in  $(a, b)$ . Thus,

$$\left| \int_a^b K_n(x, t) dt \right| = \int_a^b |K_n(x, t)| dt.$$

This and (2.30) imply (2.31). □

**Lemma 2.16.** Assume  $a = \min_{0 \leq k \leq n} x_k$  and  $b = \max_{0 \leq k \leq n} x_k$ . Assume  $0 \leq m \leq n$  and  $x \in [a, b]$ . Then,  $K_m(x, \cdot)$  changes its sign in  $(a, b)$  exactly  $n - m$  times. In particular,  $K_n(x, \cdot)$  does not change its sign in  $(a, b)$ .

*Proof.* The statement is trivially true for the case  $n = 0$ , since  $K_0(x, \cdot) = 0$  by (2.26). Assume  $n \geq 1$ . We divide our proof into three steps.

*Step 1.* If  $0 \leq m \leq n - 1$ , then the function  $K_m(x, \cdot) \in C[a, b]$  changes its sign in  $(a, b)$  at least once. This follows from an application of (2.27) to  $f = Q_{m+1} \in \mathcal{P}_{m+1} \subseteq \mathcal{P}_n$  with  $Q_{m+1}(t) = t^{m+1}/(m+1)!$  for which  $L_n Q_{m+1} = Q_{m+1}$ :

$$0 = Q_{m+1}(x) - (L_n Q_{m+1})(x) = \int_a^b K_m(x, t) Q_{m+1}^{(m+1)}(t) dt = \int_a^b K_m(x, t) dt.$$

*Step 2.* Let  $0 \leq m \leq n - 1$ . If  $K_m(x, \cdot)$  changes sign in  $(a, b)$  exactly  $k$  times, then  $K_{m+1}(x, \cdot)$  changes its sign in  $(a, b)$  at most  $k - 1$  times. This follows from the fact that  $(d/dt)K_{m+1}(x, t) = -K_m(x, t)$  and  $K_{m+1}(x, a) = K_{m+1}(x, b) = 0$ , and an application of Rolle's Theorem.

*Step 3.* For  $0 \leq m \leq n$ ,  $K_m(x, \cdot)$  changes its sign in  $(a, b)$  exactly  $n - m$  times. In particular,  $K_n(x, \cdot)$  does not change its sign in  $(a, b)$ .

The function  $K_0(x, \cdot)$  is a piecewise constant. It has jumps at  $x_0, \dots, x_n, x$ . By the assumption of the lemma, two of these points are  $a$  and  $b$ . Therefore,  $K_0(x, \cdot)$  can change

its sign in  $(a, b)$  at most  $n$  times. By Step 2 and an argument of induction,  $K_m(x, \cdot)$  can change its sign in  $(a, b)$  at most  $n - m$  times. If for some  $m$  with  $0 \leq m \leq n - 1$ ,  $K_m(x, \cdot)$  changes its sign in  $(a, b)$  less than  $n - m$  times, then by Step 2 and induction,  $K_{n-1}(x, \cdot)$  changes its sign in  $(a, b)$  less than  $n - (n - 1) = 1$  times. By step 1, this is impossible. Thus, for each  $m$  with  $0 \leq m \leq n - 1$ ,  $K_m(x, \cdot)$  changes its sign in  $(a, b)$  exactly  $n - m$  times. By Step 2,  $K_n(x, \cdot)$  does not change its sign in  $(a, b)$ .  $\square$

## 2.4 Hermite Interpolation and Divided Differences with Repeated Points

**Theorem 2.17.** *Let  $x_1, \dots, x_n$  be  $n$  distinct points in  $[a, b]$ . Let  $y_1, \dots, y_n$  and  $y'_1, \dots, y'_n$  be  $2n$  real numbers. Then there exists a unique  $p \in \mathcal{P}_{2n-1}$  such that*

$$p(x_k) = y_k, \quad p'(x_k) = y'_k, \quad k = 1, \dots, n. \quad (2.32)$$

Moreover,  $p$  is given by

$$p(x) = \sum_{k=1}^n y_k [1 - 2l'_k(x_k)(x - x_k)][l_k(x)]^2 + \sum_{k=1}^n y'_k (x - x_k)[l_k(x)]^2, \quad (2.33)$$

where  $l_k \in \mathcal{P}_{n-1}$  ( $k = 1, \dots, n$ ) are the Lagrange basis polynomials associated with  $x_1, \dots, x_n$ .

*Proof.* Define for each integer  $k$  with  $1 \leq k \leq n$

$$\phi_k(x) = [1 - 2l'_k(x_k)(x - x_k)][l_k(x)]^2, \quad (2.34)$$

$$\psi_k(x) = (x - x_k)[l_k(x)]^2. \quad (2.35)$$

Clearly, all  $\phi_k, \psi_k$  are polynomials of degree  $2n - 1$ . Moreover,

$$\begin{aligned} \phi'_k(x) &= -2l'_k(x_k)[l_k(x)]^2 + [1 - 2l'_k(x_k)(x - x_k)]2l_k(x)l'_k(x), \\ \psi'_k(x) &= [l_k(x)]^2 + 2(x - x_k)l_k(x)l'_k(x). \end{aligned}$$

Therefore,

$$\phi_k(x_j) = \delta_{kj}, \quad \phi'_k(x_j) = 0, \quad \psi_k(x_j) = 0, \quad \psi'_k(x_j) = \delta_{kj}, \quad j, k = 1, \dots, n. \quad (2.36)$$

By the definition of  $\phi_k$  and  $\psi_k$  (cf. (2.34) and (2.35)), the polynomial  $p$  defined in (2.33) is

$$p = \sum_{k=1}^n y_k \phi_k + \sum_{k=1}^n y'_k \psi_k.$$

Clearly,  $p \in \mathcal{P}_{2n-1}$ . Moreover, (2.32) and (2.36) imply (2.32).

To prove the uniqueness, we assume  $q \in \mathcal{P}_{2n-1}$  also satisfies that  $q(x_k) = y_k$  and  $q'(x_k) = y'_k$  for  $k = 1, \dots, n$ . Then  $r := p - q \in \mathcal{P}_{2n-1}$  and  $r(x_k) = r'(x_k) = 0$  for all  $k = 1, \dots, n$ . Therefore,  $r$  has  $2n$  roots, counting the multiplicity of each root. Hence  $r = 0$  and  $q = p$ .  $\square$

We call  $p$  the *Hermite interpolation polynomial*, or *Hermite interpolant*, of  $y_k, y'_k$  at  $x_1, \dots, x_n$ . If  $y_k = f(x_k)$  and  $y'_k = f'(x_k)$  ( $k = 1, \dots, n$ ) for some  $f \in C^1[a, b]$ , then  $p$  is called the Hermite interpolation polynomial (or Hermite interpolant) of  $f$  at  $x_1, \dots, x_n$ .

**Example.** Consider  $n = 2$ ,  $x_1 = a$ , and  $x_2 = b$ . We have

$$l_1(x) = \frac{x - b}{a - b} \quad \text{and} \quad l_2(x) = \frac{x - a}{b - a}.$$

Therefore,

$$\begin{aligned} \phi_1(x) &= \left[ 1 + \frac{2(x - a)}{b - a} \right] \left( \frac{x - b}{b - a} \right)^2, \\ \phi_2(x) &= \left[ 1 - \frac{2(x - b)}{b - a} \right] \left( \frac{x - a}{b - a} \right)^2, \\ \psi_1(x) &= \frac{(x - a)(x - b)^2}{(b - a)^2}, \\ \psi_2(x) &= \frac{(x - a)^2(x - b)}{(b - a)^2}. \end{aligned}$$

The Hermite interpolation polynomial  $p_3 \in \mathcal{P}_3$  that interpolates  $y_1, y_2$  and  $y'_1, y'_2$  at  $x_1 = a, x_2 = b$  is

$$\begin{aligned} p_3(x) &= y_1\phi_1(x) + y_2\phi_2(x) + y'_1\psi_1(x) + y'_2\psi_2(x) \\ &= y_1 \left[ 1 + \frac{2(x - a)}{b - a} \right] \left( \frac{x - b}{b - a} \right)^2 + y_2 \left[ 1 - \frac{2(x - b)}{b - a} \right] \left( \frac{x - a}{b - a} \right)^2 \\ &\quad + y'_1 \frac{(x - a)(x - b)^2}{(b - a)^2} + y'_2 \frac{(x - a)^2(x - b)}{(b - a)^2}. \end{aligned}$$

**Theorem 2.18 (The remainder of Hermite interpolation).** Let  $f \in C^{2n}[a, b]$ . Let  $H_{2n-1}f \in \mathcal{P}_{2n-1}$  be the Hermite interpolation polynomial of  $f$  at  $x_1, \dots, x_n$ . Then for any  $x \in [a, b]$  there exists  $\xi = \xi(x) \in [a, b]$  such that

$$f(x) - (H_{2n-1}f)(x) = \frac{f^{(2n)}(\xi(x))}{(2n)!} (x - x_1)^2 \cdots (x - x_n)^2. \quad (2.37)$$

*Proof.* Let  $x \in [a, b]$ . If  $x = x_k$  for some  $i$  then  $\xi(x) \in [a, b]$  can be any number. So, let us assume  $x \neq x_k$  ( $k = 1, \dots, n$ ). Let

$$g(t) = f(t) - (H_{2n-1}f)(t) - \lambda(t - x_1)^2 \cdots (t - x_n)^2,$$

where  $\lambda \in \mathbb{R}$  is so chosen that  $g(x) = 0$ , i.e.,

$$\lambda = \frac{f(x) - (H_{2n-1}f)(x)}{(x - x_1)^2 \cdots (x - x_n)^2}.$$

Notice that  $g = 0$  at  $n + 1$  distinct points  $x, x_1, \dots, x_n$ . Thus it follows from Rolle's Theorem that there exist  $\xi_1, \dots, \xi_n \in [a, b]$  with  $\xi_j \neq x$  and  $\xi_j \neq x_k$  ( $k = 1, \dots, n$ ) for each  $j$  ( $1 \leq j \leq n$ ). Notice also that  $g' = 0$  at  $x_1, \dots, x_n$ . Therefore,  $g' = 0$  at  $2n$  points  $x, x_1, \dots, x_n, \xi_1, \dots, \xi_n$ . Applying Rolle's Theorem repeatedly, we conclude that  $g^{(2n)} = 0$  at some point  $\xi = \xi(x) \in [a, b]$ . This, together with the fact that

$$g^{(2n)}(t) = f^{(2n)}(t) - \lambda(2n)!$$

and the definition of  $\lambda$ , leads to (2.37). □

## 2.5 Convergence of Interpolation Polynomials

Let  $n \geq 0$  be an integer,  $x_0^{(n)}, \dots, x_n^{(n)}$  be  $n + 1$  distinct points in  $[a, b]$ , and  $L_n : C[a, b] \rightarrow \mathcal{P}_n$  the associated Lagrange interpolator. Does  $\{L_n f(x)\}_{n=0}^\infty$  converge to  $f(x)$  for any  $f \in C[a, b]$  and any  $x \in [a, b]$ ? It turns out there are many negative results.

**Runge's example.** Consider  $[a, b] = [-5, 5]$  and  $x_k^{(n)}$  evenly spaced in  $[-5, 5]$ , i.e.,  $x_k^{(n)} = -5 + 10k/n$  ( $k = 0, \dots, n; n = 1, \dots$ ). For  $f(x) = 1/(1 + x^2)$ , Runge proved that there exists  $\kappa \approx 3.63338$  such that

$$\lim_{n \rightarrow \infty} (L_n f)(x) = f(x) \quad \text{if and only if} \quad |x| < \kappa.$$

See more details in [5] (Section 3.4 of Chapter 6).

**Bernstein (1918).** For  $[a, b] = [-1, 1]$ , evenly spaced interpolation points  $x_k^{(n)} \in [-1, 1]$  ( $k = 0, \dots, n; n = 1, \dots$ ), and the function  $f(x) = |x|$ , Bernstein (19??) proved that

$$\lim_{n \rightarrow \infty} L_n f(x) = f(x) \quad \text{if and only if} \quad x \in \{0, 1, -1\}.$$

**Faber (1914).** In 1914, Faber proved the following: For any given sequence of interpolation points  $x_k^{(n)} \in [a, b]$  ( $k = 0, \dots, n; n = 0, \dots$ ), there exists  $f \in C[a, b]$  such that  $\|L_n f - f\|_{C[a, b]} \not\rightarrow 0$ .

**Bernstein (1931).** In 1931, Bernstein proved the following result: For any given sequence of interpolation points  $x_k^{(n)} \in [a, b]$  ( $k = 0, \dots, n; n = 0, \dots$ ), there exist  $f \in C[a, b]$  and  $x \in [a, b]$  such that  $(L_n f)(x) \not\rightarrow f(x)$ .

**Erdős and Vértesi (1980).** In 1980, Erdős and Vértesi proved the following striking negative result: For any given sequence of interpolation points  $x_k^{(n)} \in [a, b]$  ( $k = 0, \dots, n; n = 0, \dots$ ), there exist  $f \in C[a, b]$  such that  $(L_n f)(x) \not\rightarrow f(x)$  for almost all  $x \in [a, b]$ .

There are also some positive results.

**Theorem 2.19.** For any sequence of Lagrange interpolators  $L_n : C[a, b] \rightarrow \mathcal{P}_n$  ( $n = 0, \dots$ ),  $\|L_n f - f\|_{C[a, b]} \rightarrow 0$  for any  $f \in C[a, b]$  that is the restriction onto  $[a, b]$  of an entire function.

**Theorem 2.20.** For any  $f \in C[a, b]$ , there exist  $n + 1$  distinct points  $x_0^{(n)}, \dots, x_n^{(n)} \in [a, b]$  for each  $n \geq 0$  such that

$$\lim_{n \rightarrow \infty} \|f - L_n f\|_{C[a, b]} = 0,$$

where  $L_n : C[a, b] \rightarrow \mathcal{P}_n$  is the Lagrange interpolator associated with  $x_0^{(n)}, \dots, x_n^{(n)}$  ( $n = 0, \dots$ ).

*Proof.* Fix  $f \in C[a, b]$ . If  $f \in \mathcal{P}$ , then we can choose  $x_0^{(n)}, \dots, x_n^{(n)} \in [a, b]$  to be any  $n + 1$  distinct points for each  $n \geq 0$ . Clearly,  $L_n f = f$  for  $n$  sufficiently large.

Assume  $f \notin \mathcal{P}$ . Let  $n \geq 0$  be an integer. Let  $p_n \in \mathcal{P}_n$  be the best uniform approximation of  $f$  in  $\mathcal{P}_n$ . Then it follows from the Chebyshev Alternation Theorem that there exist  $n + 1$  distinct points  $x_k^{(n)}$  ( $k = 0, \dots, n$ ) such that

$$f(x_k^{(n)}) - p_n(x_k^{(n)}) = 0, \quad k = 0, \dots, n.$$

Therefore,  $p_n = L_n f$  is the Lagrange interpolation polynomial of  $f$  at  $x_0^{(n)}, \dots, x_n^{(n)}$ . Consequently, we have by Proposition 1.5 that

$$\|f - L_n f\|_{C[a, b]} = \|f - p_n\|_{C[a, b]} = \min_{q \in \mathcal{P}_n} \|f - q\|_{C[a, b]} \rightarrow 0 \quad \text{as } n \rightarrow \infty,$$

proving the theorem. □

**Theorem 2.21.** Let  $L_{n-1} : C[-1, 1] \rightarrow \mathcal{P}_{n-1}$  be the Lagrange interpolator associated with the  $n$  roots of the Chebyshev polynomial  $T_n$  ( $n = 1, \dots$ ). Then for any  $f \in C^2[-1, 1]$

$$\|L_n f - f\|_{C[-1, 1]} = O\left(\frac{1}{\sqrt{n}}\right) \quad \text{as } n \rightarrow \infty.$$

**Theorem 2.22 (Erdős–Turán (1937)).** Let  $x_1^{(n)}, \dots, x_n^{(n)}$  be the  $n$  distinct roots of orthogonal polynomials  $Q_n$  ( $n = 1, \dots$ ) in  $L_\rho^2(a, b)$ . For each  $n \geq 1$ , let  $L_{n-1} : C[a, b] \rightarrow \mathcal{P}_{n-1}$  be the Lagrange interpolator associated with  $x_1^{(n)}, \dots, x_n^{(n)}$ . Then

$$\lim_{n \rightarrow \infty} \int_a^b \rho(x) [f(x) - (L_{n-1} f)(x)]^2 dx = 0 \quad \forall f \in C[a, b]. \quad (2.38)$$

To prove this theorem, we need the following lemma.

**Lemma 2.23.** Let  $x_1^{(n)}, \dots, x_n^{(n)}$  be the  $n$  distinct roots of orthogonal polynomials  $Q_n$  ( $n = 1, \dots$ ) in  $L_\rho^2(a, b)$ . For each  $n \geq 1$ , let  $l_1^{(n)}, \dots, l_n^{(n)}$  be the Lagrange basis polynomials associated with  $x_1^{(n)}, \dots, x_n^{(n)}$ . Then

$$\int_a^b \rho(x) l_j^{(n)}(x) l_k^{(n)}(x) dx = 0 \quad \text{if } 1 \leq j, k \leq n, \text{ and } j \neq k, \quad (2.39)$$

$$\sum_{k=1}^n \int_a^b \rho(x) \left[ l_k^{(n)}(x) \right]^2 dx = \int_a^b \rho(x) dx. \quad (2.40)$$

*Proof.* Without loss of generality, we assume that  $n \geq 2$ . Fix  $j, k$  with  $1 \leq j, k \leq n$  and  $j \neq k$ . The polynomial  $(x - x_k^{(n)}) l_k^{(n)}(x)$  in  $\mathcal{P}_n$  has  $n$  simple roots  $x_1^{(n)}, \dots, x_n^{(n)}$ . Thus, there exists a constant  $\alpha_k^{(n)}$  such that  $(x - x_k^{(n)}) l_k^{(n)}(x) = Q_n(x)$  for all  $x$ . The polynomial  $l_j^{(n)}(x) / (x - x_k^{(n)})$  has degree  $n - 2$ . Therefore, we have by the orthogonality that

$$\begin{aligned} \int_a^b \rho(x) l_j^{(n)}(x) l_k^{(n)}(x) dx &= \int_a^b \rho(x) \left[ \frac{l_j^{(n)}(x)}{x - x_k^{(n)}} \right] (x - x_k^{(n)}) l_k^{(n)}(x) dx \\ &= \alpha_k^{(n)} \int_a^b \rho(x) \left[ \frac{l_j^{(n)}(x)}{x - x_k^{(n)}} \right] Q_n(x) dx \\ &= 0, \end{aligned}$$

proving (2.39).

By (2.6) with  $p_n(x) = 1$ , we have  $\sum_{k=1}^n l_k^{(n)}(x) = 1$  identically. Thus, it follows from (2.39) that

$$\begin{aligned} \int_a^b \rho(x) dx &= \int_a^b \rho(x) \left[ \sum_{k=1}^n l_k^{(n)}(x) \right]^2 dx \\ &= \sum_{j,k=1}^n \int_a^b \rho(x) l_j^{(n)}(x) l_k^{(n)}(x) dx \\ &= \sum_{k=1}^n \int_a^b \rho(x) \left[ l_k^{(n)}(x) \right]^2 dx. \end{aligned}$$

This is (2.40). □

*Proof of Theorem 2.22.* Let  $n \geq 2$  and let  $p_{n-1} \in \mathcal{P}_{n-1}$  be the best uniform approximation of  $f$  in  $\mathcal{P}_{n-1}$ . We have

$$\begin{aligned} &\int_a^b \rho(x) [f(x) - L_{n-1}f(x)]^2 dx \\ &\leq 2 \int_a^b \rho(x) [f(x) - p_{n-1}(x)]^2 dx + 2 \int_a^b \rho(x) [p_{n-1}(x) - L_{n-1}f(x)]^2 dx. \end{aligned} \quad (2.41)$$

By Proposition 1.5,

$$\int_a^b \rho(x) [f(x) - p_{n-1}(x)]^2 dx \leq \|f - p_{n-1}\|_{C[a,b]}^2 \int_a^b \rho(x) dx \rightarrow 0 \quad \text{as } n \rightarrow \infty. \quad (2.42)$$

It follows from Proposition 2.5, Lemma 2.23, and Proposition 1.5 that

$$\begin{aligned}
& \int_a^b \rho(x) [L_{n-1}f(x) - p_{n-1}(x)]^2 dx \\
&= \int_a^b \rho(x) \{[L_{n-1}(f - p_{n-1})](x)\}^2 dx \\
&= \int_a^b \rho(x) \sum_{j,k=1}^n \left[ f(x_j^{(n)}) - p_{n-1}(x_j^{(n)}) \right] \left[ f(x_k^{(n)}) - p_{n-1}(x_k^{(n)}) \right] l_j^{(n)}(x) l_k^{(n)}(x) dx \\
&= \sum_{j,k=1}^n \left[ f(x_j^{(n)}) - p_{n-1}(x_j^{(n)}) \right] \left[ f(x_k^{(n)}) - p_{n-1}(x_k^{(n)}) \right] \int_a^b \rho(x) l_j^{(n)}(x) l_k^{(n)}(x) dx \\
&= \sum_{k=1}^n \left[ f(x_k^{(n)}) - p_{n-1}(x_k^{(n)}) \right]^2 \int_a^b \rho(x) \left[ l_k^{(n)}(x) \right]^2 dx \\
&\leq \|f - p_{n-1}\|_{C[a,b]}^2 \sum_{k=1}^n \int_a^b \rho(x) \left[ l_k^{(n)}(x) \right]^2 dx \\
&= \|f - p_{n-1}\|_{C[a,b]}^2 \int_a^b \rho(x) dx \\
&\rightarrow 0 \quad \text{as } n \rightarrow \infty.
\end{aligned}$$

This, together with (2.41) and (2.42), implies (2.38).  $\square$

For any integer  $n \geq 1$  and any  $n$  distinct points  $x_1, \dots, x_n \in [a, b]$ , we define the *Fajér–Hermite operator*  $F_n : C[a, b] \rightarrow \mathcal{P}_{2n-1}$  by

$$F_n f = \sum_{k=1}^n f(x_k) \phi_k(x) \quad \forall f \in C[a, b],$$

where  $\phi_k$  is defined in (2.34).

**Theorem 2.24.** *Let  $F_n : C[-1, 1] \rightarrow \mathcal{P}_{2n-1}$  be the Fajér–Hermite operator associated with the zeros of Chebyshev polynomial  $T_n$ . Then*

$$\lim_{n \rightarrow \infty} \|f - F_n f\| = 0 \quad \forall f \in C[-1, 1].$$

*Proof.* By Bohman–Korovkin Theorem, we need only to show ...  $\square$

## 2.6 Piecewise Polynomial Interpolation

## 2.7 Cubic Splines

## 2.8 Trigonometric Interpolation and Fast Fourier Transforms

### Exercises

1. Find the polynomial  $p \in \mathcal{P}_3$  of the form  $p(x) = c_0 + c_1x + c_2x^2 + c_3x^3$  that interpolates a given function  $f \in C[0, 3]$  at  $x = 0, 2, 3$ .
2. Let  $x_0 = 2, x_1 = 3, x_2 = 5, x_3 = 6$  and  $y_0 = 5, y_1 = 2, y_2 = 3, y_3 = 4$ . Let  $p \in \mathcal{P}_3$  be the unique polynomial that interpolates  $y_j$  at  $x_j$  ( $j = 0, 1, 2, 3$ ). Calculate  $p$  by using:  
(a) Lagrange's formula; and (b) Newton's formula.
3. Let  $f(x) = x^4 - x^2 + 17x + 1$ . Let  $p \in \mathcal{P}_{20}$  interpolates  $f$  at  $x_j = 2^j$  ( $j = 0, \dots, 20$ ). Compute  $p(0)$ .
4. Find an approximation of  $\sqrt{3}$  with the values of the function  $f(x) = 3^x$  at  $x_0 = 0, x_1 = 1$ , and  $x_2 = 2$  using  
(a) Aitken's iterative linear interpolation method;  
(b) Neville's iterative linear interpolation method.
5. Let  $x_0, \dots, x_n$  be  $n + 1$  distinct real numbers. Let  $l_j(x)$  be the associated Lagrange basis polynomials. Show that

$$\sum_{j=0}^n (x - x_j)^k l_j(x) = 0 \quad \forall k = 1, \dots, n.$$

6. Let  $x_0, \dots, x_n$  be  $n + 1$  distinct real numbers. Let  $l_j(x)$  be the associated Lagrange basis polynomials. Show for any  $j$  with  $1 \leq j \leq n$  that

$$\sum_{i=0}^n |l_i(x)| = \left| \sum_{i=0}^{j-1} (-1)^i l_i(x) - \sum_{i=j}^n (-1)^i l_i(x) \right| \quad \forall x \in (x_{j-1}, x_j).$$

7. Recall for  $n \geq 1$  that the Chebyshev polynomial  $T_n(x)$  has  $n$  distinct roots  $x_j = \cos \theta_j$  with  $\theta_j = (2j - 1)\pi/2n$  ( $j = 1, \dots, n$ ). Denote by  $L_{n-1} : C[-1, 1] \rightarrow \mathcal{P}_{n-1}$  the associated Lagrange interpolation operator. Show that

$$(L_{n-1}f)(x) = \frac{1}{n} \sum_{j=1}^n f(x_j) \frac{(-1)^{j-1} \sin \theta_j T_n(x)}{x - x_j} \quad \forall f \in C[-1, 1].$$

8. Let  $Q_n \in \mathcal{P}_n$  ( $n = 0, 1, \dots$ ) be orthonormal polynomials in  $L^2_\rho[a, b]$ . Fix  $n \geq 2$ . Let  $x_1, \dots, x_n$  be the  $n$  distinct roots of  $Q_n(x)$  in  $(a, b)$ , and  $l_1, \dots, l_n$  be the associated Lagrange basis polynomials.

- (a) Prove that  $l_1, \dots, l_n$  are orthogonal in  $L_\rho^2[a, b]$ .  
 (b) Prove the identity

$$\sum_{j=1}^n \int_a^b \rho(x) [l_j(x)]^2 dx = \int_a^b \rho(x) dx.$$

9. Let  $x_0, \dots, x_n$  be  $n+1$  distinct points in  $[a, b]$  and  $L_n : C[a, b] \rightarrow \mathcal{P}_n$  the corresponding Lagrange interpolation operator. Show that

$$\|L_n f\|_{C[a,b]} \leq \lambda_n \|f\|_{C[a,b]} \quad \forall f \in C[a, b],$$

where

$$\lambda_n = \max_{a \leq x \leq b} \sum_{j=0}^n |l_j(x)|$$

and  $l_0, \dots, l_n$  are the Lagrange basis polynomials associated with  $x_0, \dots, x_n$ . Show also that there exists a nonzero  $\tilde{f} \in C[a, b]$  depending on  $x_0, \dots, x_n$  such that

$$\|L_n \tilde{f}\|_{C[a,b]} = \lambda_n \|\tilde{f}\|_{C[a,b]}.$$

10. Let  $\{x_j\}_{j=0}^\infty$  be a sequence of equidistant points  $x_j = x_0 + jh$  with  $h > 0$ . Define for each  $j \geq 0$

$$\Delta^0 f(x_j) = f(x_j) \quad \text{and} \quad \Delta^k f(x_j) = \Delta^{k-1} f(x_{j+1}) - \Delta^{k-1} f(x_j), \quad k = 1, \dots$$

- (a) Let  $f \in C^n[x_0, x_n]$ . Prove that

$$f[x_0, \dots, x_n] = \frac{1}{n! h^n} \Delta^n f(x_0)$$

and that

$$\Delta^n f(x_0) = h^n f^{(n)}(\xi)$$

for some  $\xi \in [x_0, x_n]$ .

- (b) Let  $f \in C^{n+1}[x_0, x_n]$ . Let  $p_n \in \mathcal{P}_n$  be the unique Lagrange polynomial that interpolates  $f$  at  $x_0, \dots, x_n$ . Let  $t$  be a real number. Show that

$$p_n(x_0 + th) = \frac{\pi_n(t)}{n!} \sum_{j=0}^n (-1)^{n-j} \binom{n}{j} \frac{f(x_j)}{t-j}$$

and that

$$p_n(x_0 + th) = f(x_0) + \frac{\pi_0(t)}{1!} \Delta f(x_0) + \frac{\pi_1(t)}{2!} \Delta^2 f(x_0) + \dots + \frac{\pi_{n-1}(t)}{n!} \Delta^n f(x_0),$$

where

$$\pi_0(t) = t, \quad \text{and} \quad \pi_j(t) = t(t-1)\cdots(t-j), \quad j = 1, \dots, n.$$

Show also that

$$f(x_0 + th) - p_n(x_0 + th) = \pi_n(t)h^{n+1} \frac{f^{(n+1)}(\eta)}{(n+1)!}$$

for some  $\eta$  in an interval containing all  $x_0, \dots, x_n$  and  $x_0 + th$ .

11. Let  $N \geq 1$  be an integer,  $h = (b-a)/N$ , and  $x_j = a + jh$ ,  $j = 0, \dots, N$ . For any  $f \in C[a, b]$ , let  $I_h f \in C[a, b]$  be such that  $I_h f \in \mathcal{P}_1$  on each  $[x_{j-1}, x_j]$  ( $j = 1, \dots, N$ ) and  $I_h f(x_j) = f(x_j)$  ( $j = 0, \dots, N$ ).

(a) Let  $f \in C^2[a, b]$ . Denote  $M_2 = \max_{a \leq x \leq b} |f''(x)|$ . Show that

$$\max_{a \leq x \leq b} |f(x) - (I_h f)(x)| \leq \frac{1}{8} M_2 h^2.$$

(b) Let  $f \in C^3[a, b]$ . Denote  $M_k = \max_{a \leq x \leq b} |f^{(k)}(x)|$  for  $k = 2$  and  $3$ . Show that

$$\max_{1 \leq j \leq N} |f'(m_j) - (I_h f)'(m_j)| \leq \frac{M_3}{24} h^2,$$

where  $m_j = (x_{j-1} + x_j)/2$  is the midpoint of the interval  $[x_{j-1}, x_j]$  ( $j = 1, \dots, N$ ), and that

$$\max_{1 \leq j \leq N} \sup_{x_{j-1} < x < x_j} |f'(x) - (I_h f)'(x)| \leq \frac{M_2}{2} h + \frac{M_3}{24} h^2.$$

12. For each integer  $n \geq 0$ , let  $x_0^{(n)}, \dots, x_n^{(n)}$  be  $n+1$  distinct points in  $[a, b]$ . Let  $L_n : C[a, b] \rightarrow \mathcal{P}_n$  be the associated Lagrange interpolation operator. Let  $f \in C^\infty[a, b]$  satisfy for some constant  $M > 0$  that

$$\|f^{(k)}\|_{C[a,b]} \leq M \quad \forall k \geq 1.$$

Show that

$$\|f - L_n f\|_{C[a,b]} \rightarrow 0 \quad \text{as } n \rightarrow \infty.$$

13. Let  $f \in C[a, b]$ . Show that, for each integer  $n \geq 1$ , there exist  $n$  distinct points  $x_1^{(n)}, \dots, x_n^{(n)}$  such that

$$\|f - L_{n-1} f\|_{C[a,b]} \rightarrow 0 \quad \text{as } n \rightarrow \infty,$$

where  $L_{n-1} f \in \mathcal{P}_{n-1}$  is the Lagrange interpolation polynomial of  $f$  at  $x_1^{(n)}, \dots, x_n^{(n)}$ .

14. Let  $n \geq 1$  be an integer. Define

$$\pi(t) = \prod_{j=0}^n (t - j) \quad \text{and} \quad l_j(t) = \frac{\pi(t)}{(t - j)\pi'_n(j)} \quad j = 0, \dots, n.$$

Show that

$$|\pi(t)| \leq n! \quad \forall t \in [0, n],$$

that

$$|l_j(t)| \leq \binom{n}{j} \quad \forall t \in [0, n], \quad j = 0, \dots, n,$$

and that

$$\sum_{j=0}^n |l_j(t)| \leq 2^n.$$

15. Let  $f \in C[a, b]$ . Let  $n \geq 1$  be a fixed integer. For each integer  $N \geq 1$ , let  $H_N = (b - a)/N$  and  $x_j^{(N)} = a + jH_N$ ,  $j = 0, \dots, N$ . Let  $p_n^{(N)} \in C[a, b]$  satisfy for each  $j$  with  $1 \leq j \leq N$  that the restriction of  $p_n^{(N)}$  on the subinterval  $[x_{j-1}^{(N)}, x_j^{(N)}]$  is the Lagrange interpolation polynomial in  $\mathcal{P}_n$  that interpolates  $f$  at the  $n + 1$  points  $x_{j-1}^{(N)} + kH_N/n$ ,  $k = 0, \dots, n$ .

(a) Show that

$$\|f - p_n^{(N)}\|_{C[a,b]} \leq 2^n \omega_f(H_N),$$

where  $\omega_f$  is the modulus of continuity of  $f$ , and that

$$\|f - p_n^{(N)}\|_{C[a,b]} \rightarrow 0 \quad \text{as } N \rightarrow \infty.$$

(b) If  $f \in C^{n+1}[a, b]$ , show that

$$\|f - p_n^{(N)}\|_{C[a,b]} \leq \frac{\|f^{(n+1)}\|_{C[a,b]}}{n + 1} \left(\frac{H_N}{n}\right)^{n+1}.$$

16. Let  $f(x) = \sin x$ ,  $[a, b] = [0, 1]$ , and  $n = 1$ . Find an integer  $N \geq 1$ , as small as possible, such that

$$\|f - p_n^{(N)}\|_{C[a,b]} \leq 1.25 \times 10^{-9},$$

where  $p_n^{(N)} \in C[0, 1]$  is the piecewise Lagrange interpolation polynomial of  $f$  defined as in the previous problem.

17. Let  $x_1, \dots, x_n$  be  $n$  distinct points and  $l_1, \dots, l_n$  the associated Lagrange basis polynomials. Prove the identity

$$\sum_{j=1}^n (x - x_j)^2 l'_j(x_j) l_j^2(x) = 0.$$

18. Let  $N \geq 1$  be an integer and  $x_j = j2\pi/N$ ,  $j = 0, \dots, N-1$ . Show for any integers  $k$  and  $l$  that

$$\sum_{j=0}^{N-1} e^{ikx_j} e^{-ilx_j} = \begin{cases} N & \text{if } k \equiv l \pmod{N}, \\ 0 & \text{if } k \not\equiv l \pmod{N}. \end{cases}$$

19. Let  $n \geq 0$  be an integer,

$$x_0 = \frac{\pi}{2(n+1)}, \quad \text{and} \quad x_j = x_0 + \frac{j\pi}{2(n+1)}, \quad j = 1, \dots, n.$$

Let  $g \in C[0, \pi]$ . Prove that there exists a unique  $T_n \in \text{span}\{1, \cos x, \dots, \cos nx\}$  such that

$$T_n(x_j) = g(x_j) \quad j = 0, \dots, n.$$

Moreover,

$$T_n(x) = \frac{\gamma_0}{2} + \sum_{k=1}^n \gamma_k \cos kx,$$

where

$$\gamma_k = \frac{2}{n+1} \sum_{j=0}^n g(x_j) \cos kx_j, \quad k = 0, \dots, n.$$

20. Let  $N \geq 1$  be an integer. Let

$$\Pi_N = \{(a_k)_{k=-\infty}^{\infty} : a_k \in \mathbb{C}, a_{k+N} = a_k, \forall k = 0, \pm 1, \dots\}$$

denote the space of all bi-infinite,  $N$ -periodic complex sequences. For any  $\mathbf{a} = (a_k)_{k=-\infty}^{\infty} \in \Pi_N$  and  $\mathbf{b} = (b_k)_{k=-\infty}^{\infty} \in \Pi_N$ , define the convolution  $\mathbf{c} = \mathbf{a} * \mathbf{b} \in \Pi_N$  by  $\mathbf{c} = (c_k)_{k=-\infty}^{\infty}$  with

$$c_k = \frac{1}{N} \sum_{j=0}^{N-1} a_j b_{k-j} \quad \forall k = 0, \pm 1, \dots$$

Prove that the discrete Fourier transform converts convolution into multiplication:

$$(\mathcal{F}_N \mathbf{c})_k = (\mathcal{F}_N \mathbf{a})_k (\mathcal{F}_N \mathbf{b})_k \quad \forall k = 0, \pm 1, \dots$$

21. Let  $n \geq 1$  be an integer and  $\Delta = \{a = x_0 < \dots < x_n = b\}$ . Suppose that  $s \in \mathcal{S}_3(\Delta)$  and  $f \in H^4(a, b)$  satisfy

$$s(x_j) = f(x_j) \quad j = 0, \dots, n.$$

Suppose also that one of the following conditions is satisfied:

- (a)  $s'(a) = f'(a)$  and  $s'(b) = f'(b)$ ;
- (b)  $s''(a) = f''(a)$  and  $s''(b) = f''(b)$ ;

(c)  $f \in H_p^4(a, b)$  and  $s \in H_p^3(a, b)$ .

Show that

$$\int_a^b [f''(x) - s''(x)]^2 dx = \int_a^b [f(x) - s(x)] f^{(4)}(x) dx.$$

22. Let  $\Delta = \{a = x_0 < \cdots < x_n = b\}$  be a partition of  $[a, b]$ . Consider the boundary condition

$$s^{(k)}(x_0) = s^{(k)}(x_n) = 0, \quad k = 0, 1, 2.$$

- (a) Show that any cubic spline on  $\Delta$  satisfying the given boundary condition vanishes identically if  $1 \leq n \leq 3$ .  
 (b) Show that any cubic spline on  $\Delta$  satisfying the given boundary condition is uniquely determined by its value at  $x_2$  if  $n = 4$ .  
 (c) Let  $n = 4$  and  $x_j = -2, -1, 0, 1, 2$ . Find explicitly the cubic spline  $s \in \mathcal{S}_3(\Delta)$  that satisfies that given boundary condition and that  $s(0) = 1$ .
23. Let  $n \geq 1$  be an integer and  $\Delta = \{a = x_0 < \cdots < x_n = b\}$  be a partition of  $[a, b]$ . Denote by  $\mathcal{S}$  the set of all cubic splines  $s$  on  $\Delta$  that satisfy  $s''(x_0) = s''(x_n) = 0$ .  
 (a) Show that for each  $j$  with  $0 \leq j \leq n$ , there exists a unique  $S_j \in \mathcal{S}$  that satisfies

$$S_j(x_k) = \delta_{jk} \quad k = 0, \dots, n.$$

- (b) Let  $f \in C[a, b]$ . Show that

$$S(x) = \sum_{j=0}^n f(x_j) S_j(x)$$

is the unique spline in  $\mathcal{S}$  that interpolates  $f$  at  $x_0, \dots, x_n$ .

- (c) What is the dimension of  $\mathcal{S}$ ?

24. Let  $x_j = a + jh$  ( $j = 0, \dots, n$ ) with  $h > 0$ . Denote by  $\mathcal{S}_3(\Delta)$  the set of splines determined by these knots  $x_j$  ( $j = 0, \dots, n$ ). Define  $S_j \in \mathcal{S}_3(\Delta)$  ( $j = 0, \dots, n$ ) by

$$S_j(x_k) = \delta_{jk} \quad j, k = 0, \dots, n \quad \text{and} \quad S_j''(x_0) = S_j''(x_n) = 0.$$

Fix  $j$  with  $0 \leq j \leq n$ . Show that the moments  $M_1, \dots, M_{n-1}$  of  $S_j$  are given by

$$M_i = \begin{cases} -\frac{1}{\rho_i} M_{i+1} & i = 1, \dots, j-2, \\ -\frac{1}{\rho_{n-i}} M_{i-1} & i = j+2, \dots, n-1, \end{cases}$$

$$\left\{ \begin{array}{l} M_j = -\frac{6(2 + 1/\rho_{j-1} + 1/\rho_{n-j-1})}{h^2(4 - 1/\rho_{j-1} - 1/\rho_{n-j-1})} \\ M_{j-1} = \frac{6h^{-2} - M_j}{\rho_{j-1}} \\ M_{j+1} = \frac{6h^{-2} - M_j}{\rho_{n-j-1}} \end{array} \right. \quad j \neq 0, 1, n-1, n,$$

where

$$\rho_1 = 4 \quad \text{and} \quad \rho_i = 4 - 1/\rho_{i-1} \quad i = 2, \dots$$

# Chapter 3

## Numerical Integration

### 3.1 The Basics

**Definition 3.1 (Numerical quadrature).** Let  $x_1, \dots, x_n$  be  $n$  distinct points in  $[a, b]$  and  $A_1, \dots, A_n \in \mathbb{R}$ . We call

$$\int_a^b f(x) dx \approx \sum_{k=1}^n A_k f(x_k) \quad (3.1)$$

a numerical quadrature with  $x_1, \dots, x_n$  quadrature points and  $A_1, \dots, A_n$  coefficients.

We say that the quadrature (3.1) is exact for an integrable function  $f : [a, b] \rightarrow \mathbb{R}$ , if

$$\int_a^b f(x) dx = \sum_{k=1}^n A_k f(x_k).$$

**Definition 3.2 (Degree of precision).** The degree of precision of a numerical quadrature (3.1) is the smallest integer  $m \geq 0$  such that the quadrature (3.1) is exact for  $f_j(x) = x^j$ ,  $j = 0, 1, \dots, m$  but is not exact for  $f_{m+1}(x) = x^{m+1}$ .

Equivalently, the degree of precision of (3.1) is  $m$  if and only if (3.1) is exact for all  $f \in \mathcal{P}_m$  but is not exact for some  $f \in \mathcal{P}_{m+1}$ .

Since the quadrature (3.1) is determined by  $2n$  parameters  $x_1, \dots, x_n$  and  $A_1, \dots, A_n$ , we expect that the degree of precision of (3.1) can not exceed  $2n - 1$ . This is indeed true.

**Proposition 3.3.** The degree of precision of any numerical quadrature (3.1) is  $\leq 2n - 1$ .

*Proof.* Let  $p_{2n}(x) = \prod_{k=1}^n (x - x_k)^2$ . Then  $p_{2n} \in \mathcal{P}_{2n}$ . Moreover,  $\int_a^b p_{2n}(x) dx > 0$  and  $\sum_{k=1}^n A_k p_{2n}(x_k) = 0$ . Thus, the degree of precision  $\leq 2n - 1$ .  $\square$

If the quadrature points of the numerical quadrature (3.1) are known, then one can use the *method of determined coefficients* to find the coefficients  $A_1, \dots, A_n$  so that the degree

of precision of the quadrature can be as high as possible.

**An example of the method of undetermined coefficients.** Find  $A_1$  and  $A_2$  such that the numerical quadrature

$$\int_0^1 f(x)dx \approx A_1f(0) + A_2f(1)$$

has the degree of precision as high as possible.

We choose  $A_1$  and  $A_2$  so that this quadrature is exact for  $f(x) = 1$  and  $f(x) = x$ :

$$\begin{aligned}\int_0^1 dx &= 1 = A_1 + A_2, \\ \int_0^1 x dx &= \frac{1}{2} = A_2.\end{aligned}$$

Solving these two equations, we obtain that  $A_1 = A_2 = 1/2$ . The quadrature thus becomes

$$\int_0^1 f(x)dx \approx \frac{1}{2}[f(0) + f(1)].$$

To find out the degree of precision of this quadrature, we check its exactness for  $f(x) = x^2$ . We have

$$\begin{aligned}\int_0^1 x^2 dx &= \frac{1}{3}, \\ \frac{1}{2}(0^2 + 1^2) &= \frac{1}{2}.\end{aligned}$$

Thus, this quadrature is not exact for  $f(x) = x^2$ . Consequently, the degree of precision of this quadrature is 1.

In the rest of this section, we give a few examples of simple numerical quadrature. For each of these examples, we determine the degree of precision, give the related composite formula, and derive its error formula.

**The left-endpoint rectangle rule.**

$$\int_a^b f(x)dx \approx f(a)(b - a). \tag{3.2}$$

To find out the degree of precision of this quadrature, we check

$$\int_a^b 1 dx = 1(b - a)$$

$$\int_a^b x dx = \frac{1}{2}(b^2 - a^2) \neq a(b - a)$$

So, the degree of precision is  $m = 0$ . Let  $f \in C^1[a, b]$ . We have

$$\begin{aligned} \int_a^b f(x) dx - f(a)(b - a) &= \int_a^b [f(x) - f(a)] dx \\ &= \int_a^b f'(\xi(x))(x - a) dx \\ &= f'(\xi) \int_a^b (x - a) dx, \end{aligned}$$

where we have used the Generalized Mean-Value Theorem<sup>1</sup> for integrals.

**Composite left-endpoint rectangle rule.** Let  $f \in C[a, b]$ . Let  $N \geq 1$  be an integer. Define  $h = (b - a)/N$  and  $x_j = a + jh$ ,  $j = 0, \dots, N$ . If we apply the left-endpoint rectangle rule to each interval  $[x_{j-1}, x_j]$  ( $1 \leq j \leq N$ ), we obtain

$$\int_a^b f(x) dx = \sum_{j=1}^N \int_{x_{j-1}}^{x_j} f(x) dx \approx \sum_{j=1}^N f(x_{j-1})(x_j - x_{j-1}) = h \sum_{j=1}^N f(x_{j-1}) = h \sum_{j=0}^{N-1} f(x_j).$$

Error:

$$\begin{aligned} f &\in C^1[a, b] \\ \int_a^b f(x) dx - h \sum_{j=0}^{N-1} f(x_j) &= \sum_{j=1}^N \left[ \int_{x_{j-1}}^{x_j} f(x) dx - f(x_{j-1})h \right] \\ &= \sum_{j=1}^N \frac{1}{2} h^2 f'(\xi_j) \\ &= \sum_{j=1}^N \frac{1}{2} h \cdot \frac{b - a}{N} \cdot f'(\xi_j) \\ &= \frac{1}{2} (b - a) h f'(\xi). \end{aligned}$$

---

<sup>1</sup>**Generalized Mean-Value Theorem for Integrals.** Let  $u \in C[a, b]$ . Let  $v : [a, b] \rightarrow \mathbb{R}$  be integrable with  $v(x) \geq 0$  for all  $x \in [a, b]$  or  $v(x) \leq 0$  for all  $x \in [a, b]$ . Then there exists  $\xi \in [a, b]$  such that

$$\int_a^b u(x)v(x) dx = u(\xi) \int_a^b v(x) dx.$$

Since

$$\min_{x \in [a,b]} f'(x) \leq \frac{1}{N} \sum_{j=1}^N f'(\xi_j) \leq \max_{x \in [a,b]} f'(x),$$

it follows from the Intermediate-Value Theorem <sup>2</sup> that

**The midpoint rectangular rule.**

$$\int_a^b f(x)dx \approx f\left(\frac{a+b}{2}\right)(b-a)$$

$$\begin{aligned} \int_a^b 1dx &= 1(b-a) \\ \int_a^b xdx &= \frac{1}{2}(b^2 - a^2) = \left(\frac{a+b}{2}\right)(b-a) = \frac{1}{2}(b^2 - a^2) \\ \int_a^b x^2 dx &= \frac{1}{3}(b^3 - a^3) \neq \left(\frac{a+b}{2}\right)^2(b-a) \end{aligned}$$

$$\begin{aligned} \frac{1}{3}(b-a)(b^2 - ba + a^2) &\neq \frac{1}{4}(a+b)^2(b-a) \\ \Leftrightarrow \frac{1}{3}(b^2 - ba + a^2) &\neq \frac{1}{4}(a^2 + 2ab + b^2) \\ \Leftrightarrow 4b^2 - 4ba + 4a^2 &\neq 3a^2 + 6ab + 3b^2 \\ \Leftrightarrow b^2 + a^2 - 2ab &\neq 0 \Leftrightarrow a \neq b \end{aligned}$$

So the degree of precision is  $m = 1$ .

Let  $f \in C^2[a, b]$

$$\begin{aligned} &\int_a^b f(x)dx - f\left(\frac{a+b}{2}\right)(b-a) \\ &= \int_a^b [f(x) - f\left(\frac{a+b}{2}\right)]dx \\ &= \int_a^b \left[f'\left(\frac{a+b}{2}\right)\left(x - \frac{a+b}{2}\right) + \frac{1}{2}f''(\xi(x))\left(x - \frac{a+b}{2}\right)^2\right]dx \\ &= \frac{1}{2} \int_a^b f''(\xi(x))\left(x - \frac{a+b}{2}\right)^2 dx \end{aligned}$$

---

<sup>2</sup>**The Intermediate-Value Theorem.** If  $f \in C[a, b]$  and  $\mu \in \mathbb{R}$  satisfy  $\min_{x \in [a,b]} f(x) \leq \mu \leq \max_{x \in [a,b]} f(x)$ , then there exist  $\xi \in [a, b]$  such that  $f(\xi) = \mu$ .

$$\begin{aligned}
&= \frac{1}{2} f''(\xi) \int_a^b \left(x - \frac{a+b}{2}\right)^2 dx \\
&= \frac{1}{24} (b-a)^3 f''(\xi).
\end{aligned}$$

### The composite mid-point rule

$$\begin{aligned}
\int_a^b f(x) dx &= \sum_{j=1}^N \int_{x_{j-1}}^{x_j} f(x) dx \\
&\approx \sum_{j=1}^N f\left(\frac{x_{j-1} + x_j}{2}\right) (x_j - x_{j-1}) \\
&= h \sum_{j=1}^N f(x_{j-\frac{1}{2}}) \quad x_{j-\frac{1}{2}} = \frac{1}{2}(x_{j-1} + x_j)
\end{aligned}$$

Error:  $f \in C^2[a, b]$

$$\begin{aligned}
&\int_a^b f(x) dx - h \sum_{j=1}^N f(x_{j-\frac{1}{2}}) \\
&= \sum_{j=1}^N \left[ \int_{x_{j-1}}^{x_j} f(x) dx - f\left(\frac{x_{j-1} + x_j}{2}\right) h \right] \\
&= \sum_{j=1}^N \frac{1}{24} h^3 f''(\xi_j) \\
&= \frac{(b-a)}{24} h^2 f''(\xi) \quad \xi \in [a, b].
\end{aligned}$$

### The trapezoidal rule.

$$\begin{aligned}
\int_a^b f(x) dx &\approx \frac{1}{2} [f(a) + f(b)] (b-a) \\
\int_a^b 1 dx &= \frac{1}{2} (1+1)(b-a) \\
\int_a^b x dx &= \frac{1}{2} (b^2 - a^2) = \frac{1}{2} [a+b](b-a) \\
\int_a^b x^2 dx &= \frac{1}{3} (b^3 - a^3) \neq \frac{1}{2} (a^2 + b^2)(b-a) \\
m &= 1
\end{aligned}$$

Let  $f \in C^2[a, b]$ .

$$\begin{aligned} & \int_a^b f(x)dx - \frac{1}{2}[f(a) + f(b)](b - a) \\ &= \int_a^b \left\{ f(x) - \frac{1}{2}[f(a) + f(b)] \right\} dx \\ &= -\frac{1}{12}(b - a)^3 f''(\xi), \text{ for some } \xi \in [a, b]. \end{aligned}$$

### Composite Rule

$$\int_a^b f(x)dx \approx \frac{h}{2} [f(x_0) + f(x_N)] + h \sum_{j=1}^{N-1} f(x_j)$$

$$\int_a^b f(x)dx - \left\{ \frac{h}{2} [f(x_0) + f(x_N)] + h \sum_{j=1}^{N-1} f(x_j) \right\} = -\frac{(b - a)h^2}{12} f''(\xi)$$

## 3.2 Interpolatory Quadrature

Let  $x_0, \dots, x_n$  be  $n + 1$  distinct points in  $[a, b]$ . Let  $f \in C[a, b]$ . The Lagrange interpolation polynomial  $L_n f \in \mathcal{P}_n$  of  $f$  at  $x_0, \dots, x_n$  is given by

$$(L_n f)(x) = \sum_{k=0}^n f(x_k) l_k(x),$$

where  $l_k(x) (k = 0, \dots, n)$  are the Lagrange basis polynomials associated with  $x_0, \dots, x_n$ .

$$l_k(x) = \prod_{\substack{i=0 \\ i \neq k}}^n \frac{x - x_i}{x_k - x_i}, \quad k = 0, \dots, n.$$

The approximation

$$\int_a^b f(x)dx \approx \int_a^b (L_n f)(x)dx = \sum_{k=0}^n \left[ \int_a^b l_k(x)dx \right] f(x_k)$$

leads to the following:

**Definition 3.4 (Interpolatory quadrature).** *The interpolatory quadrature associated with  $n + 1$  distinct points  $x_0, \dots, x_n$  in  $[a, b]$  is the numerical quadrature*

$$\int_a^b f(x)dx \approx \sum_{k=0}^n A_k f(x_k)$$

with

$$A_k = \int_a^b l_k(x) dx, \quad k = 0, \dots, n, \quad (3.3)$$

where  $l_0, \dots, l_n$  are the Lagrange basis polynomials associated with  $x_0, \dots, x_n$ .

**Theorem 3.5 (Characterization of interpolatory quadrature).** *Let  $x_0, \dots, x_n$  be  $n+1$  distinct points in  $[a, b]$ . A numerical quadrature*

$$\int_a^b f(x) dx \approx \sum_{k=0}^n B_k f(x_k) \quad (3.4)$$

*is an interpolatory quadrature if and only if its degree of precision is  $\geq n$ .*

*Proof. The “if” part.* Suppose the degree of precision of (3.4) is  $\geq n$ . Let  $l_0, \dots, l_n \in \mathcal{P}_n$  be the Lagrange basis polynomials associated with  $x_0, \dots, x_n$ . Since each  $l_j \in \mathcal{P}_n$ , we have

$$\int_a^b l_j(x) dx = \sum_{k=0}^n B_k l_j(x_k) = \sum_{k=0}^n B_k \delta_{jk} = B_j.$$

Thus, the quadrature (3.4) is interpolatory.

*The “only if” part.* Suppose (3.4) is interpolatory. Then the coefficients are given by

$$B_k = \int_a^b l_k(x) dx, \quad k = 0, \dots, n.$$

Let  $f \in \mathcal{P}_n$ . Then  $L_n f = f$  by Proposition 2.5. Consequently,

$$\int_a^b f(x) dx = \int_a^b (L_n f)(x) dx = \sum_{k=0}^n \int_a^b l_k(x) dx f(x_k) = \sum_{k=0}^n B_k f(x_k).$$

This implies that the degree of precision of (3.4) is  $\geq n$ . □

**Definition 3.6 (Newton-Cotes formula).** *Let  $n \geq 1$  be an integer. A (closed) Newton-Cotes formula is an interpolatory quadrature*

$$\int_a^b f(x) dx \approx \sum_{k=0}^n A_k f(x_k) \quad (3.5)$$

*with the quadrature points  $x_k = a + k(b - a)/n$  ( $k = 0, \dots, n$ ).*

By the definition of interpolatory quadrature, the coefficients  $A_k$  ( $k = 0, \dots, n$ ) in the Newton-Cotes formula (3.5) are given by (3.3) with  $l_k \in \mathcal{P}_n$  ( $k = 0, \dots, n$ ) the Lagrange basis polynomials associated with the evenly distributed quadrature points  $x_k$  ( $k = 0, \dots, n$ ).

**Examples.** (1) Consider the Newton-Cotes formula with  $n = 1$ . We have  $x_0 = a$ ,  $x_1 = b$ , and

$$A_0 = \int_a^b l_0(x) dx = \int_a^b \frac{x-b}{a-b} dx = \frac{1}{2}(b-a),$$

$$A_1 = \int_a^b l_1(x) dx = \int_a^b \frac{x-a}{b-a} dx = \frac{1}{2}(b-a).$$

Thus, the formula is

$$\int_a^b f(x) dx \approx \frac{1}{2}(b-a)f(x_0) + \frac{1}{2}(b-a)f(x_1) = \frac{1}{2}(b-a)[f(x_0) + f(x_1)].$$

This is exactly the trapezoidal rule.

(2) Consider the Newton-Cotes formula with  $n = 2$ . We have  $x_0 = a$ ,  $x_1 = (a+b)/2 =: c$ ,  $x_2 = b$ , and

$$A_0 = \int_a^b l_0(x) dx = \int_a^b \frac{(x-c)(x-b)}{(a-c)(a-b)} dx = \frac{b-a}{6},$$

$$A_1 = \int_a^b l_1(x) dx = \int_a^b \frac{(x-a)(x-b)}{(c-a)(c-b)} dx = \frac{2(b-a)}{3},$$

$$A_2 = \int_a^b l_2(x) dx = \int_a^b \frac{(x-a)(x-c)}{(b-a)(b-c)} dx = \frac{b-a}{6}.$$

The formula is

$$\int_a^b f(x) dx \approx \frac{b-a}{6} \left[ f(a) + 4f\left(\frac{a+b}{2}\right) + f(b) \right].$$

This is called *Simpson's rule*.

In the case  $[a, b] = [-1, 1]$ , this becomes

$$\int_{-1}^1 f(x) dx \approx \frac{1}{3}[f(-1) + 4f(0) + f(1)].$$

We can verify directly that this is exact for  $f(x) = 1$ ,  $x$ ,  $x^2$ . In fact, it is also exact for  $f(x) = x^3$  but not for  $f(x) = x^4$ . Therefore, the degree of precision of Simpson's rule is 3.

**Theorem 3.7 (Error formula for Newton-Cotes formula).** Consider a Newton-Cotes formula

$$\int_a^b f(x) dx \approx \sum_{k=0}^n A_k f(x_k). \quad (3.6)$$

(1) If  $n$  is even and  $f \in C^{n+2}[a, b]$ , then exists  $\xi \in (a, b)$  such that

$$\int_a^b f(x)dx - \sum_{k=0}^n A_k f(x_k) = \frac{f^{(n+2)}(\xi)}{(n+2)!} \mu_n, \quad (3.7)$$

where

$$\mu_n = \int_a^b x(x-x_0) \cdots (x-x_n) dx < 0.$$

(2) If  $n$  is odd and  $f \in C^{n+1}[a, b]$ , then there exists  $\eta \in (a, b)$  such that

$$\int_a^b f(x)dx - \sum_{k=0}^n A_k f(x_k) = \frac{f^{(n+1)}(\eta)}{(n+1)!} \nu_n, \quad (3.8)$$

where

$$\nu_n = \int_a^b (x-x_0) \cdots (x-x_n) dx < 0.$$

**Colloary 3.8.** The degree of precision of the Newton-Cotes formula (3.6) with quadrature points  $x_k = a + k(b-a)/n$  ( $k = 0, \dots, n$ ) is  $n$  if  $n$  is odd and  $n+1$  if  $n$  is even.

*Proof.* Suppose  $n \geq 1$  is even. By (3.7), the quadrature (3.6) is exact for all  $f \in \mathcal{P}_{n+1}$ . Setting  $f(x) = x^{n+2}$  in (3.7), we see that the right-hand side of (3.7) is  $\mu_n \neq 0$ . Thus, the degree of precision in this case is  $n+1$ . The same argument applies to the case  $n$  is odd.  $\square$

To prove Theorem 3.7, we first prove the following:

**Lemma 3.9.** Let  $n \geq 1$  be an even number,  $h = (b-a)/n$ , and  $x_k = a + kh$  ( $k = 0, \dots, n$ ). Let

$$\omega_n(x) = (x-x_0) \cdots (x-x_n) \quad \text{and} \quad \Omega_n(x) = \int_a^x \omega_n(t) dt. \quad (3.9)$$

Then  $\Omega_n(a) = \Omega_n(b) = 0$  and  $\Omega_n(x) > 0$  for all  $x \in (a, b)$ .

*Proof.* It is obvious that  $\Omega_n(a) = \int_a^a \omega_n(t) dt = 0$ . Since  $n$  is even, we have by the change of variable  $x = a + h(t + n/2)$  that

$$\Omega_n(b) = \int_a^b \omega(x) dx = h^{n+2} \int_{-\frac{n}{2}}^{\frac{n}{2}} t \prod_{k=1}^{\frac{n}{2}} (t^2 - k^2) dt = 0.$$

Let an index  $j$  be such that  $1 \leq j \leq n/2$ . We claim:

$$|\omega_n(x)| > |\omega_n(x+h)| \quad \forall x \in (x_{2j-2}, x_{2j-1}). \quad (3.10)$$

To see this, let us fix an  $x \in (x_{2j-2}, x_{2j-1})$ . Let  $x = a + ht$  for some  $t \in \mathbb{R}$ . Clearly,  $2j - 2 < t < 2j - 1$ . Thus,  $t$  is not an integer. Moreover, since  $1 \leq j \leq n/2$ , we have  $0 < t + 1 < n/2$ . From  $x = a + ht$  and  $x_k = a + kh$  ( $k = 0, \dots, n$ ), we then obtain that

$$\left| \frac{\omega_n(x+h)}{\omega_n(x)} \right| = \left| \frac{(t+1)t \cdot (t-1) \dots (t-n+1)}{t(t-1) \dots (t-n)} \right| = \left| \frac{t+1}{t-n} \right| = \frac{t+1}{n-t} < 1,$$

where the last inequality is equivalent to the true fact that  $t + 1/2 < n/2$ . This proves (3.10).

Consider now  $[x_0, x_2] = [x_0, x_1] \cup [x_1, x_2]$ . Since  $n$  is even,  $\omega_n(x) < 0$  on  $(-\infty, x_0)$  and  $\omega_n(x) > 0$  on  $(x_0, x_1)$ . Thus, by the fact that  $\Omega_n(x_0) = 0$ , we have  $\Omega_n(x) > 0$  on  $(x_0, x_1]$ . Let  $x \in (x_1, x_2]$ . Then  $x - h \in (x_0, x_1]$ . Hence,  $\omega_n(t) > 0$  for any  $t \in (x_0, x - h)$ . This and (3.10) with  $j = 1$  imply that  $\omega_n(t) + \omega_n(t+h) > 0$  for any  $t \in (x_0, x - h)$ . Therefore, by the change of variable  $s = t - h$  and (3.10) for  $j = 1$ ,

$$\begin{aligned} \Omega_n(x) &= \int_a^x \omega_n(t) dt = \int_{x_0}^{x_1} \omega_n(t) dt + \int_{x_1}^x \omega_n(t) dt \\ &\geq \int_{x_0}^{x-h} \omega_n(t) dt + \int_{x_0}^{x-h} \omega_n(s+h) ds = \int_{x_0}^{x-h} [\omega_n(t) + \omega_n(t+h)] dt > 0. \end{aligned}$$

Hence,  $\Omega_n(x) > 0$  on  $[x_0, x_2]$ . Since  $\Omega'_n(x) = \omega_n(x) > 0$  in  $(x_2, x_3)$  and  $\Omega_n(x_2) > 0$ , we have  $\Omega_n(x) > 0$  on  $[x_2, x_3]$ . A similar argument then leads to  $\Omega_n(x) > 0$  on  $[x_2, x_4]$ . Continuing this process, we have  $\Omega_n(x) > 0$  for  $x \in (x_0, x_{2j-1}]$  with  $2j - 1 = n/2$  or  $2j - 1 = (n/2) - 1$ . In the latter case, we have  $\Omega_n(x) > 0$  in  $(x_{2j-1}, x_{n/2})$ , since  $\Omega'_n(x) = \omega_n(x) > 0$  in  $(x_{2j-1}, x_{n/2})$  and  $\Omega_n(x) > 0$  on  $(x_0, x_{2j-1}]$ . Therefore,  $\Omega_n(x) > 0$  on  $(x_0, x_{n/2}]$ .

If  $x \in (x_{n/2}, x_n)$  then  $x_{n/2} - (x - x_{n/2}) \in (x_0, x_{n/2})$  and hence  $\Omega_n(x_{n/2} - (x - x_{n/2})) > 0$ . Moreover, by the change of variable  $s = t - x_{n/2}$ , we have

$$\int_{x_{n/2} - (x - x_{n/2})}^{x_{n/2} + (x - x_{n/2})} \omega_n(t) dt = \int_{-(x - x_{n/2})}^{x - x_{n/2}} \omega_n(s + x_{n/2}) ds = \int_{-(x - x_{n/2})}^{x - x_{n/2}} s \prod_{k=1}^{n/2} [s^2 - (kh)^2] ds = 0.$$

Therefore,

$$\begin{aligned} \Omega_n(x) &= \int_{x_0}^x \omega_n(t) dt \\ &= \int_{x_0}^{x_{n/2} - (x - x_{n/2})} \omega_n(t) dt + \int_{x_{n/2} - (x - x_{n/2})}^{x_{n/2} + (x - x_{n/2})} \omega_n(t) dt \\ &= \Omega_n((x_{n/2} - (x - x_{n/2}))) > 0. \end{aligned}$$

The proof is complete. □

*Proof of Theorem 3.7.* Let  $\omega_n$  and  $\Omega_n$  be given as in (3.9). Since (3.6) is an interpolatory formula, we have by Theorem 2.10 and Proposition ? (on the properties of divided differences with repeated points) that

$$\begin{aligned} e_n(f) &:= \int_a^b f(x)dx - \sum_{k=0}^n A_k f(x_k) \\ &= \int_a^b f(x)dx - \int_a^b (L_n f)(x)dx \\ &= \int_a^b f[x_0, \dots, x_n, x] \omega_n(x) dx \quad \forall f \in C^1[a, b]. \end{aligned} \quad (3.11)$$

*Case 1:  $n$  is even and  $f \in C^{n+2}[a, b]$ .* By integration by parts, we obtain from (3.11), Lemma 3.9, Proposition (?) (on properties on divided differences with repeated points), and the Generalized Mean-Value Theorem for integrals that

$$\begin{aligned} e_n(f) &= \int_a^b f[x_0, \dots, x_n, x] \Omega'_n(x) dx \\ &= f[x_0, \dots, x_n, x] \Omega_n(x) \Big|_{x=a}^{x=b} - \int_a^b \Omega(x) \frac{d}{dx} f[x_0, \dots, x_n, x] dx \\ &= - \int_a^b \Omega_n(x) \frac{d}{dx} f[x_0, \dots, x_n, x] dx \\ &= - \int_a^b \Omega_n(x) \frac{f^{(n+2)}(\xi(x)) dx}{(n+2)!} \\ &= - \frac{f^{(n+2)}(\xi)}{(n+2)!} \int_a^b \Omega_n(x) dx \end{aligned}$$

for some  $\xi \in (a, b)$ . By integration by parts and Lemma 3.9, we then have

$$\int_a^b \Omega_n(x) dx = x \Omega_n(x) \Big|_a^b - \int_a^b x \Omega'_n(x) dx = - \int_a^b x \omega_n(x) dx = -\mu_n.$$

Since  $\int_a^b \Omega_n(x) dx > 0$  by Lemma 3.9,  $\mu_n < 0$ .

*Case 2:  $n$  is odd and  $f \in C^{n+1}[a, b]$ .* It follows from (3.11) that

$$e_n(f) = \int_a^{b-h} \omega_n(x) f[x_0, \dots, x_n, x] dx + \int_{b-h}^b \omega_n(x) f[x_0, \dots, x_n, x] dx =: I_1 + I_2. \quad (3.12)$$

Since  $\omega_n(x)$  does not change sign in  $(b-h, b)$ , we have by the Generalized Mean-Value Theorem for integrals that

$$I_2 := \int_{b-h}^b \omega_n(x) f[x_0, \dots, x_n, x] dx = \frac{f^{(n+1)}(\eta')}{(n+1)!} \int_{b-h}^b \omega_n(x) dx \quad (3.13)$$

for some  $\eta' \in (a, b)$ .

By the definition of divided differences, we have

$$\begin{aligned}
I_1 &:= \int_a^{b-h} \omega_n(x) f[x_0, \dots, x_n, x] dx \\
&= \int_a^{b-h} \omega_{n-1}(x)(x - x_n) \left( \frac{f[x_0, \dots, x_{n-1}, x] - f[x_0, \dots, x_{n-1}, x_n]}{x - x_n} \right) dx \\
&= \int_a^{b-h} \Omega'_{n-1}(x) f[x_0, \dots, x_{n-1}, x] dx - \int_a^{b-h} \Omega'_{n-1}(x) f[x_0, \dots, x_{n-1}, x_n] dx \\
&=: J_1 - J_2.
\end{aligned}$$

Since  $n - 1$  is even, by Lemma 3.9 we have  $\Omega_{n-1}(a) = \Omega_{n-1}(b - h) = 0$ . Consequently,

$$J_2 = f[x_0, \dots, x_{n-1}, x_n] \int_a^{b-h} \Omega'_{n-1}(x) dx = f[x_0, \dots, x_{n-1}, x_n] [\Omega_{n-1}(b - h) - \Omega_{n-1}(a)] = 0. \quad (3.14)$$

Again by Lemm 3.9,  $\Omega_{n-1}(x) > 0$  on  $(a, b - h)$ . By integration by parts, Propertion? (on properties of divided differences with repeated points), and the Generalized Mean-Value Theorem for integrals, we get

$$\begin{aligned}
J_1 &:= \int_a^{b-h} \Omega'_{n-1}(x) f[x_0, \dots, x_{n-1}, x] dx \\
&= - \int_a^{b-h} \Omega_{n-1}(x) \frac{d}{dx} f[x_0, \dots, x_{n-1}, x] dx \\
&= - \frac{f^{(n+1)}(\eta'')}{(n+1)!} \int_a^{b-h} \Omega_{n-1}(x) dx \quad (3.15)
\end{aligned}$$

for some  $\eta'' \in (a, b)$ .

From (3.12)–(3.15), we obtain

$$e_n(f) = -[A f^{(n+1)}(\eta') + B f^{(n+1)}(\eta'')],$$

where

$$\begin{aligned}
A &= -\frac{1}{(n+1)!} \int_{b-h}^b \omega_n(x) dx, \\
B &= \frac{1}{(n+1)!} \int_a^{b-h} \Omega_{n-1}(x) dx.
\end{aligned}$$

Clearly,  $\omega_n(x) > 0$  for any  $x > b$ . Thus,  $\omega_n(x) < 0$  in  $(b - h, b)$ . This implies that  $A > 0$ . Since  $n - 1$  is even, we have  $B > 0$  by Lemma 3.9. The fact that

$$\min_{a \leq x \leq b} f^{(n+1)}(x) \leq \frac{A f^{(n+1)}(\eta') + B f^{(n+1)}(\eta'')}{A + B} \leq \max_{a \leq x \leq b} f^{(n+1)}(x)$$

and the Intermediate-Value Theorem now imply that

$$e_n(f) = -(A + B)f^{(n+1)}(\eta) \quad (3.16)$$

for some  $\eta \in (a, b)$ . Again since  $n - 1$  is even, we obtain by Lemma 3.9 that

$$\begin{aligned} \int_a^{b-h} \omega_n(x) dx &= \int_a^{b-h} \Omega'_{n-1}(x)(x-b) dx \\ &= \Omega_{n-1}(x)(x-b) \Big|_a^{b-h} - \int_a^{b-h} \Omega_{n-1}(x) dx \\ &= - \int_a^{b-h} \Omega_{n-1}(x) dx. \end{aligned}$$

Consequently,

$$\begin{aligned} A + B &= \frac{1}{(n+1)!} \int_{b-h}^b \omega_n(x) dx + \frac{1}{(n+1)!} \int_a^{b-h} \Omega_{n-1}(x) dx \\ &= \frac{1}{(n+1)!} \int_{b-h}^b \omega_n(x) dx - \frac{1}{(n+1)!} \int_a^{b-h} \omega_n(x) dx \\ &= - \frac{1}{(n+1)!} \int_a^b \omega_n(x) dx. \end{aligned} \quad (3.17)$$

This, together (3.16), leads to (3.8). Since  $A$  and  $B$  are positive, then we have by (3.17) that

$$\nu_n = \int_a^b \omega_n(x) dx = -(n+1)!(A+B) < 0,$$

completing the proof.  $\square$

### 3.3 Peano Kernel and Error Representation

**Theorem 3.10 (Peano kernel and error representation for numerical quadrature).**

*Assume the degree of precision of a given numerical quadrature*

$$\int_a^b f(x) dx \approx \sum_{k=0}^n A_k f(x_k)$$

*is  $m$ . Then*

$$\int_a^b f(x) - \sum_{k=0}^n A_k f(x_k) = \int_a^b \tilde{K}_m(t) f^{(m+1)}(t) dt \quad \forall f \in C^{(m+1)}[a, b], \quad (3.18)$$

*where*

$$\tilde{K}_m(t) = \frac{1}{m!} \left[ \int_a^b (x-t)_+^m dx - \sum_{k=0}^n A_k (x_k - t)_+^m \right]. \quad (3.19)$$

*Proof.* Let  $f \in C^{(m+1)}[a, b]$ . By the Taylor expansion,

$$\begin{aligned} f(x) &= f(a) + f'(a)(x-a) + \cdots + \frac{f^{(m)}(a)}{m!}(x-a)^m + \frac{1}{m!} \int_a^b (x-t)_+^m f^{(m+1)}(t) dt \\ &= Q_m(x) + R_m(x). \end{aligned}$$

Since the degree of precision of (3.18) is  $m$  and

$$Q_m(x) = f(a) + f'(a)(x-a) + \cdots + \frac{f^{(m)}(a)}{m!}(x-a)^m$$

is a polynomial of degree  $\leq m$ , the quadrature (3.18) is exact for  $Q_m$ . Thus,

$$\begin{aligned} &\int_a^b f(x) dx - \sum_{k=0}^n A_k f(x_k) \\ &= \int_a^b Q_m(x) dx - \sum_{k=0}^n A_k Q_m(x_k) + \int_a^b R_m(x) dx - \sum_{k=0}^n A_k R_m(x_k) \\ &= \int_a^b R_m(x) dx - \sum_{k=0}^n A_k R_m(x_k) \\ &= \frac{1}{m!} \int_a^b \left[ \int_a^b (x-t)_+^m dx - \sum_{k=0}^n A_k (x_k - t)_+^m \right] f^{(m+1)}(t) dt \\ &= \int_a^b \tilde{K}_m(t) f^{(m+1)}(t) dt. \end{aligned}$$

The proof is complete. □

We call  $\tilde{K}_m$  defined in (3.19) the Peano kernel for the numerical quadrature (3.18).

## 3.4 Euler–Maclaurin Formula, Richardson Extrapolation, and Romberg Algorithm

## 3.5 Weighted Gaussian Quadrature

Let  $\rho$  be a weight function on  $[a, b]$ . We consider a numerical quadrature

$$\int_a^b \rho(x) f(x) dx \approx \sum_{k=1}^n A_k f(x_k), \quad (3.20)$$

where  $n \geq 1$  is an integer,  $x_1, \dots, x_n \in [a, b]$  are  $n$  distinct points, and  $A_1, \dots, A_n \in \mathbb{R}$ . It follows from Proposition 3.3 that the degree of precision of any numerical quadrature (3.20)

is less than or equal to  $2n - 1$ . On the other hand, by ?, the degree of precision of (3.20) is greater than or equal to  $n - 1$ , if it is interpolatory, i.e., the coefficients are given by

$$A_k = \int_a^b \rho(x) l_k(x) dx, \quad k = 1, \dots, n, \quad (3.21)$$

where

$$l_k(x) = \prod_{\substack{j=1 \\ j \neq k}}^n \frac{x - x_j}{x_k - x_j}, \quad k = 1, \dots, n,$$

are the Lagrange basis polynomials associated with  $x_1, \dots, x_n$ .

We want to choose the quadrature points  $x_1, \dots, x_n$  so that the formula (3.20) has the degree of precision as high as possible. Let  $f \in C[a, b]$ . Let  $L_{n-1} : C[a, b] \rightarrow \mathcal{P}_{n-1}$  be the Lagrange interpolator associated with  $x_1, \dots, x_n$ . By Theorem 2.10, we have the error

$$\begin{aligned} e_n(f) &:= \int_a^b \rho(x) f(x) dx - \sum_{k=1}^n A_k f(x_k) \\ &= \int_a^b \rho(x) [f(x) - (L_{n-1}f)(x)] dx \\ &= \int_a^b \rho(x) f[x_1, \dots, x_n, x] \prod_{k=1}^n (x - x_k) dx. \end{aligned}$$

If  $f \in \mathcal{P}_m$  for some integer  $m \geq n$ , then by Proposition 2.11 the divided difference  $f[x_1, \dots, x_n, x]$  is a polynomial in  $\mathcal{P}_{m-n}$ . We thus want to choose  $x_1, \dots, x_n$  so that

$$\int_a^b \rho(x) p(x) \prod_{k=1}^n (x - x_k) dx = 0 \quad \forall p \in \mathcal{P}_{m-n} \quad (3.22)$$

with possibly  $m = n, \dots, 2n - 1$ . This will not hold true for  $m = 2n$  by Proposition 3.3. It is then clear that if  $\omega_n(x) := \prod_{k=1}^n (x - x_k)$  is the  $n$ th orthogonal polynomial in  $\tilde{\mathcal{P}}_n$ , i.e.,  $x_1, \dots, x_n$  are roots of an  $n$ th orthogonal polynomial, then (3.22) will hold true for all  $m = n, \dots, 2n - 1$ . This means that  $e_n(f) = 0$  for any  $f \in \mathcal{P}_m$  with  $m = n, \dots, 2n - 1$ . Clearly,  $e_n(f) = 0$  for any  $f \in \mathcal{P}_m$  with  $m = 0, \dots, n - 1$ , if (3.20) is interpolatory. Therefore, the highest possible degree of precision is achieved by an interpolatory quadrature with quadrature points roots of orthogonal polynomials.

**Definition 3.11 (Weighted Gaussian quadrature).** *A numerical quadrature (3.20) is called a weighted Gaussian quadrature, if*

- (1) *the quadrature points  $x_1, \dots, x_n$  are the  $n$  simple roots of an orthogonal polynomial of degree  $n$  in  $L_\rho^2(a, b)$ ,*
- (2) *the quadrature is interpolatory, i.e., the coefficients  $A_1, \dots, A_n$  are given by (3.21).*

**Theorem 3.12 (Characterization of weighted Gaussian quadrature).** *A numerical quadrature (3.20) is a weighted Gaussian quadrature if and only if it has the degree of precision  $2n - 1$ .*

*Proof. The “if” part.* Assume the degree of precision of the numerical quadrature (3.20) is  $2n - 1$ . Let  $Q_n(x) = \prod_{k=1}^n (x - x_k)$  and  $q \in \mathcal{P}_{n-1}$ . Then  $qQ_n \in \mathcal{P}_{2n-1}$ . Since the degree of precision of (3.20) is  $2n - 1$ ,

$$\int_a^b \rho(x)q(x)Q_n(x)dx = \sum_{k=1}^n A_k q(x_k)Q_n(x_k) = 0.$$

Therefore,  $Q_n \in \mathcal{P}_n$  is an  $n$ th orthogonal polynomial in  $L^2_\rho(a, b)$ , and hence  $x_1, \dots, x_n$  are roots of this polynomial. Moreover, since the quadrature is exact for all polynomials in  $\mathcal{P}_{n-1}$ , it is interpolatory by ?. Thus, it is a weighted Gaussian quadrature by Definition 3.11.

*The “only if” part.* Assume (3.20) is a weighted Gaussian quadrature. For  $p \in \mathcal{P}_{2n-1}$ , there exist  $q \in \mathcal{P}_{n-1}$  and  $r \in \mathcal{P}_{n-1}$  with  $\deg r < \deg p$  such that

$$p(x) = q(x)\tilde{Q}_n(x) + r(x),$$

where  $\tilde{Q}_n(x) = \prod_{k=1}^n (x - x_k)$  in  $\tilde{P}_n$  is the  $n$ th orthogonal polynomial in  $L^2_\rho(a, b)$ . Clearly,  $p(x_k) = r(x_k)$  for  $k = 1, \dots, n$ . Thus, by the orthogonality and the fact that the weighted Gaussian quadrature (3.20) is exact for any polynomial in  $\mathcal{P}_{n-1}$ , we have

$$\int_a^b \rho(x)p(x) dx = \int_a^b \rho(x)q(x)\tilde{Q}_n(x) dx + \int_a^b \rho(x)r(x) dx = \sum_{k=1}^n A_k r(x_k) = \sum_{k=1}^n A_k p(x_k).$$

Therefore, the degree of precision of (3.20) is greater than or equal to  $2n - 1$ ; and is in fact exactly  $2n - 1$  by Proposition 3.3.  $\square$

**Theorem 3.13 (Error of weighted Gaussian quadrature).** *Let (3.20) be a weighted Gaussian quadrature. For any  $f \in C^{2n}[a, b]$ , there exists  $\xi \in [a, b]$  such that*

$$\int_a^b \rho(x)f(x)dx - \sum_{k=1}^n A_k f(x_k) = \frac{f^{(2n)}(\xi)}{(2n)!} \int_a^b \rho(x) \prod_{k=1}^n (x - x_k)^2 dx.$$

*Proof.* Let  $p \in \mathcal{P}_{2n-1}$  be the Hermite interpolation polynomial of  $f$  at  $x_1, \dots, x_n$ . Then it follows from Theorem 2.18 that for each  $x \in [a, b]$  there exists  $\xi(x) \in [a, b]$  such that

$$f(x) - p(x) = \frac{f^{(2n)}(\xi(x))}{(2n)!} \prod_{k=1}^n (x - x_k)^2.$$

Since the weighted Gaussian quadrature (3.20) has the degree of precision  $2n - 1$  and since  $p(x_k) = f(x_k)$  ( $k = 1, \dots, n$ ), we have by the Generalized Mean-value Theorem for integrals ? that

$$\int_a^b \rho(x)f(x)dx = \int_a^b \rho(x)p(x)dx + \int_a^b \rho(x) \frac{f^{(2n)}(\xi(x))}{(2n)!} \prod_{k=1}^n (x - x_k)^2 dx$$

$$\begin{aligned}
&= \sum_{k=1}^n A_k p(x_k) + \frac{f^{(2n)}(\xi)}{(2n)!} \int_a^b \rho(x) \prod_{k=1}^n (x - x_k)^2 dx \\
&= \sum_{k=1}^n A_k f(x_k) + \frac{f^{(2n)}(\xi)}{(2n)!} \int_a^b \rho(x) \prod_{k=1}^n (x - x_k)^2 dx,
\end{aligned}$$

completing the proof.  $\square$

The following is a useful property of a weighted Gaussian quadrature:

**Proposition 3.14.** *The coefficients of a weighted Gaussian quadrature are all positive.*

*Proof.* Let (3.20) be a weighted Gaussian quadrature. Let  $l_j \in \mathcal{P}_{n-1}$  ( $j = 1, \dots, n$ ) be the Lagrange basis polynomials associated with the quadrature points  $x_1, \dots, x_n \in (a, b)$ . Since each  $l_j^2 \in \mathcal{P}_{2n-2}$  and the weighted Gaussian quadrature (3.20) has the degree of precision  $2n - 1$ , we have

$$0 < \int_a^b \rho(x) [l_j(x)]^2 dx = \sum_{k=1}^n A_k [l_j(x_k)]^2 = A_j,$$

completing the proof.  $\square$

### Gaussian quadrature

$$\int_{-1}^1 f(x) dx \approx \sum_{k=1}^n A_k f(x_k). \quad (3.23)$$

The Legendre polynomials  $P_n$  ( $n = 0, \dots$ ) are orthogonal polynomials in  $L^2[-1, 1]$ . Recall:

$$\begin{aligned}
P_n(x) &= \frac{1}{2^n n!} \frac{d^n}{dx^n} [(x^2 - 1)^n] \\
\int_{-1}^1 P_m(x) P_n(x) dx &= \begin{cases} 0 & m \neq n \\ \frac{2}{2n+1} & m = n. \end{cases}
\end{aligned}$$

Each  $P_n$  ( $n \geq 1$ ) has  $n$  roots in  $(-1, 1)$ .

$$\begin{aligned}
P_0(x) &= 1 \\
P_1(x) &= x \\
P_2(x) &= \frac{1}{2}(3x^2 - 1) \\
P_3(x) &= \frac{1}{2}(5x^3 - 3x) \\
\int_{-1}^1 f(x) dx &\approx 2f(0) \\
\int_{-1}^1 f(x) dx &\approx f\left(-\frac{1}{\sqrt{3}}\right) + f\left(\frac{1}{\sqrt{3}}\right)
\end{aligned}$$

$$\int_{-1}^1 f(x)dx \approx \frac{5}{9}f\left(-\sqrt{\frac{3}{5}}\right) + \frac{8}{9}f(0) + \frac{5}{9}f\left(\sqrt{\frac{3}{5}}\right)$$

**Theorem 3.15.** *The coefficients of the Gaussian quadrature (3.23) are given by*

$$A_k = \frac{2}{(1-x_k^2)[P'_n(x_k)]^2}, \quad k = 1, \dots, n.$$

*Proof.* The Lagrange basis polynomials  $l_k \in \mathcal{P}_{n-1}$  ( $k = 1, \dots, n$ ) associated with the roots  $x_1, \dots, x_n$  of the  $n$ th Legendre polynomial  $P_n$  are given by

$$l_k(x) = \frac{P_n(x)}{(x-x_k)P'_n(x_k)} = \prod_{\substack{j=1 \\ j \neq k}}^n \frac{x-x_j}{x_k-x_j}, \quad k = 1, \dots, n. \quad (3.24)$$

Fix an integer  $k$  with  $1 \leq k \leq n$ . By integration by parts and the orthogonality, we obtain that

$$S_k := \int_{-1}^1 l_k(x)P'_n(x)dx = l_k(x)P_n(x) \Big|_{-1}^1 - \int_{-1}^1 l'_k(x)P_n(x)dx = l_k(x)P_n(x) \Big|_{-1}^1.$$

Since  $l_k P'_n \in \mathcal{P}_{2n-2}$  and the Gaussian quadrature (3.23) has the degree of precision  $2n-1$ , we also have

$$S_k = \int_{-1}^1 l_k(x)P'_n(x)dx = \sum_{j=1}^n A_j l_k(x_j)P'_n(x_j) = A_k P'_n(x_k).$$

Consequently, by these two equations for  $S_k$ , (3.24), and Part (5) of Theorem 1.26, we obtain

$$\begin{aligned} A_k &= \frac{l_k(x)P_n(x) \Big|_{x=-1}^{x=1}}{P'_n(x_k)} = \frac{[P_n(x)]^2 \Big|_{x=-1}^{x=1}}{(x-x_k)[P'_n(x_k)]^2 \Big|_{x=-1}^{x=1}} \\ &= \left( \frac{[P_n(1)]^2}{1-x_k} - \frac{[P_n(-1)]^2}{-1-x_k} \right) \frac{1}{[P'_n(x_k)]^2} = \frac{2}{(1-x_k^2)[P'_n(x_k)]^2}, \end{aligned}$$

completing the proof. □

### Remainder

$$\frac{f^{(2n)}(\xi)}{(2n)!} \int_{-1}^1 \prod_{k=1}^n (x-x_k)^2 dx = \frac{f^{(2n)}(\xi)}{(2n)!} \int_{-1}^1 \left( \frac{2^n n!}{(2n)!} \right)^2 P_n^2(x) dx = f^{(2n)}(\xi) \frac{2^{2n+1} (n!)^4}{(2n+1)[(2n)!]^3}$$

### The Gauss-Chebyshev quadrature

$$\int_{-1}^1 \frac{f(x)}{\sqrt{1-x^2}} dx \approx \sum_{k=1}^n A_k f(x_k)$$

$x_1, \dots, x_n \in (-1, 1)$  are roots of  $T_n(x) = \cos(n \arccos x)$

$$x_k = \cos \theta_k = \cos \frac{(2k-1)\pi}{2n}, \quad k = 1, \dots, n,$$

$$A_k = \int_{-1}^1 \frac{l_k(x)}{\sqrt{1-x^2}} dx = \frac{\pi}{n}, \quad k = 1, \dots, n.$$

*Proof.* We show that the formula is exact for all  $T_0, \dots, T_{2n-1}$ . Then, the degree of precision is  $\geq 2n-1$ . But any numerical quadrature with  $n$  quadrature points has degree of precision  $\leq 2n-1$ . Thus, this has the degree of precision exactly  $2n-1$ . Therefore, this is the weighted Gaussian quadrature.

Notice that By the change of variable  $x = \cos \theta$ ,

$$\int_{-1}^1 \frac{T_m(x)}{\sqrt{1-x^2}} dx = \int_0^\pi \cos m\theta d\theta = \begin{cases} \pi & \text{if } m = 0, \\ 0 & \text{if } m \neq 0. \end{cases}$$

For  $m = 0$ , we have

$$\frac{\pi}{n} \sum_{k=1}^n T_0(x_k) = \pi.$$

Thus the formula is exact for  $m = 0$ .

Consider now  $1 \leq m \leq 2n-1$ . Denote by  $i$  the complex unit, i.e.,  $i^2 = -1$ . We have  $e^{im\pi} \neq 1$ . Denote by  $\mathcal{R}(z)$  the real part of a complex number  $z$ . We have

$$\begin{aligned} \sum_{k=1}^n T_m(x_k) &= \sum_{k=1}^n \cos \left( \frac{m(2k-1)\pi}{2n} \right) \\ &= \sum_{k=1}^n \mathcal{R} \left( e^{im(2k-1)\pi/(2n)} \right) \\ &= \mathcal{R} \left( e^{-im\pi/(2n)} \sum_{k=1}^n e^{imk\pi/n} \right) \\ &= \mathcal{R} \left( e^{-im\pi/(2n)} e^{im\pi/n} \frac{e^{im\pi} - 1}{e^{im\pi/n} - 1} \right) \\ &= [(-1)^m - 1] \mathcal{R} \left( \frac{e^{im\pi/(2n)}}{e^{im\pi/n} - 1} \right) \\ &= [(-1)^m - 1] \mathcal{R} \left( \frac{e^{im\pi/(2n)} (e^{-im\pi/n} - 1)}{|e^{im\pi/n} - 1|^2} \right) \\ &= [(-1)^m - 1] \mathcal{R} \left( \frac{e^{-im\pi/(2n)} - e^{im\pi/(2n)}}{|e^{im\pi/n} - 1|^2} \right) \end{aligned}$$

$$= 0.$$

$$\int_{-1}^1 \frac{T_m(x)}{\sqrt{1-x^2}} dx = \frac{\pi}{n} \sum_{k=1}^n T_m(x_k), \quad m = 0, \dots, 2n-1.$$

□

Error:  $\frac{\pi}{(2n)!2^{2n-1}} f^{(2n)}(\xi).$

### 3.6 Convergence of Sequences of Numerical Quadrature

**Theorem 3.16 (Convergence of sequences of numerical quadrature).** *Given a sequence of numerical quadrature*

$$\int_a^b \rho(x)f(x) dx \approx \sum_{k=1}^n A_k^{(n)} f(x_k^{(n)}), \quad n = 1, \dots,$$

where  $\rho$  is a weight function on  $[a, b]$ . Suppose

$$(1) \quad \lim_{n \rightarrow \infty} \sum_{k=1}^n A_k^{(n)} p(x_k^{(n)}) = \int_a^b \rho(x)p(x) dx \quad \forall p \in \mathcal{P}, \quad (3.25)$$

$$(2) \quad \sup_{n \geq 1} \sum_{k=1}^n |A_k^{(n)}| < \infty. \quad (3.26)$$

Then

$$\lim_{n \rightarrow \infty} \sum_{k=1}^n A_k^{(n)} f(x_k^{(n)}) = \int_a^b \rho(x)f(x) dx \quad \forall f \in C[a, b].$$

*Proof.* Let  $f \in C[a, b]$ . Denote

$$I(f) = \int_a^b \rho(x)f(x) dx.$$

Denote also for each integer  $n \geq 1$

$$I_n(f) = \sum_{k=1}^n A_k^{(n)} f(x_k^{(n)}).$$

Clearly, each  $I_n : C[a, b] \rightarrow \mathbb{R}$  is linear. Setting

$$M = \sup_{n \geq 1} \sum_{k=1}^n |A_k^{(n)}|,$$

we have

$$|I_n(f)| \leq \sum_{k=1}^n |A_k^{(n)}| \cdot \left| f(x_k^{(n)}) \right| \leq M \|f\|_{C[a,b]} \quad \forall f \in C[a,b] \quad \forall n \geq 1.$$

Now, let  $f \in C[a,b]$  and let  $\varepsilon > 0$ . By the First Weierstrass Approximation Theorem, there exists  $p \in \mathcal{P}$  such that

$$\|f - p\|_{C[a,b]} < \frac{\varepsilon}{2 \left( M + \int_a^b \rho(x) dx \right)}.$$

By the first assumption (3.25), there exists an integer  $N \geq 1$  such that

$$|I_n(p) - I(p)| < \frac{\varepsilon}{2} \quad \forall n \geq N.$$

Therefore, for any  $n \geq N$ ,

$$\begin{aligned} |I_n(f) - I(f)| &\leq |I_n(f - p)| + |I_n(p) - I(p)| + |I(p) - I(f)| \\ &< M \|f - p\|_{C[a,b]} + \frac{\varepsilon}{2} + \|p - f\|_{C[a,b]} \int_a^b \rho(x) dx \\ &< \varepsilon. \end{aligned}$$

This completes the proof. □

**Colloary 3.17.** *Given a sequence of interpolatory numerical quadrature*

$$\int_a^b \rho(x) f(x) dx \approx \sum_{k=1}^n A_k^{(n)} f(x_k^{(n)}), \quad n = 1, \dots$$

*Suppose all the coefficients  $A_k^{(n)}$  ( $k = 1, \dots, n; n = 1, \dots$ ) are positive. Then*

$$\lim_{n \rightarrow \infty} \sum_{k=1}^n A_k^{(n)} f(x_k^{(n)}) = \int_a^b \rho(x) f(x) dx \quad \forall f \in C[a,b]. \quad (3.27)$$

*Proof.* Let  $p \in \mathcal{P}$ . Then there exists an integer  $N \geq 1$  such that  $p \in \mathcal{P}_N$ . Since an interpolatory quadrature with  $n$  quadrature points has the degree of precision  $\geq n - 1$ . Therefore,

$$\sum_{k=1}^n A_k^{(n)} p(x_k^{(n)}) = \int_a^b \rho(x) p(x) dx \quad \forall n \geq N.$$

This shows that the assumption (3.25) in the above theorem holds true. We have also for all  $n \geq 1$  that

$$\int_a^b \rho(x) dx = \sum_{k=1}^n A_k^{(n)} = \sum_{k=1}^n |A_k^{(n)}|,$$

where we used the fact that all  $A_k^{(n)}$  ( $k = 1, \dots, n; n = 1, \dots$ ) are positive. Thus, the second assumption (3.26) in the above theorem holds true. The desired convergence (3.27) then follows from Theorem 3.16.  $\square$

A direct consequence of this corollary and Proposition 3.14 is the following:

**Colloary 3.18 (Convergence of weighted Gaussian quadrature).** *For any sequence of weighted Gaussian quadrature*

$$\int_a^b \rho(x)f(x) dx \approx \sum_{k=1}^n A_k^{(n)} f(x_k^{(n)}), \quad n = 1, \dots,$$

we have

$$\lim_{n \rightarrow \infty} \sum_{k=1}^n A_k^{(n)} f(x_k^{(n)}) = \int_a^b \rho(x)f(x) dx \quad \forall f \in C[a, b]. \quad \square$$

## Exercises

- Use (1) the left endpoint rectangle rule, (2) the midpoint rectangle rule, (3) the trapezoid rule, (4) Simpson's rule, and (5) the two-point Gaussian quadrature to compute the integral

$$\int_0^1 \sin x dx$$

with the number of subintervals of equal length  $n = 1, \dots, 10$ , respectively. Compute also all the corresponding absolute errors.

- Make a table of six columns with one for  $n$  and the other five for the the computed values by the five rules, respectively. Keep eight digits after decimal points.
  - In a single plot, display five curves showing the absolute errors in the log-log scale (i.e.  $\log(\text{error})$  vs.  $\log(n)$ ) for the five corresponding rules.
  - Discuss convergence rates for these quadrature rules based on the computational result.
- Use the trapezoid rule and Simpson's rule to compute the integral

$$\int_0^1 \sin x dx$$

with the number of subintervals of equal length  $n = 2^k$ ,  $k = 0, \dots, 8$ . Then, apply the Richardson extrapolation procedure to the computed values with the trapezoid rule for the pairs  $(2k - 1, 2k)$ ,  $k = 1, \dots, 8$ . Compute all the corresponding absolute errors.

- (a) Make a table of four columns with one for  $n$ , one for the computed values by the trapezoid rule, one for that by the Richardson extrapolation, and one for that by Simpson's rule. Keep eight digits after decimal points.
- (b) In a single plot, display three curves showing in the log-log scale the absolute errors (i.e.  $\log(\text{error})$  vs.  $\log(n)$ ) for the two corresponding rules and the Richardson extrapolation.
- (c) Discuss the computational result in terms of convergence rates.
3. Given a numerical integration formula on  $[-1, 1]$

$$\int_{-1}^1 g(t) dt \approx \sum_{j=1}^n a_j g(t_j). \quad (3.28)$$

Define, for an interval  $[a, b]$ ,  $A_j = (b - a)a_j/2$  and  $x_j = [(b - a)t_j + a + b]/2$ ,  $j = 1, \dots, n$ . Show that the numerical integration formula on  $[a, b]$

$$\int_a^b f(x) dx \approx \sum_{j=1}^n A_j f(x_j)$$

has the same degree of precision as that of the formula (3.28).

4. Find  $A, B, C$  such that the weighted numerical quadrature

$$\int_{-2}^2 |x|f(x) dx \approx Af(-1) + Bf(0) + Cf(1)$$

is exact for polynomials of degree as high as possible. What is the degree of precision of the quadrature?

5. Let  $h > 0$ . Find  $A, B, C, D$  so that the numerical quadrature

$$\int_{-h}^h f(x) dx \approx Af(-h) + Bf(0) + Cf(h) + Dhf'(h)$$

is exact for polynomials of degree as high as possible. What is the degree of precision of the quadrature?

6. Find  $A, B, C, D$  such that the numerical quadrature

$$\int_0^1 f(x) dx \approx Af(0) + Bf(1) + Cf''(0) + Df''(1)$$

is exact for polynomials of degree as high as possible. What is the degree of precision of the quadrature?

7. Consider an interpolatory quadrature

$$\int_a^b f(x) dx \approx \sum_{k=0}^n A_k f(x_k).$$

Define for each integer  $j \geq 0$

$$F_j(t) = \int_a^b (x-t)_+^j dx - \sum_{k=1}^n A_k (x_k - t)_+^j.$$

Show that

$$\int_a^b F_j(t) dt = 0 \quad j = 0, \dots, n-1.$$

8. Consider the trapezoidal formula

$$\int_a^b f(x) dx \approx \frac{1}{2}(b-a)[f(a) + f(b)].$$

- (a) Show that the degree of precision of the formula is  $m = 1$ .
- (b) Calculate explicitly the Peano kernel  $K_1$  of the formula and show that the kernel does not change sign in  $[a, b]$ .
- (c) Let  $f \in C^2[a, b]$ . Show that there exists  $\xi \in (a, b)$  such that

$$\int_a^b f(x) dx - \frac{1}{2}(b-a)[f(a) + f(b)] = -\frac{1}{12}(b-a)^3 f''(\xi).$$

- (d) Let  $N \geq 1$  be an integer,  $h = (b-a)/N$ , and  $x_j = a + jh$ ,  $j = 0, \dots, N$ . Prove for  $f \in C^2[a, b]$  the error formula for the composite trapezoidal formula

$$\int_a^b f(x) dx - \left\{ \frac{h}{2}[f(a) + f(b)] + h \sum_{j=1}^{N-1} f(x_j) \right\} = -\frac{(b-a)f''(\eta)}{12}h^2,$$

where  $\eta \in (a, b)$  depends on  $f$ .

9. Find an integer  $N \geq 1$ , as small as possible, so that

$$\left| \int_0^1 e^x dx - T_N \right| \leq 10^{-12},$$

where  $T_N$  is the numerical integration value (without round-off error) of the function  $e^x$  over  $[0, 1]$  using the composite trapezoidal rule with  $N$  subintervals of equal length.

10. Let  $p_3 \in \mathcal{P}_3$  be the Hermite interpolation polynomial of  $f \in C^1[a, b]$  determined by

$$p_3(a) = f(a), \quad p_3'(a) = f'(a), \quad p_3(b) = f(b), \quad p_3'(b) = f'(b).$$

(a) Show that

$$\int_a^b p_3(x) dx = \frac{1}{2}(b-a)[f(a) + f(b)] - \frac{1}{12}(b-a)^2[f'(b) - f'(a)].$$

(b) Determine the degree of precision of the numerical quadrature

$$\int_a^b f(x) dx \approx \frac{1}{2}(b-a)[f(a) + f(b)] - \frac{1}{12}(b-a)^2[f'(b) - f'(a)]. \quad (3.29)$$

(c) Calculate explicitly the Peano kernel of the numerical quadrature (3.29).

(d) Derive the error formula for the numerical quadrature (3.29).

(e) Let  $N \geq 1$  be an integer,  $h = (b-a)/N$ , and  $x_j = a + jh$ ,  $j = 0, \dots, N$ . Derive the composite integration formula based on the formula (3.29). Show that the composite formula is the same as the composite trapezoidal formula for functions  $f \in C^1[a, b]$  such that  $f'(a) = f'(b)$ .

11. Let  $p \in \mathcal{P}_5$  be the Hermite interpolation polynomial of  $f \in C^1[-1, 1]$  determined by

$$p(x_j) = f(x_j) \quad \text{and} \quad p'(x_j) = f'(x_j), \quad j = 0, 1, 2,$$

where  $x_0 = -1, x_1 = 0, x_2 = 1$ .

(a) Show that

$$\int_{-1}^1 p(x) dx = \frac{1}{15} [7f(-1) + 16f(0) + 7f(1) + f'(-1) - f'(1)].$$

(b) Show that the degree of precision of the numerical integration formula

$$\int_{-1}^1 f(x) dx \approx \frac{1}{15} [7f(-1) + 16f(0) + 7f(1) + f'(-1) - f'(1)]. \quad (3.30)$$

is  $m = 5$ .

(c) Derive the error formula for the numerical integration formula (3.30).

(d) Derive the composite numerical integration formula corresponding to the formula (3.30).

12. Let  $a \leq x_0 < \dots < x_n \leq b$ . Show that there exist  $n + 1$  real numbers  $\gamma_0, \dots, \gamma_n$  such that

$$\int_a^b p(x) dx = \sum_{j=0}^n \gamma_j p(x_j) \quad \forall p \in \mathcal{P}_n.$$

13. Consider the Newton-Cotes formula

$$\int_a^b f(x) dx \approx \sum_{j=0}^n A_j f(x_j)$$

with  $n + 1$  points  $x_j = a + j(b-a)/n$ ,  $j = 0, \dots, n$ .

(a) Show that  $A_j = A_{n-j}$  for  $j = 0, \dots, [n/2]$ .

(b) Show by direct calculation that the degree of precision of the formula is  $n$  if  $n$  is odd and is  $n + 1$  if  $n$  is even.

14. Let  $n \geq 2$  be an even number,  $\omega_n(x) = \prod_{j=0}^n (x - j)$ , and

$$\Omega_n(x) = \int_0^x \omega_n(t) dt.$$

Show that  $\Omega_n(0) = \Omega_n(n) = 0$  and that  $\Omega_n(x) > 0$  for all  $x \in (0, n)$ .

15. Let  $x_1, \dots, x_n$  be  $n$  distinct points in  $[a, b]$  and  $A_1, \dots, A_n$  be  $n$  real numbers. If the degree of precision of the weighted numerical integration formula

$$\int_a^b \rho(x) f(x) dx \approx \sum_{j=1}^n A_j f(x_j)$$

with  $\rho$  a weight function on  $[a, b]$  is  $2n - 1$ , then it must be the weighted Gaussian formula on  $[a, b]$  with the weight function  $\rho$ .

16. Let  $n \geq 2$  be an integer and  $x_1, \dots, x_n$  the  $n$  distinct roots in  $(-1, 1)$  of the  $n$ th Legendre polynomial  $P_n$ . Set

$$l_j(x) = \frac{P_n(x)}{(x - x_j)P_n'(x_j)} \quad \text{and} \quad A_j = \int_{-1}^1 l_j(x) dx, \quad j = 1, \dots, n.$$

- (a) Show that

$$\int_{-1}^1 p(x)q(x) dx = \sum_{j=1}^n A_j p(x_j)q(x_j) \quad \forall p, q \in \mathcal{P}_{n-1}.$$

- (b) Show that

$$A_j = \int_{-1}^1 [l_j(x)]^2 dx > 0, \quad j = 1, \dots, n.$$

17. Let  $\{Q_n\}_{n=0}^\infty$  be a system of orthogonal polynomials with each  $\deg Q_n = n$  with respect to the inner product in  $L_\rho^2[a, b]$ , where  $\rho$  is a weight function on  $[a, b]$ . Fix  $n \geq 1$ . Let  $x_1, \dots, x_n$  be the  $n$  distinct roots of  $Q_n$  in  $(a, b)$ . Let

$$\int_a^b \rho(x) f(x) dx \approx \sum_{j=1}^n A_j f(x_j)$$

be the corresponding weighted Gaussian quadrature. Show that

$$\sum_{j=1}^n A_k Q_k(x_j) = 0, \quad k = 1, \dots, 2n - 1.$$

18. (Gautschi) Consider a weighted Gaussian formula

$$\int_a^b \rho(x)f(x) dx \approx \sum_{j=1}^n A_j f(x_j)$$

with  $\rho$  a weight function on  $[a, b]$ . Show that for any  $f \in C[a, b]$  the error

$$e_n(f) = \int_a^b \rho(x)f(x) dx - \sum_{j=1}^n A_j f(x_j)$$

satisfies

$$|e_n(f)| \leq 2 \left( \int_a^b \rho(x) dx \right) E_{2n-1}(f),$$

where

$$E_{2n-1}(f) = \min_{q \in \mathcal{P}_{2n-1}} \|f - q\|_{C[a,b]}.$$

19. Let  $Q_n \in \mathcal{P}_n$  be the  $n$ th orthogonal polynomial with respect to the weight  $\rho$  on  $[a, b]$ ,  $n = 0, \dots$ . Fix  $n \geq 1$ . Let  $x_1, \dots, x_n$  be the  $n$  distinct roots of  $Q_n$  in  $(a, b)$ . Let

$$\int_a^b \rho(x)f(x) dx \approx \sum_{j=1}^n A_j f(x_j)$$

be the corresponding weighted Gaussian quadrature. Show that

$$A_j = \frac{a_n}{a_{n-1} Q'_n(x_j) Q_{n-1}(x_j)}, \quad j = 1, \dots, n,$$

where  $a_k$  is the leading coefficient of  $Q_k$  ( $k = 0, \dots$ ).

20. Let  $n \geq 1$  be an integer. The Gauss-Chebyshev quadrature is the weighted Gaussian quadrature on  $[-1, 1]$  with the weight  $1/\sqrt{1-x^2}$  using  $x_j = \cos(2j-1)\pi/2n$  ( $j = 1, \dots, n$ ), the  $n$  roots of the  $n$ th Chebyshev polynomial  $T_n(x) = \cos(n \arccos x)$ . Show that the Gauss-Chebyshev formula is given by

$$\int_{-1}^1 \frac{f(x)}{\sqrt{1-x^2}} dx \approx \frac{\pi}{n} \sum_{j=1}^n f(x_j).$$

21. (a) Show for any  $f \in C[0, 1]$  that

$$\int_0^1 B_n f(x) dx = \frac{1}{n+1} \sum_{k=0}^n f\left(\frac{k}{n}\right), \quad n = 0, \dots,$$

where  $B_n f$  ( $n = 0, \dots$ ) are the Bernstein polynomials of  $f$ .

(b) Show that the degree of precision of the numerical integration formula

$$\int_0^1 f(x) dx \approx \frac{1}{n+1} \sum_{k=0}^n f\left(\frac{k}{n}\right)$$

is  $m = 1$  for all  $n = 0, \dots$

(c) Show that

$$\lim_{n \rightarrow \infty} \frac{1}{n+1} \sum_{k=0}^n f\left(\frac{k}{n}\right) = \int_0^1 f(x) dx \quad \forall f \in C[0, 1].$$

22. (Bernoulli polynomials) Define

$$B_0(x) = 1, \quad B_1(x) = x - \frac{1}{2},$$

$$B'_{n+1}(x) = (n+1)B_n(x) \quad \text{and} \quad \int_0^1 B_{n+1}(x) dx = 0, \quad n = 2, \dots$$

(a) Prove that, for each  $n \geq 0$ ,  $B_n$  is a polynomial of degree  $n$  with leading coefficient 1.

(b) Prove the identities

$$\begin{aligned} B_n(x+1) - B_n(x) &= nx^{n-1}, & n = 0, \dots, \\ B_n(1-x) &= (-1)^n B_n(x), & n = 0, \dots \end{aligned}$$

(c) Prove the identities

$$\begin{aligned} B_n(0) &= B_n(1), & n = 2, \dots, \\ B_{2n+1}(0) &= 0, & n = 1, \dots \end{aligned}$$

23. Prove

$$\sum_{k=1}^n k^3 = \left(\frac{n(n+1)}{2}\right)^2$$

by the Euler–Maclaurin summation formula.

24. Let  $f \in C[a, b]$  and denote by  $I(f)$  the integral of  $f$  over  $[a, b]$ . Let  $N \geq 1$  be an integer,  $h = (b-a)/2N$ , and  $x_j = a + jh$ ,  $j = 0, \dots, 2N$ . Let  $T_N$ ,  $T_{2N}$ , and  $S_N$  denote, respectively, the approximate value of  $I(f)$  by the composite trapezoidal rule with  $N$  subintervals  $[x_{2j-1}, x_{2j}]$ ,  $j = 1, \dots, N$ , by the composite trapezoidal rule with  $2N$  subintervals  $[x_{j-1}, x_j]$ ,  $j = 0, \dots, 2N$ , and by the composite Simpson rule with  $N$  subintervals  $[x_{2j-1}, x_{2j}]$ ,  $j = 1, \dots, N$ . Prove that the Richardson extrapolation using  $T_N$  and  $T_{2N}$  leads to exactly  $S_N$ , i.e.,

$$S_N = \frac{4T_{2N} - T_N}{3}.$$

# Bibliography

- [1] E. W. Cheney. *Introduction to Approximation Theory*. McGraw-Hill, 1966.
- [2] P. J. Davis. *Interpolation and Approximation*. Blaisdell, 1963.
- [3] P. J. Davis and P. H. Rabinowitz. *Methods of Numerical Integration*. Academic press, 1975.
- [4] H. Engles. *Numerical Quadrature and Cubature*. Academic Press, 1980.
- [5] E. I. Isaacson and H. B. Keller. *Analysis of Numerical Methods*. Dover, 1994.
- [6] P. P. Korovkin. *Linear Operators and Approximation Theory*. Hindustan, 1960.
- [7] P. M. J. D. Powell. *Approximation Theory and Methods*. Cambridge University Press, 1981.
- [8] T. J. Rivlin. *Introduction to the Approximation of Functions*. Blaisdell, 1969 (Dover, 1987).
- [9] G. Szegő. *Orthogonal Polynomials*. Amer. Math. Soc., 1959.
- [10] J. Todd. *Introduction to the Constructive Theory of Functions*. Academic Press, 1963.
- [11] L.-Z. Xu, R.-H. Wang, and Y.-S. Zhou. *Theory and Methods of Function Approximation*. Shanghai Sci. Tech. Press, 1983 (in Chinese).