

19. AUTOMORPHISM GROUP OF S_n

Definition-Lemma 19.1. *Let G be a group.*

*The **automorphism group of G** , denoted $\text{Aut}(G)$, is the subgroup of $A(S_n)$ of all automorphisms of G .*

Proof. We check that $\text{Aut}(G)$ is closed under products and inverses.

Suppose that ϕ and $\psi \in \text{Aut}(G)$. Let $\xi = \phi \circ \psi$. If g and $h \in G$ then

$$\begin{aligned}\xi(gh) &= (\phi \circ \psi)(gh) \\ &= \phi(\psi(gh)) \\ &= \phi(\psi(g)\psi(h)) \\ &= \phi(\psi(g))\phi(\psi(h)) \\ &= (\phi \circ \psi)(g)(\phi \circ \psi)(h) \\ &= \xi(g)\xi(h).\end{aligned}$$

Thus $\xi = \phi \circ \psi$ is a group homomorphism. Thus $\text{Aut}(G)$ is closed under products.

Now let $\xi = \phi^{-1}$. If g and $h \in G$ then we can find g' and h' such that $g = \phi(g')$ and $h = \phi(h')$. It follows that

$$\begin{aligned}\xi(gh) &= \xi(\phi(g')\phi(h')) \\ &= \xi(\phi(g'h')) \\ &= g'h' \\ &= \xi(g)\xi(h).\end{aligned}$$

Thus $\xi = \phi^{-1}$ is a group homomorphism. Thus $\text{Aut}(G)$ is closed under inverses. \square

Lemma 19.2. *Let G be a group and let $a \in G$. ϕ_a is the automorphism of G given by conjugation by a , $\phi(g) = aga^{-1}$.*

If a and $b \in G$ then

$$\phi_{ab} = \phi_a \phi_b.$$

Proof. Both sides are functions from G to G . We just need to check that they have the same effect on any element g of G :

$$\begin{aligned}(\phi_a \circ \phi_b)(g) &= \phi_a(\phi_b(g)) \\ &= \phi_a(bgb^{-1}) \\ &= a(bgb^{-1})a^{-1} \\ &= (ab)g(ab)^{-1} \\ &= \phi_{ab}(g).\end{aligned}$$

\square

Definition-Lemma 19.3. We say that an automorphism ϕ of G is **inner** if $\phi = \phi_a$ for some a . The **inner automorphism group** of G , denoted $\text{Inn}(G)$, is the subgroup of $\text{Aut}(G)$ given by inner automorphisms.

Proof. We check that $\text{Inn}(G)$ is closed under products and inverses.

We checked that $\text{Inn}(G)$ is closed under products in (19.2). Suppose that $a \in G$. We check that the inverse of ϕ_a is $\phi_{a^{-1}}$. We have

$$\begin{aligned}\phi_a \phi_{a^{-1}} &= \phi_{aa^{-1}} \\ &= \phi_e,\end{aligned}$$

which is clearly the identity function. Thus $\text{Inn}(G)$ is closed under inverses. \square

Definition-Lemma 19.4. Let G be a group.

Then the inner automorphism group is a normal subgroup of $A(G)$. The quotient group $\text{Aut}(G)/\text{Inn}(G)$ is called the **outer automorphism group** of G , denoted $\text{Out}(G)$.

Proof. Let f be a permutation of G and let ϕ_a be an inner automorphism. Let $b = f(a)$. We check $f\phi_a f^{-1} = \phi_b$. Since both sides are functions from G to G we just need to check they have the same effect on every element g of G . Suppose that $g = f(h)$. We have

$$\begin{aligned}(f\phi_a f^{-1})(g) &= (f\phi_a f^{-1})(f(h)) \\ &= f\phi_a(h) \\ &= f(aha^{-1}) \\ &= f(a)f(h)f(a^{-1}) \\ &= bgb^{-1} \\ &= \phi_b(g).\end{aligned}$$

\square

Lemma 19.5. Let G be a group with centre Z .

Then $\text{Inn}(G) \simeq G/Z$.

Proof. Define a function

$$A: G \longrightarrow \text{Inn}(G) \quad \text{by sending} \quad a \longrightarrow \phi_a.$$

A is a group homomorphism by (19.2). A is clearly surjective. We identify the kernel. $a \in \text{Ker } A$ if and only if ϕ_a is the identity if and only if $\phi_a(g) = g$ for all $g \in G$ if and only if $aga^{-1} = g$ for all $g \in G$ if and only if $ag = ga$ for all $g \in G$ if and only if $a \in Z$.

Now apply the first Isomorphism Theorem. \square

Theorem 19.6. $\text{Aut}(S_n) = \text{Inn}(S_n) \simeq S_n$ unless

- (1) $n = 2$ when $\text{Aut}(S_n) = \text{Inn}(S_n) = \{e\}$.
- (2) $n = 6$ when $\text{Inn}(S_n) = S_n$ and $\text{Out}(S_n) = \mathbb{Z}_2$.

Observe that (19.6) says that most automorphisms of S_n are inner. We first compute the centre of S_n :

Lemma 19.7. *The centre of S_n is S_n unless $n = 2$.*

Proof. We may assume that $n \geq 3$. Suppose that $\sigma \in S_n$ is not the identity. Pick i such that $j = \sigma(i) \neq i$. Pick $k \notin \{i, j\}$ and let $\tau = (j, k)$. Then $\tau\sigma\tau^{-1}$ sends i to k . Thus

$$\tau\sigma\tau^{-1} \neq \sigma,$$

so that σ does not belong to the centre. □

Note that an inner automorphism of S_n preserves cycle type. We show the converse is true.

Lemma 19.8. *Suppose that G is a group and S is a set of generators of G .*

If ϕ_1 and ϕ_2 are two automorphisms of G that agree on S then $\phi_1 = \phi_2$.

Proof. Let H be the largest subset of G on which ϕ_1 and ϕ_2 agree. We show that H is a subgroup of G . $e \in H$ and so H is non-empty. Suppose that g and h belong to H . We have

$$\begin{aligned} \phi_1(gh) &= \phi_1(g)\phi_1(h) \\ &= \phi_2(g)\phi_2(h) \\ &= \phi_2(gh). \end{aligned}$$

Thus $gh \in H$. Thus H is closed under products.

Suppose that $g \in H$. We have

$$\begin{aligned} \phi_1(g^{-1}) &= \phi_1(g)^{-1} \\ &= \phi_2(g)^{-1} \\ &= \phi_2(g^{-1}). \end{aligned}$$

Thus $g^{-1} \in H$ and so H is closed under inverses. Thus H is a subgroup of G .

As H contains S , $H = G$, and so $\phi_1 = \phi_2$. □

Lemma 19.9. *If $\phi \in \text{Aut}(S_n)$ sends transpositions to transpositions then ϕ is inner.*

Proof. The transpositions $(i, i+1)$, $1 \leq i \leq n-1$ generate S_n . Suppose that $(\alpha, \beta) = \phi(i, i+1)$ and $(\gamma, \delta) = \phi(i+1, i+2)$. Possibly rearranging we may assume that $\beta = \delta$. Thus we may assume that there are a_1, a_2, \dots, a_n such that $(a_i, a_{i+1}) = \phi(i, i+1)$. Let $\tau(i) = a_i$. Then τ is a permutation of the first n natural numbers and ϕ and ϕ_τ agree on the generators $(i, i+1)$. (19.8) implies that $\phi = \phi_\tau$ so that ϕ is inner. \square

Lemma 19.10. *Let $C \subset S_n$ be a conjugacy class with $\binom{n}{2}$ elements of order 2.*

Then either C consists of transpositions or $n = 6$ and C consists of the product of three disjoint transpositions.

Proof. The only permutations of order two are the product of k disjoint transpositions. In this case the cycle type is $1^{n-2k}2^k$. Conjugacy in S_n is determined by cycle type. The number of permutations with cycle type 2^k is

$$\frac{1}{k!} \binom{n}{2} \binom{n-2}{2} \cdots \binom{n+2-2k}{2}.$$

For this to equal $\binom{n}{2}$ we must have

$$\frac{1}{(k-1)!} \binom{n-2}{2} \cdots \binom{n+2-2k}{2} = k.$$

Note that the LHS counts the number of permutations with cycle type $1^{n-2k}2^{k-1}$.

If $k = 1$ then both sides are equal to one. So suppose $k \geq 2$. The number of permutations in S_{n-2} which are the product of $k-1$ disjoint transpositions is at least the number of ways to pair $2(k-1)$ objects. As $k > 1$, this is at least the number of ways to pair the first element with any other element, which is $n-2-1 = n-3$. So we must have $n-3 \leq k$, that is, $n \leq k+3$.

As $2k \leq n$ we must have $k \leq 3$. In this case $n \leq 6$. If $k = 3$ then $n = 6$ and we get equality. If $k = 2$ then $4 \leq n \leq 5$. If $n = 4$ the LHS is 1, not 2, and if $n = 5$ the LHS is 3, not 2. \square

If we put everything we have done together it remains to show that if $n = 6$ then the outer automorphism is non-trivial.

Lemma 19.11. *The order of $\text{Out}(S_6)$ is at most two.*

Further the order is two if and only if there is an automorphism ϕ of S_6 which sends a transposition to a product of three disjoint transpositions.

Proof. We have already seen that an automorphism is inner if it fixes the subset of all transpositions. By (19.10) if we don't send a transposition to a transposition then we must send it to product of three disjoint transpositions. \square

It is actually suprisingly involved to write down an automorphism ϕ which sends a transposition to a product of three disjoint transpositions. The problem is that there are too many choices. Outer automorphisms are really equivalence classes, left cosets of the inner automorphism group. Writing down an explicit automorphism which is not inner is somehow completely the opposite to what we have done so far, there don't seem to be any natural choices.

In our case there are $6! = 720$ inner automorphisms and so ϕ belongs to a left coset with 720 elements. We start by figuring out how ϕ acts on the other conjugacy classes. It is useful to write down a table of conjugacy classes, the order of a typical element and their sizes:

Type	Order	Size
e	1	1
$(1, 2)$	2	15
$(1, 2)(3, 4)$	2	45
$(1, 2)(3, 4)(5, 6)$	2	15
$(1, 2, 3)$	3	40
$(1, 2, 3)(4, 5, 6)$	3	40
$(1, 2, 3, 4)$	4	90
$(1, 2, 3, 4)(5, 6)$	4	90
$(1, 2, 3, 4, 5)$	5	144
$(1, 2, 3, 4, 5, 6)$	6	120
$(1, 2)(3, 4, 5)$	6	120

As a check, the sum of the numbers in the last column is $720 = 6!$ the order of S_n .

Note that all of these conjugacy classes come in pairs C_1 and C_2 , where the order of the elements of C_1 and C_2 are the same and the cardinality of C_1 and C_2 is the same, with three exceptions. Presumably an outer automorphism switches C_1 and C_2 . $(1, 2)$ is paired with $(1, 2)(3, 4)(5, 6)$; $(1, 2, 3)$ is paired with $(1, 2, 3)(4, 5, 6)$; $(1, 2, 3, 4)$ is paired with $(1, 2, 3, 4)(5, 6)$; $(1, 2, 3, 4, 5, 6)$ is paired with $(1, 2, 3)(4, 5)$. The classes represented by e , $(1, 2)(3, 4)$ and $(1, 2, 3, 4, 5)$ are paired with themselves. This suggests that 5-cycles play a special role.

The construction of an outer automorphism is quite involved; the interested reader might look online for all of the details. The idea is to

find an injective group homomorphism $\pi: S_5 \longrightarrow S_6$ which is different from the obvious inclusion.

Take the complete graph with 5 vertices and colour the ten edges red and blue so that there is one red 5-cycle and one blue 5-cycle. After a little bit of drawing pictures, it is not hard to see there are six ways to do this. Permuting the five vertices permutes the six ways to colour. This defines π .

Note that the kernel of π is one of the following normal subgroups: $\{e\}$, A_5 and S_5 . It is not hard to check that $(1, 2, 3) \in A_5$ is not in the kernel so that the kernel is $\{e\}$ and so π is injective. It is also not hard to see that the transposition $(1, 2)$ is sent to a product of three disjoint transpositions.

Let H be the image of S_5 . Then H is a subgroup of S_6 of index $6 = 6!/5!$. S_6 acts on the left cosets of H in S_6 and this defines a homomorphism $\phi: S_6 \longrightarrow S_6$. Again the kernel is one of three possible normal subgroups $\{e\}$, A_6 or S_6 . It is again easy to see the kernel of ϕ is $\{e\}$. It follows that ϕ is injective, so that ϕ is a bijection. Once again, it is not hard to check that the image of a transposition is not a transposition, so that ϕ corresponds to an outer automorphism.