

On Arithmetic Progressions of Cycle Lengths in Graphs

Jacques Verstraëte

Department of Pure Mathematics and Mathematical Statistics

Centre for Mathematical Sciences

Wilberforce Road, Cambridge CB3 0WB

England August 1999.

jbav@microsoft.com

Abstract

A recently posed question of Häggkvist and Scott's asked whether or not there exists a constant c such that if G is a graph of minimum degree ck then G contains cycles of k consecutive even lengths. In this paper we answer the question by proving that for $k \geq 2$, a bipartite graph of average degree at least $4k$ and girth g contains cycles of $(g/2 - 1)k$ consecutive even lengths. We also obtain a short proof of the theorem of Bondy and Simonovits, that a graph of order n and size at least $8(k-1)n^{1+1/k}$ has a cycle of length $2k$.

Erdős and Burr [4] conjectured that for every odd number k , there is a constant c_k such that for every natural number m , every graph of average degree at least c_k contains a cycle of length m modulo k . Erdős and Burr [4] settled their conjecture in the case $m = 2$ and Robertson (see [4]) settled the case $m = 0$. The full conjecture was resolved by Bollobás [1], who proved the conjecture with $c_k = 2[(k+1)^k - 1]/k$. In this paper, we show that $c_k = 8k$ will do. Thomassen [11] later showed cycles of all even lengths modulo k are obtained under the hypothesis that the average degree is at least $4k(k+1)$, without requiring k to be odd. Thomassen [10] also proved that if G is a graph of minimum degree at least three and girth at least $2(k^2 + 1)(3 \cdot 2^{k^2+1} + (k^2 + 1)^2 - 1)$, then G contains cycles of all even lengths modulo k .

Bondy and Vince [3] proved that in a graph in which all but at most two vertices have degree at least three, there exist two cycles whose lengths differ by at most two. This answered a conjecture of Erdős and was also studied by Häggkvist and Scott [8]. Recently, Häggkvist and Scott [7], considered extending this to considering arithmetic progressions of cycle lengths in graphs. Häggkvist and Scott [7] proved that if G is a graph of minimum degree at least $300k^2$ then G contains $k + 1$ consecutive even cycle lengths. The same authors asked if a linear bound on the minimum degree is possible. In this paper, we answer the question of Häggkvist and Scott in the following theorem.

Theorem 1 *Let $k \geq 2$ be a natural number and G a bipartite graph of average degree at least $4k$ and girth g . Then there exist cycles of $(g/2 - 1)k$ consecutive even lengths in G . Moreover, the shortest of these cycles has length at most twice the radius of G .*

This result generalises the above-mentioned result of Bollobás and partly generalises that of Thomassen, insofar as Thomassen's result is valid for graphs of minimum degree at least three, whereas the result above requires average degree at least eight. The graph $K_{k,n-k}$ shows that we require the average degree to be at least about $2k$ to ensure the conclusion of Theorem 1.

The following lemma lies at the heart of the proof of Theorem 1. It was originally inspired by methods used by Gyárfás, Komlós and Szemerédi [6]. Whilst this paper was being written, the lemma was discovered to be implicit in a lemma of Bondy and Simonovits [2]. Nevertheless, the proof is short and is retained here for completeness.

Lemma 2 *Let H be a graph comprising a cycle with a chord. Let (A, B) be a non-trivial partition of $V(H)$. Then H contains A - B paths of every length less than $|H|$, unless H is bipartite with bipartition (A, B) .*

Proof. Label the vertices of the cycle $0, 1, \dots, n - 1$ where $n = |H|$. Suppose H does not contain A - B paths of every length less than n , and let

m be the smallest integer for which there is no A - B path of length m not using the chord; $m > 1$ since (A, B) is a non-trivial partition of $V(H)$. We remark also that $m \leq n/2$, or H would contain A - B paths of all lengths less than n .

Now $\chi(j) = \chi(j + m)$ for every $j \in V(H)$, where χ is the characteristic function of A (label arithmetic is modulo n). Let $d = \text{hcf}(n, m)$. Then there are integers p and q such that $pm + qn = d$; hence $\chi(j) = \chi(j + d)$ for every j . But then there is no A - B path of length d round the cycle; thus $d = m$ and $m \mid n$. In particular, A - B paths of every length less than m exist by the definition of m , so A - B paths of every length other than multiples of m exist by periodicity of χ .

We find paths of the remaining lengths km , $1 \leq k \leq n/m - 1$, using the chord. Suppose first that the chord joins two vertices within distance m on the cycle, say 0 and r where $1 < r \leq m$. There exist A - B paths of length $m+r-1$ round the cycle; thus $\chi(j) \neq \chi(j+m+r-1)$ for some j , $-m < j \leq 0$. But $j + m + r - 1 \geq r$, so the path $j, j + 1, \dots, 0, r, r + 1, \dots, j + km + r - 1$ is an A - B path of length km provided $j + km + r - 1 < n + j$, which holds for all the desired $k \leq n/m - 1$.

So we may suppose the chord is $0r$, where $m < r < n - m$. Let $-m < j < 0$ and consider the paths $j, j + 1, \dots, 0, r, r - 1, \dots, r - j - m + 1$ and $m + j, m + j - 1, \dots, 0, r, r + 1, \dots, r - j - 1$, of length m . If either of them is an A - B path we can extend it, by m vertices at a time, to A - B paths of lengths km , $k \geq 1$, until the number of unused vertices in the two arcs defined by the chord is less than m in each arc. At this point $km + 1 \geq n - 2(m - 1)$, and as $m \mid n$, $km = n - m$ as desired. Likewise, if either of the two paths $0, r, r - 1, \dots, r - m + 1$ and $0, r, r + 1, r + m - 1$ is an A - B path then H contains paths of all lengths less than $|H|$.

Thus it follows, as $\chi(j) = \chi(m + j)$, that for $-m < j < 0$ we have $\chi(r - j - m + 1) = \chi(r - j - 1)$ and that $\chi(r - m + 1) = \chi(r + m - 1)$, implying $\chi(r - j + 1) = \chi(r - j - 1)$ and $\chi(r + m + 1) = \chi(r + m - 1)$. So $\chi(v + 2) = \chi(v)$ for all $r \leq v < r + m$, and so for all $v \in V(H)$. Hence $m = 2$.

We conclude, therefore, that $|H|$ is even and the vertices of the cycle are alternately in A and in B . It is immediately seen that, under these circumstances, if the chord joins two vertices in the same class then H contains A – B paths of all lengths less than $|H|$. Consequently, the chord joins A to B , so H is bipartite, with bipartition (A, B) . \square

Lemma 3 *Let $k \geq 2$ be a natural number and let G be a graph of average degree at least $2k$ and girth g . Then G contains a cycle of length at least $(g - 2)k + 2$, with at least one chord.*

Proof. It is easily seen that a graph G of average degree at least $2k$ contains a subgraph H of minimum degree at least $k + 1$. If P is a longest path in H , then an endvertex v of P has all its neighbours on P . Some neighbour u of v is at distance at least $(g - 2)k + 1$ from v on P . Hence $P + uv$ is a cycle of length at least $(g - 2)k + 2$. As $k + 1 \geq 3$, this cycle has at least one chord. \square

Proof of Theorem 1. We may assume that G is connected and let the radius of G be $\text{rad}(G)$. Choose a central vertex $v_0 \in V(G)$, and let V_i denote the set of vertices a distance i from v_0 in G . Then there exists l such that $V_l \cup V_{l+1}$ spans a graph G' with at least $k|V_l \cup V_{l+1}|$ edges. By Lemma 3, find $H \subset G'$ comprising a cycle, of length at least $(g - 2)k + 2$, with a chord. Let T' be a minimal subtree of T , restricted to $\bigcup_{i \leq l} V_i$, such that T' contains $V(H) \cap V_l$. The minimality of T' ensures that it branches at its root. Now let A be the set of vertices of H in one of these branches and let $B = V(H) \setminus A$. By Lemma 2, and as (A, B) is not the bipartition of H , there are A – B paths of all lengths up to $(g - 2)k + 1$, all disjoint from $T' - \text{end}(T')$. Each A – B path of even length s , together with a subpath of T' between the ends of such a path, gives rise to a cycle of length $s + 2r$, where r is the distance from V_l to the root of T' . Note that, as G is bipartite, all paths of even length with one end in A have their other end in V_l . This gives cycles $C_{2r+2}, C_{2r+4}, \dots, C_{2r+(g-2)k}$, of $(g/2 - 1)k$ consecutive even lengths, and since v_0 is a central vertex, $2r + 2 \leq \text{rad}(G)$, as required. \square

We define the even girth of a graph G to be the length of a shortest even cycle in G . Theorem 1 easily extends to general graphs, as is shown by the following corollary.

Corollary 4 *Let $k \geq 2$ be a natural number, and let G be a graph of average degree at least $8k$ and even girth g . Then there are cycles of $(g/2 - 1)k$ consecutive even lengths in G .*

Proof. This follows from the observation that a graph of average degree at least $8k$ has a spanning bipartite subgraph of average degree at least $4k$, and then applying Theorem 1 to this bipartite subgraph. \square

In the case of graphs of average degree at least $6k$, we may also argue as follows. Given a vertex v_0 , let V_l denote the vertices a distance l from v_0 . Then either there exists l such that $V_l \cup V_{l+1}$ spans a bipartite graph of average degree at least $2k$, or there exists l such that V_l spans a graph of average degree at least $2k$. In the former case, the method of Theorem 1 gives cycles of all even lengths in an integer interval of form $[2r+1, 2r+(g-2)k+1]$ and the latter case gives (also following the proof of Theorem 1) cycles of all odd lengths in an integer interval of form $[2r+1, 2r+(g-2)k+1]$. So we have the following theorem:

Theorem 5 *Let $k \geq 2$ be a natural number, and let G be a graph of average degree at least $6k$ and girth g . Then, for some odd number $r \geq 3$, there exist cycles of all even lengths or all odd lengths in the interval $[r, r+(g-2)k]$.*

The above result recalls the result of Bondy and Vince [3], that if G is a graph with at most two vertices of degree at most two, then G contains cycles of two consecutive lengths or two consecutive even lengths. This was proved using a technique of Thomassen and Toft [12]. In comparison, note that Theorem 5 requires average degree at least $6k$ where $k \geq 2$. Therefore, to ensure two cycles of consecutive lengths or consecutive even lengths, we require average degree at least twelve, which is higher than what is required in the context of Bondy and Vince's results.

The following result is proved in the same way as Theorem 1:

Corollary 6 *Let $k \geq 2$ be a natural number, and suppose that G has chromatic number at least $2k + 2$ and girth at least g . Then G contains cycles of $k(g - 2)$ consecutive lengths.*

The idea is that some level of a breadth-first search tree induces a graph of chromatic number at least $k + 1$. Such a graph contains an odd cycle of length at least $k(g - 2) + 1$ with a chord. We then apply the colouring lemma to deduce that G contains cycles of $k(g - 2)$ consecutive lengths. In a sense, this generalizes a result of Gyárfás who showed that if G has chromatic number at least $2k + 2$, then G contains cycles of k distinct odd lengths. These ideas may also be used to give a relatively short proof that $r(C_{2k+1}, K_n) \ll n^{1+1/(k+1)}$. However, better results are easily obtain — for example, it is possible to show that $r(C_{2k+1}, K_n) \ll n^{1+1/(k+1)}(\log_2 n)^{-1/k}$.

The cycles we have obtained are all very close together in the sense that they share many vertices. Häggkvist and Scott [7] asked if it was possible, under an appropriate bound on the size of the graph, to find disjoint cycles of k consecutive even lengths. This question also remains open, noting that a bound of order at least k^2 on the average degree would be required for disjoint cycles of k consecutive even lengths. This is shown, for example, by $K_{l, n-l}$ where $l < 2k + k(k - 1)/2$ and n is sufficiently large.

We remark that from Theorem 1 we may obtain a result on extremal numbers for even cycles, that slightly improves the result obtained by Bondy and Simonovits [2] (see Corollary 9). From their paper, it follows that a graph of order n and size at least $90kn^{1+1/k}$ contains a cycle of length $2k$. Two simple lemmas are required before proving our result. The first lemma is a special case of a lemma of Kostochka and Pyber [9].

Lemma 7 *Let G be a graph of order n and size at least $cn^{1+1/k}$, where $c \geq 1$. Then G contains a subgraph of average degree at least c and radius at most k .*

Proof. We may assume that G has minimum degree at least $cn^{1/k}$. Let v_0 be an arbitrary vertex in G and define H_i to be the subgraph of G induced by

vertices at distance at most i from v_0 . Define $r = \min\{i : e(H_i) \geq \frac{1}{2}c|H_i|\}$. Clearly $e(H_i) \geq \frac{1}{2}cn^{1/k}|H_{i-1}|$ for all i and so, by definition of r , $|H_{r-1}| > n^{1/k}|H_{r-2}|$ which gives $|H_{r-1}| > n^{(r-1)/k}$. Since $|H_{r-1}| \leq n$, $r - 1 < k$ and so $r \leq k$ and H_r is the desired subgraph. \square

Lemma 8 *Let G be a graph of order n with $e(G) \geq 2n^{1+1/k}$ where $k \geq 2$. Then G has girth at most $2k + 1$.*

Proof. By Lemma 6, G has a subgraph H of radius at most k and average degree at least two. So H contains a cycle and some cycle in H has length at most $2k + 1$. \square

In particular, if G is bipartite and $k = 2$ in Lemma 7, then G has a cycle of length four. For comparison, a standard result states that a graph which has at least $n^{3/2}/2 + n/4$ edges contains a cycle of length four. We are able, using Theorem 1, to show the existence of longer even cycles:

Theorem 9 *Let G be a bipartite graph of order n and girth g , and of size at least $4\lceil 2(k-1)/(g-2) \rceil n^{1+1/k}$, where $k \geq 2$ is an integer. Then G has a cycle of length $2k$.*

Proof. By Lemma 7, either G contains a cycle of length $2k$ or $g < 2k$ and $k \geq 3$. In the latter case, $\lceil 2(k-1)/(g-2) \rceil \geq 2$. Lemma 6 shows that G contains a subgraph of average degree at least $4\lceil 2(k-1)/(g-2) \rceil$ and of radius at most k . By Theorem 1, there are cycles of at least $k-1$ consecutive even lengths in H , the shortest length being at most $2k$. So one of these cycles must have length exactly $2k$. \square

As a corollary to Theorem 8, we slightly improve the result of Bondy and Simonovits [2]:

Corollary 10 *Let G be a graph of order n and size at least $8(k-1)n^{1+1/k}$, where $k \geq 2$. Then G contains a cycle of length $2k$.*

Acknowledgements

I would like to thank Andrew Thomason for his many helpful suggestions.

References

- [1] Bollobás, B., Cycles Modulo k , *Bull. London Math. Soc.* **9** (1977) 97–98.
- [2] Bondy, J., Simonovits, M., Cycles of Even Length in Graphs, *J. Combinatorial Theory B* **16** (1974) 97–105.
- [3] Bondy, J., Vince, A. Cycles in a graph whose lengths differ by one or two, *J. Graph Theory* **27** (1998) 11–15.
- [4] Erdős, P., Some recent problems and results in graph theory, combinatorics, and number theory, Proc. Seventh S–E Conf. Combinatorics, Graph Theory and Computing, *Utilitas Math.*, Winnipeg, (1976) 3–14.
- [5] Erdős, P., Some of my recent problems in Combinatorial Number Theory, Geometry and Combinatorics in: *Graph Theory, Combinatorics and Algorithms*, Volume 1, Proc. Seventh Quadrennial International Conference on the Theory and Applications of Graphs, Y. Alavi and A. Schwenk eds., John Wiley and Sons (1995) 335–349.
- [6] Gyárfás, A., Komlós, J., Szemerédi, E., On the distribution of cycle lengths in graphs, *J. Graph Theory* **8** (1984) 441–466.
- [7] Häggkvist, R., Scott, A., Arithmetic progressions of cycles, preprint.
- [8] Häggkvist, R., Scott, A., Cycles of nearly equal length in cubic graphs, preprint.
- [9] Kostochka, A., Pyber, L., Small topological complete subgraphs of dense graphs, *Combinatorica* **8**(1) (1988) 83–86.
- [10] Thomassen, C., Girth in Graphs, *J. Combinatorial Theory B* **35** (1983) 129–141.
- [11] Thomassen, C., Paths, Circuits and Subdivisions in: *Selected Topics in Graph Theory 3*, L. Beineke, R. Wilson eds., Academic Press (1988) 97–133.

- [12] Thomassen, C., Toft, B., Non-separating induced cycles in graphs, *J. Combinatorial Theory B* **31** (1981) 199–224.