

A Note on Bipartite Graphs without $2k$ -Cycles.

ASSAF NAOR and JACQUES VERSTRAËTE

Microsoft Research, One Microsoft Way, Redmond, WA 98052-6399, United States.

The question of the maximum number $\text{ex}(m, n, C_{2k})$ of edges in an m by n bipartite graph without a cycle of length $2k$ is addressed in this note. For each $k \geq 2$, it is shown that

$$\text{ex}(m, n, C_{2k}) \leq \begin{cases} (2k-3)[(mn)^{\frac{k+1}{k}} + m + n] & \text{if } k \text{ is odd.} \\ (2k-3)[m^{\frac{k+2}{k}} n^{\frac{1}{2}} + m + n] & \text{if } k \text{ is even.} \end{cases}$$

1. Introduction

In this note, we study the maximum number of edges in an m by n bipartite graph containing no $2k$ -cycles. This problem was studied in the papers by Erdős, Sós and Sárközy [3] and by Györi [4], in the context of a number theoretic problem. It is also related to the size of subsets of points in projective planes, such as arcs and caps. A connection to a geometric problem involving points and lines in Euclidean space is described in de Caen and Szekély [1].

Throughout the material to follow, $\gamma(m, n, g, k)$ denotes the maximum number of edges in an m by n bipartite graph of girth at least $2g$ containing no $2k$ -cycle. In particular, we write $\gamma(m, n, 2, k) = \text{ex}(m, n, C_{2k})$, which is the maximum number of edges in a $2k$ -cycle-free m by n bipartite graph. For $d, g \geq 2$, let $c(d, g)$ be the largest integer such that every bipartite graph of average degree at least $2d$ and girth at least $2g$ contains a cycle of length at least $c(d, g)$ with at least one chord. Our main result is as follows:

Theorem 1. *Let $k, g \geq 2$ be integers. Then, for any d such that $c(d, g) \geq 2(k - g + 1)$,*

$$\gamma(m, n, g, k) \leq \begin{cases} 2d \left[(mn)^{\frac{k+1}{2k}} + m + n \right] & \text{if } k \text{ is odd.} \\ 2d \left[m^{\frac{k+2}{2k}} n^{\frac{1}{2}} + m + n \right] & \text{if } k \text{ is even.} \end{cases}$$

Theorem 1 depends explicitly on the parameter $c(d, g)$. Let us make a few remarks about $c(d, g)$. First, a straightforward argument shows that $c(d, g) \geq 2d(g - 1) + 1$. Stronger results were obtained by Erdős, Faudree, Rousseau, and Schelp [2]. We deduce from their paper that $c(d, g) \geq d^{g/4}$. In particular, by using these two bounds on $c(d, g)$ in Theorem 1, we respectively obtain the following two corollaries:

Corollary 2.

$$ex(m, n, C_{2k}) \leq \begin{cases} (2k-3) \left[(mn)^{\frac{k+1}{2k}} + m + n \right] & \text{if } k \text{ is odd.} \\ (2k-3) \left[m^{\frac{k+2}{2k}} n^{\frac{1}{2}} + m + n \right] & \text{if } k \text{ is even.} \end{cases}$$

Corollary 3. For any number $\delta > 0$, there exists a constant $c(\delta)$ such that

$$\gamma(m, n, \delta \log k, k) \leq \begin{cases} c(\delta) \left[(mn)^{\frac{k+1}{2k}} + m + n \right] & \text{if } k \text{ is odd.} \\ c(\delta) \left[m^{\frac{k+2}{2k}} n^{\frac{1}{2}} + m + n \right] & \text{if } k \text{ is even.} \end{cases}$$

Using the requirement on $c(d, g)$ from Theorem 1 and the inequality $c(d, g) \geq d^{g/4}$, a short computation shows that we may certainly take $c(\delta) = 4^{4/\delta}$ in Corollary 3. Recent results of Hoory [5] and of Lam [7] show that $\gamma(m, n, k, k)$ satisfies the bounds given in Corollary 3, with $c(\delta) = 1$. In the case $k \in \{2, 3, 5\}$, the existence of rank two geometries known as generalized polygons (see [6] for constructions) show that the constant $c(\delta) = 1$ is best possible when $m = n$. The strength in Corollary 3 is that it shows that excluding cycles of length $O(\log k)$ has, to within an absolute constant factor, the same effect on the upper bounds as excluding all cycles of length at most $2k$. On the other hand, our next result gives an indication that $\gamma(m, n, k, k)$ and $\gamma(m, n, 2, k)$ may differ substantially:

Theorem 4. For all integers m, n and $k \geq 3$,

$$\gamma((k-1)m, n, k, k) \geq (k-1) \cdot \gamma(m, n, 2, k).$$

If we make the assumption that for each even positive integer k , $\gamma(m, n, 2, k)$ is asymptotically $c_1(mn)^{1/2+1/(2k)} + c_2(m+n)$ as $m, n \rightarrow \infty$, for some constants c_1, c_2 , then Theorem 4 gives

$$\liminf_{m, n \rightarrow \infty} \frac{\gamma(m, n, k, k)}{\gamma(m, n, 2, k)} \geq \sqrt{k-1}.$$

Similar observations may be made for odd values of k , namely

$$\liminf_{m, n \rightarrow \infty} \frac{\gamma(m, n, k, k)}{\gamma(m, n, 2, k)} \geq (k-1)^{1/2-1/(2k)}.$$

One can deduce from these observations and the constructions of generalized quadrangles and hexagons that

$$\begin{aligned} \liminf_{n \rightarrow \infty} \frac{\gamma(n, 2n, 3, 3)}{\gamma(n, 2n, 2, 3)} &\geq 2^{1/3} \\ \liminf_{n \rightarrow \infty} \frac{\gamma(n, 2n, 5, 5)}{\gamma(n, 2n, 2, 5)} &\geq 4^{1/5} \end{aligned}$$

The first inequality comes from [8] and the second inequality is implicit in the work of Lazebnik, Ustimenko and Woldar [6]. The next section is devoted to proving Theorem 1, and the construction for Theorem 4 is presented in Section 3.

2. Proof of Theorem 1

The main tool in our proof will be the first theorem in [9]. Although the next proposition is not the statement of this theorem, it is straightforward to verify, from the proof appearing in [9]:

Proposition 5. *Let G be a bipartite graph of average degree at least $4d$ and girth $2g$. Then G contains cycles of $\frac{1}{2}c(2d, 2g)$ consecutive even lengths, the shortest of which has length at most twice the radius of G .*

Proof of Theorem 1 We proceed by induction on $m+n$. Suppose, for a contradiction, that G is a bipartite graph without cycles of length at most $2g-2$ and containing no $2k$ -cycle, with parts A and B of sizes m and n , respectively, and with more edges than the corresponding upper bound in Theorem 1. By Proposition 5, and using the definition of d in the statement of Theorem 1, it is sufficient to show that G contains a subgraph of radius at most k and average degree at least d . Indeed, since G has even girth $2g$, Proposition 5 would then imply that there is an integer r such that $g \leq r \leq k$ and such that G contains the cycles $C_{2r}, C_{2r+2}, \dots, C_{2r+2k-2g}$. Since $2k \in [2r, 2r+2k-2g]$ whenever $g \leq r \leq k$, one of these cycles has length $2k$, as required.

Case 1. k is odd.

In this case, we may assume that the minimum degree in A is at least

$$d_A = d \frac{m^{\frac{1}{2} + \frac{1}{2k}}}{n^{\frac{1}{2} - \frac{1}{2k}}} + d.$$

Indeed, if there was a vertex $v \in A$ with degree less than d_A then by deleting it we would arrive at an $m \times (n-1)$ bipartite graph G' with

$$\begin{aligned} e(G') &> e(G) - d_A &\geq 2dm^{\frac{1}{2} + \frac{1}{2k}} \left(n^{\frac{1}{2} + \frac{1}{2k}} - \frac{1}{2}n^{-\frac{1}{2} + \frac{1}{2k}} \right) + 2d(m+n) - d \\ &> 2d \left([m(n-1)]^{\frac{1}{2} + \frac{1}{2k}} + (m+n-1) \right), \end{aligned}$$

so that G' contains a $2k$ -gon by the inductive hypothesis. Similarly, we may assume that the minimum degree in B is at least

$$d_B = d \frac{n^{\frac{1}{2} + \frac{1}{2k}}}{m^{\frac{1}{2} - \frac{1}{2k}}} + d.$$

Choose a vertex $v \in A$ and let H_r be the subgraph of G induced by vertices at distance at most r from v . Let us show that H_r has average degree at least d for some $r \leq k$. Suppose this cannot be done. Let D_r denote the set of vertices of H_r at distance exactly r from v . Then the average number of neighbors in D_{r-1} of a vertex in D_r is less than d . It follows that if $D_r \subset A$, then $|D_{r-1}|(d_B - d) < d|D_r|$ and if $D_r \subset B$, then $|D_{r-1}|(d_A - d) < d|D_r|$. Therefore,

$$|D_r| > \begin{cases} \left(\frac{d_A}{d} - 1 \right) |D_{r-1}| = \frac{m^{\frac{1}{2} + \frac{1}{2k}}}{n^{\frac{1}{2} - \frac{1}{2k}}} |D_{r-1}| & \text{if } r \text{ is odd.} \\ \left(\frac{d_B}{d} - 1 \right) |D_{r-1}| = \frac{n^{\frac{1}{2} + \frac{1}{2k}}}{m^{\frac{1}{2} - \frac{1}{2k}}} |D_{r-1}| & \text{if } r \text{ is even.} \end{cases}$$

Iterating these inequalities for $r = 1, 2, \dots, k$ we get that since k is odd,

$$|D_k| > \left(\frac{m^{\frac{1}{2} + \frac{1}{2k}}}{n^{\frac{1}{2} - \frac{1}{2k}}} \right)^{\lceil \frac{k}{2} \rceil} \left(\frac{n^{\frac{1}{2} + \frac{1}{2k}}}{m^{\frac{1}{2} - \frac{1}{2k}}} \right)^{\lfloor \frac{k}{2} \rfloor} = m.$$

On the other hand, using the fact that k is odd once more, $D_k \subset B$, so that $|D_k| \leq m$, which is a contradiction. The proof of the second part is complete.

Case 2. k is even

The proof here is similar, so we only indicate the necessary changes to the argument. The inductive hypothesis implies that the minimum degree in A is at least:

$$d'_A = d \frac{m^{\frac{1}{2} + \frac{1}{k}}}{\sqrt{n}} + d,$$

and the minimum degree in B is at least:

$$d_B = d \frac{\sqrt{n}}{m^{\frac{1}{2} - \frac{1}{k}}} + d.$$

We now start with a vertex $v \in B$, and repeat the above argument. Since k is even, $D_k \subset B$, so that $|D_k| \leq m$, but

$$|D_k| > \left(\frac{m^{\frac{1}{2} + \frac{1}{k}}}{\sqrt{n}} \right)^{\frac{k}{2}} \left(\frac{\sqrt{n}}{m^{\frac{1}{2} - \frac{1}{k}}} \right)^{\frac{k}{2}} = m,$$

so we once more arrive at a contradiction. ■

3. Proof of Theorem 4

Suppose we are given an m by n bipartite graph H , of girth at least $2k + 2$. From H , we construct a $(k - 1)m$ by n bipartite graph containing no $2k$ -cycles, and with $k - 1$ times as many edges as H . Let A, B be the parts of H , let A_1, A_2, \dots, A_{k-1} be disjoint sets, and let $\phi : \bigcup_{i=1}^{k-1} A_i \rightarrow A$ be defined so that ϕ restricted to A_i is a bijection $A_i \leftrightarrow A$. Define a new graph G with parts $\bigcup_{i=1}^{k-1} A_i$ and B , with edge set

$$E = \{ab : \phi(a)b \in H\}.$$

In words, we are taking $(k - 1)$ identical edge-disjoint copies of H which share B as one of their parts. We now show that G has no $2k$ -cycles.

Suppose, for a contradiction, that G contains a $2k$ -cycle $C = (a_1, b_1, a_2, a_2, \dots, a_k, b_k, a_1)$ with $b_i \in B$ for all $i \in \{1, 2, \dots, k\}$. Then

$$W = (\phi(a_1), b_1, \phi(a_2), b_2, \dots, \phi(a_k), b_k, \phi(a_1))$$

is a closed walk of length $2k$ in H . As H has girth at least $2k + 2$, W takes place on a tree $T \subset H$ with at most k edges. On the other hand, the tree contains the k vertices in $V(C) \cap B$, and there are at least two vertices $a, a' \in V(C) \cap A_i$ for some $i \in \{1, 2, \dots, k-1\}$,

by the pigeonhole principle. Now $\phi(a)$ and $\phi(a')$ are distinct, since ϕ restricted to A_i is a bijection. Therefore the tree has at least $k + 2$ vertices, a contradiction. Therefore the graph G is $2k$ -cycle-free.

References

- [1] de Caen, D.; Székely, L. A. The maximum size of 4- and 6-cycle free bipartite graphs on m, n vertices. *Sets, graphs and numbers* (Budapest, 1991), 135–142, *Colloq. Math. Soc. Jnos Bolyai*, 60, North-Holland, Amsterdam, 1992.
- [2] Erdős, P.; Faudree, R. J.; Rousseau, C. C.; Schelp, R. H. The number of cycle lengths in graphs of given minimum degree and girth. *Paul Erdős memorial collection. Discrete Math.* 200 (1999), no. 1-3, 55–60.
- [3] Erdős, P.; Sárközy, A.; Sós, V. T. On product representations of powers. I. *European J. Combin.* 16 (1995), no. 6, 567–588.
- [4] Györi, E. C_6 -free bipartite graphs and product representation of squares. *Graphs and combinatorics* (Marseille, 1995). *Discrete Math.* 165/166 (1997), 371–375.
- [5] Hoory, S. The size of bipartite graphs with a given girth. *J. Combin. Theory Ser. B* 86 (2002), no. 2, 215–220.
- [6] Lazebnik, F.; Ustimenko, V. A.; Woldar, A. J. Polarities and $2k$ -cycle-free graphs. *16th British Combinatorial Conference* (London, 1997). *Discrete Math.* 197/198 (1999), 503–513.
- [7] Lam, T. Graphs without cycles of even length. *Bull. Austral. Math. Soc.* 63 (2001), no. 3, 435–440.
- [8] Naor, A.; Verstraëte, J. On the Turán number for the hexagon. In preparation.
- [9] Verstraëte, J. A. Arithmetic progressions of cycle lengths in graphs. *Combin. Probab. Comput* 9 (2000) no 4. 369–373.