

**Lecture 14: 4.7 Change of basis.** Recall that if  $\mathcal{B} = \{\mathbf{b}_1, \dots, \mathbf{b}_n\}$  is a basis, then there is a unique way to write any  $\mathbf{x}$  as  $\mathbf{x} = c_1\mathbf{b}_1 + \dots + c_n\mathbf{b}_n$ . The vector

$$[\mathbf{x}]_{\mathcal{B}} = \begin{bmatrix} c_1 \\ \vdots \\ c_n \end{bmatrix}$$

is called the  **$\mathcal{B}$ -coordinate vector of  $\mathbf{x}$** . If  $\mathbf{b}_1, \dots, \mathbf{b}_n$  are vectors in  $\mathbf{R}^n$  then

$$\mathbf{x} = P_{\mathcal{B}}[\mathbf{x}]_{\mathcal{B}}, \quad \text{where} \quad P_{\mathcal{B}} = [\mathbf{b}_1 \ \dots \ \mathbf{b}_n]$$

**Ex** Suppose we have two basis  $\mathcal{B} = \{\mathbf{b}_1, \mathbf{b}_2\}$  and  $\mathcal{C} = \{\mathbf{c}_1, \mathbf{c}_2\}$  that are related by

$$\mathbf{b}_1 = 3\mathbf{c}_1 + 5\mathbf{c}_2, \quad \mathbf{b}_2 = \mathbf{c}_1 + 2\mathbf{c}_2$$

If  $[\mathbf{x}]_{\mathcal{B}} = \begin{bmatrix} 1 \\ -2 \end{bmatrix}$ , i.e.  $\mathbf{x} = \mathbf{b}_1 - 2\mathbf{b}_2$ , find  $[\mathbf{x}]_{\mathcal{C}}$ , i.e.  $(y_1, y_2)$  such that  $\mathbf{x} = y_1\mathbf{c}_1 + y_2\mathbf{c}_2$ .

**Sol**

$$[\mathbf{x}]_{\mathcal{C}} = [\mathbf{b}_1 - 2\mathbf{b}_2]_{\mathcal{C}} = [\mathbf{b}_1]_{\mathcal{C}} - 2[\mathbf{b}_2]_{\mathcal{C}} = \begin{bmatrix} [\mathbf{b}_1]_{\mathcal{C}} & [\mathbf{b}_2]_{\mathcal{C}} \end{bmatrix} \begin{bmatrix} 1 \\ -2 \end{bmatrix} = \begin{bmatrix} 3 & 1 \\ 5 & 2 \end{bmatrix} \begin{bmatrix} 1 \\ -2 \end{bmatrix} = \begin{bmatrix} 1 \\ 1 \end{bmatrix}$$

since  $[\mathbf{b}_1]_{\mathcal{C}} = \begin{bmatrix} 3 \\ 5 \end{bmatrix}$  and  $[\mathbf{b}_2]_{\mathcal{C}} = \begin{bmatrix} 1 \\ 2 \end{bmatrix}$ .

**Th** Let  $\mathcal{B} = \{\mathbf{b}_1, \dots, \mathbf{b}_n\}$  and  $\mathcal{C} = \{\mathbf{c}_1, \dots, \mathbf{c}_n\}$  be two basis for a vector space  $V$ . Then there is a unique matrix  $P_{\mathcal{C} \leftarrow \mathcal{B}}$  such that

$$[\mathbf{x}]_{\mathcal{C}} = P_{\mathcal{C} \leftarrow \mathcal{B}}[\mathbf{x}]_{\mathcal{B}}$$

The columns of  $P_{\mathcal{C} \leftarrow \mathcal{B}}$  are the  $\mathcal{C}$ -coordinate vectors of the vectors in the basis  $\mathcal{B}$ :

$$P_{\mathcal{C} \leftarrow \mathcal{B}} = \begin{bmatrix} [\mathbf{b}_1]_{\mathcal{C}} & \dots & [\mathbf{b}_n]_{\mathcal{C}} \end{bmatrix}$$

The matrix  $P_{\mathcal{C} \leftarrow \mathcal{B}}$  is called the **change-of-coordinates matrix from  $\mathcal{B}$  to  $\mathcal{C}$** . Draw the diagram of  $\mathbf{x} \in V$  and its coordinates in the two basis in  $\mathbf{R}^n$ .

The inverse change of variables must satisfy

$$P_{\mathcal{B} \leftarrow \mathcal{C}} = P_{\mathcal{C} \leftarrow \mathcal{B}}^{-1},$$

since  $P_{\mathcal{B} \leftarrow \mathcal{C}}P_{\mathcal{C} \leftarrow \mathcal{B}}$  is the identity transformation.

If  $\mathcal{C}$  is the standard basis  $\mathcal{E} = \{\mathbf{e}_1, \dots, \mathbf{e}_n\}$  then the change of coordinates

$$P_{\mathcal{E} \leftarrow \mathcal{B}} = P_{\mathcal{B}} = \begin{bmatrix} \mathbf{b}_1 & \dots & \mathbf{b}_n \end{bmatrix}.$$

In general one can write

$$P_{\mathcal{C} \leftarrow \mathcal{B}} = P_{\mathcal{C} \leftarrow \mathcal{E}}P_{\mathcal{E} \leftarrow \mathcal{B}} = P_{\mathcal{E} \leftarrow \mathcal{C}}^{-1}P_{\mathcal{E} \leftarrow \mathcal{B}} = P_{\mathcal{C}}^{-1}P_{\mathcal{B}}$$

In other words we can express a vector in the there different coordinate systems

$$z_1\mathbf{c}_1 + \dots + z_n\mathbf{c}_n = y_1\mathbf{b}_1 + \dots + y_n\mathbf{b}_n = x_1\mathbf{e}_1 + \dots + x_n\mathbf{e}_n$$

and the matrix for the transformation  $[\mathbf{x}]_{\mathcal{B}} = (y_1, \dots, y_n) \rightarrow [\mathbf{x}]_{\mathcal{C}} = (z_1, \dots, z_n)$  can be obtained by first going  $(y_1, \dots, y_n) \rightarrow (x_1, \dots, x_n)$  and then  $(x_1, \dots, x_n) \rightarrow (z_1, \dots, z_n)$ . The matrix for the last transformation is easiest obtained as the inverse of the matrix for  $(z_1, \dots, z_n) \rightarrow (x_1, \dots, x_n)$ .

**Ex** Let  $\mathbf{c}_1 = \begin{bmatrix} 1 \\ 2 \end{bmatrix}$ ,  $\mathbf{c}_2 = \begin{bmatrix} -1 \\ 1 \end{bmatrix}$ , and  $\mathbf{b}_1 = \begin{bmatrix} 3 \\ 0 \end{bmatrix}$ ,  $\mathbf{b}_2 = \begin{bmatrix} 4 \\ 5 \end{bmatrix}$ .

Find the change of coordinate matrix from the coordinates in the  $\mathcal{B} = \{\mathbf{b}_1, \mathbf{b}_2\}$  basis to the coordinates in the  $\mathcal{C} = \{\mathbf{c}_1, \mathbf{c}_2\}$  basis.

**Sol**  $P_{\mathcal{C} \leftarrow \mathcal{B}} = P_{\mathcal{C}}^{-1}P_{\mathcal{B}} = \begin{bmatrix} 1 & -1 \\ 2 & 1 \end{bmatrix}^{-1} \begin{bmatrix} 3 & 4 \\ 0 & 5 \end{bmatrix} = \frac{1}{3} \begin{bmatrix} 1 & 1 \\ -2 & 1 \end{bmatrix} \begin{bmatrix} 3 & 4 \\ 0 & 5 \end{bmatrix} = \begin{bmatrix} 1 & 3 \\ -2 & -1 \end{bmatrix}$

### 5.4 Expressing a linear transformation in terms of different bases.

**Ex** Let  $L$  be the line in  $\mathbf{R}^2$  that is spanned by the vector  $\begin{bmatrix} 3 \\ 1 \end{bmatrix}$ .

Let  $T$  be the linear transformation that projects any vector orthogonally onto  $L$ . Find the matrix  $A$  for  $T$  in the standard coordinate system.

**Sol** We now pick a coordinate system  $\mathcal{B} = \{\mathbf{b}_1, \mathbf{b}_2\}$  with  $\mathbf{b}_1$  parallel to the line and  $\mathbf{b}_2$  perpendicular to the line

$$\mathbf{b}_1 = \begin{bmatrix} 3 \\ 1 \end{bmatrix}, \quad \mathbf{b}_2 = \begin{bmatrix} -1 \\ 3 \end{bmatrix}$$

If  $\mathbf{x} = c_1 \mathbf{b}_1 + c_2 \mathbf{b}_2$  then  $T(\mathbf{x}) = c_1 \mathbf{b}_1$ . Equivalently, if  $[\mathbf{x}]_{\mathcal{B}} = \begin{bmatrix} c_1 \\ c_2 \end{bmatrix}$  then  $[T(\mathbf{x})]_{\mathcal{B}} = \begin{bmatrix} c_1 \\ 0 \end{bmatrix}$ :

$$[T(\mathbf{x})]_{\mathcal{B}} = B [\mathbf{x}]_{\mathcal{B}}, \quad B = \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}$$

The matrix  $B$  for  $T$  in the  $\mathcal{B}$  coordinate system is hence very simple.

The matrix for  $A$  for  $T$  in the standard coordinates is more complicated but one can calculate it from  $B$ :

$$\begin{array}{ccc} \mathbf{x} & \xrightarrow{A} & T(\mathbf{x}) \\ P_{\mathcal{B}} \uparrow & & \uparrow P_{\mathcal{B}} \\ [\mathbf{x}]_{\mathcal{B}} & \xrightarrow{B} & [T(\mathbf{x})]_{\mathcal{B}} \end{array}$$

where  $P_{\mathcal{B}} = [\mathbf{b}_1 \ \mathbf{b}_2] = \begin{bmatrix} 3 & -1 \\ 1 & 3 \end{bmatrix}$  and  $P_{\mathcal{B}}^{-1} = \frac{1}{10} \begin{bmatrix} 3 & 1 \\ -1 & 3 \end{bmatrix}$ . Hence

$$A = P_{\mathcal{B}} B P_{\mathcal{B}}^{-1} = \begin{bmatrix} 3 & -1 \\ 1 & 3 \end{bmatrix} \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix} \frac{1}{10} \begin{bmatrix} 3 & 1 \\ -1 & 3 \end{bmatrix} = \frac{1}{10} \begin{bmatrix} 9 & 3 \\ 3 & 1 \end{bmatrix}$$