

Lecture 5: 3.1 Covariant derivative.

(3.1.1)

$$\nabla_c T^{a_1 \dots a_k}_{b_1 \dots b_\ell} = \partial_c T^{a_1 \dots a_k}_{b_1 \dots b_\ell} + \sum_{i=1}^k \Gamma_{cd}^{a_i} T^{a_1 \dots d \dots a_k}_{b_1 \dots b_\ell} - \sum_{i=1}^{\ell} \Gamma_{cb_i}^d T^{a_1 \dots a_k}_{b_1 \dots d \dots b_\ell}$$

3.2 Curvature.

A calculation shows that

$$(\nabla_a \nabla_b - \nabla_b \nabla_a)(f \omega_c) = f(\nabla_a \nabla_b - \nabla_b \nabla_a) \omega_c$$

It follows that this quantity at a point in the manifold only depends on ω_c at that point and not on the derivatives of ω_c . Hence there is four tensor $R_{abc}{}^d$ such that

$$(3.2.1) \quad (\nabla_a \nabla_b - \nabla_b \nabla_a) \omega_c = R_{abc}{}^d \omega_d.$$

$R_{abc}{}^d$ is called the *Riemann curvature tensor*. A calculations shows that

$$0 = (\nabla_a \nabla_b - \nabla_b \nabla_a)(t^c \omega_c) = \omega_c (\nabla_a \nabla_b - \nabla_b \nabla_a) t^c + t^c (\nabla_a \nabla_b - \nabla_b \nabla_a) \omega_c$$

and hence

$$(3.2.2) \quad (\nabla_a \nabla_b - \nabla_b \nabla_a) t^c = -R_{abd}{}^c t^d.$$

By induction

$$(3.2.3) \quad (\nabla_a \nabla_b - \nabla_b \nabla_a) T^{c_1 \dots c_k}_{d_1 \dots d_\ell} = \sum_{i=1}^{\ell} R_{abd_j}{}^e T^{c_1 \dots c_k}_{d_1 \dots e \dots d_\ell} - \sum_{i=1}^k R_{abe}{}^{e_i} T^{c_1 \dots e \dots c_k}_{d_1 \dots d_\ell}$$

The Riemann curvature tensor satisfy the following properties

1. $R_{abc}{}^d = -R_{bac}{}^d$. This follows directly from (3.2.1).
2. $R_{[abc]}{}^d = 0$. Here $[abc]$ is stands for the antisymmetrization over the indices a, b, c :
 $R_{[abc]}{}^d = R_{abc}{}^d + R_{bca}{}^d + R_{cab}{}^d - R_{bac}{}^d - R_{bca}{}^d - R_{acb}{}^d$ so if we also use (1) we get:
 $R_{abc}{}^d + R_{bca}{}^d + R_{cab}{}^d = 0$.
3. $R_{abcd} = -R_{abdc}$, if $R_{abcd} = R_{abc}{}^e g_{de}$. This follows from (3.2.2) since $\nabla_a g_{bc} = 0$.
 Moreover (1)-(3) in addition implies $R_{abcd} = R_{cdab}$.
4. The Bianchi identity $\nabla_{[a} R_{bc]d}{}^e = 0$.

Using (3.1.1) and $\Gamma_{ab}^c = \Gamma_{ba}^c$ one can show that $\nabla_{[a} \nabla_b \omega_c] = 0$, from which (2) follows.
 (4) follows from using (3.2.14) and (2).