

# ON THE PRIME FACTORS OF $\binom{n}{k}$

P. ERDÖS

Hungarian Academy of Sciences, Budapest, Hungary  
and

R. L. GRAHAM

Bell Laboratories, Murray Hill, New Jersey

A well known theorem of Sylvester and Schur (see [5]) states that for  $n \geq 2k$ , the binomial coefficient  $\binom{n}{k}$  always has a prime factor exceeding  $k$ . This can be considered as a generalization of the theorem of Chebyshev: There is always a prime between  $m$  and  $2m$ . Set

$$\binom{n}{k} = u_n(k)v_n(k)$$

with

$$u_n(k) = \prod_{p < k} p^{\alpha_{\parallel} \binom{n}{k}}, \quad v_n(k) = \prod_{p \geq k} p^{\alpha_{\parallel} \binom{n}{k}}.$$

In [4] it is proved that  $v_n(k) > u_n(k)$  for all but a finite number of cases (which are tabulated there).

In this note, we continue the investigation of  $u_n(k)$  and  $v_n(k)$ . We first consider  $v_n(k)$ , the product of the large prime divisors of  $\binom{n}{k}$ .

*Theorem.*

$$\max_{1 < k \leq n} v_n(k) = e^{\frac{n}{2}(1+o(1))}.$$

*Proof.* For  $k < \epsilon n$  the result is immediate since in this case  $\binom{n}{k}$  itself is less than  $e^{n/2}$ . Also, it is clear that the maximum of  $v_n(k)$  is not achieved for  $k > n/2$ . Hence, we may assume  $\epsilon n < k \leq n/2$ . Now, for any prime

$$p \in \left( \frac{n-k}{r}, \frac{n}{r} \right]$$

with  $p \geq k$  and  $r \geq 1$ , we have  $p | v_n(k)$ . Also, if  $k^2 > n$  then  $p^2 \nmid v_n(k)$  so that in this case the contribution to  $v_n(k)$  of the primes

$$p \in \left( \frac{n-k}{r}, \frac{n}{r} \right]$$

is (by the Prime Number Theorem (PNT)) just  $e^{\frac{k}{r}(1+o(1))}$ . Thus, letting  $\frac{n}{t+1} < k \leq \frac{n}{t}$ , we obtain

$$v_n(k) = \exp \left[ \left( \sum_{r=1}^{t-1} \frac{k}{r} + \left( \frac{n}{t} - k \right) \right) (1+o(1)) \right] = \exp \left[ \left( \frac{n}{t} \sum_{r=1}^{t-1} \frac{1}{r} \right) (1+o(1)) \right] \\ \leq e^{\frac{n}{2}(1+o(1))}$$

and the theorem is proved.

It is interesting to note that since

$$\frac{n}{t} \sum_{r=1}^{t-1} \frac{1}{r} = \frac{1}{2}$$

for both  $t = 2$  and  $t = 3$  then

$$\lim_n v_n(k)^{1/n} = e^{1/2}$$

for any  $k \in \left(\frac{n}{3}, \frac{n}{2}\right)$ .

In Table 1, we tabulate the least value  $k^*(n)$  of  $k$  for which  $v_n(k)$  achieves its maximum value for selected values of  $n \leq 200$ . It seems likely that infinitely often  $k^*(n) = \frac{n}{2}$  but we are at present far from being able to prove this.

Table 1

$n$	$k^*(n)$	$n$	$k^*(n)$	$n$	$k^*(n)$
2	1	10	2	18	8
3	1	11	3	19	9
4	2	12	6	20	10
5	2	13	4	50	22
6	2	14	4	100	42
7	3	15	5	200	100
8	4	16	6		
9	2	17	7		

Note that

$$v_7(0) < v_7(1) < v_7(2) < v_7(3).$$

It is easy to see that for  $n > 7$ , the  $v_n(k)$  cannot increase monotonically for  $0 \leq k \leq \frac{n}{2}$ .

Next, we mention several results concerning  $u_n(k)$ . To begin with, note that while  $u_n(k) = 1$  for  $0 \leq k \leq \frac{n}{2}$ , this behavior is no longer possible for  $n > 7$ . In fact, we have the following more precise statement.

**Theorem.** For some  $k \leq (2 + o(1)) \log n$ , we have  $u_n(k) > 1$ .

*Proof.* Suppose  $u_n(k) = 1$  for all  $k \leq (2 + \epsilon) \log n$ . Choose a prime  $p < (1 + \epsilon) \log n$  which does not divide  $n + 1$ . Such a prime clearly exists (for large  $n$ ) by the PNT. Since  $p \nmid n + 1$  then for some  $k$  with  $p < k < 2p$ ,

$$p^2 \mid n(n-1) \dots (n-k+1), \quad p^2 \nmid k!$$

Thus,  $p \mid u_n(k)$  and since

$$k < 2p < (2 + 2\epsilon) \log n,$$

the theorem is proved.

In the other direction we have the following result.

**Fact.** There exist infinitely many  $n$  so that for all  $k \leq (1/2 + o(1)) \log n$ ,  $u_n(k) = 1$ .

*Proof.* Choose  $n + 1 = [\text{l.c.m. } \{1, 2, \dots, t\}]^2$ . By the PNT,  $n = e^{(2+o(1))t}$ . Clearly, if  $m \leq t$  then  $m \nmid \binom{n}{t}$ . Thus,

$$u_n(k) = 1 \quad \text{for } k \leq \left(\frac{1}{2} + o(1)\right) \log n$$

as claimed.

In Table 2 we list the least value  $n^*(k)$  of  $n$  such that  $u_n(i) = 1$  for  $1 \leq i \leq k$

Table 2

$k$	$n^*(k)$
1	1
2	2
3	3
4	7
5	23
6	71

Of course, for  $k \leq 2$ ,  $u_n(k) = 1$  is automatic. By a theorem of Mahler [11], it follows that

$$u_n(k) < n^{1+\epsilon}$$

for  $k \geq 3$  and large  $n$ . It is well known that if  $p^\alpha \mid \binom{n}{k}$  then  $p^\alpha \leq n$ . Consequently,

$$u_n(k) \leq n^{\pi(k)},$$

where  $\pi(k)$  denotes the number of primes not exceeding  $k$ . It seems likely that the following stronger estimate holds:

$$(*) \quad u_n(k) < n^{(1+o(1))(1-\gamma)\pi(k)}, \quad k \geq 5,$$

where  $\gamma$  denotes Euler's constant. It is easy to prove (\*) for certain ranges of  $k$ . For example, suppose  $k$  is relatively large compared to  $n$ , say,  $k = n/t$  for a large fixed  $t$ . Of course, any prime  $p \in (n - n/t, n)$  divides  $v_n(k)$  and by the PNT

$$\prod_{n(1-1/t) < p < n} p = e^{(1+o(1))n/t}.$$

More generally, if  $rp \in (n - n/t, n)$  with  $r < t$  then  $p \geq k$  and  $p \mid v_n(k)$  so that again by the PNT

$$\prod_{\frac{n}{r} \left(1 - \frac{1}{t}\right) < p < \frac{n}{r}} p = e^{(1+o(1))n/rt}.$$

Thus

$$\begin{aligned} v_n(k) &\geq \prod_{1 \leq r < t} \prod_{\frac{n}{t} \left(1 - \frac{1}{t}\right) < p < \frac{n}{r}} p = \exp \left( (1+o(1)) \sum_{1 \leq r < t} \frac{1}{r} \right) \frac{n}{t} \\ &= \exp \left( (1+o(1))(\log t + \gamma) \right) \frac{n}{t}. \end{aligned}$$

But by Stirling's formula we have

$$\binom{n}{n/t} = e^{\frac{n}{t} \log t + \frac{n}{t} + o\left(\frac{n}{t}\right)}$$

Thus,

$$\begin{aligned} u_n(k) = \binom{n}{k} / v_n(k) &\leq e^{\frac{n}{t} \log t + \frac{n}{t} + o\left(\frac{n}{t}\right) - (1+o(1))(\log t + \gamma) \frac{n}{t}} \\ &= e^{(1+o(1))(1-\gamma) \frac{n}{t}} = n^{(1+o(1))(1-\gamma)\pi(k)} \end{aligned}$$

which is just (\*).

In contrast to the situation for  $v_n(k)$ , the maximum value of  $u_n(k)$  clearly occurs for  $k \geq \frac{n}{2}$ . Specifically, we have the following result.

**Theorem.** The value  $\hat{k}(n)$  of  $k$  for which  $u_n(k)$  assumes its maximum value satisfies

$$\hat{k}(n) = (1+o(1)) \left( \frac{e}{e+1} \right) n.$$

**Proof.** Let  $k = (1-c)n$ . For  $c \leq \frac{1}{2}$ ,

$$v_n(k) = \prod_{n-k < p \leq n} p = e^{(1+o(1))cn}.$$

Since

$$\binom{n}{k} = \binom{n}{cn} = e^{-c \log c + (1-c) \log(1-c)} (1+o(1))n$$

then

$$u_n(k) = \binom{n}{k} / v_n(k) = e^{-(1+o(1))(c + \log c^c (1-c)^{1-c})n}.$$

A simple calculation shows that the exponent is maximized by taking  $c = \frac{1}{e+1} = 0.2689 \dots$

**Concluding remarks.** We mention here several related problems which were not able to settle or did not have time to investigate. One of the authors [8] previously conjectured that  $\binom{2n}{n}$  is never squarefree for  $n > 4$  (at present this is still open). Of course, more generally, we expect that for all  $a$ ,  $\binom{2n}{n}$  is always divisible by an  $a^{\text{th}}$  power of a prime  $> k$  if  $n > n_0(a, k)$ . We can show the much weaker result that  $n = 23$  is the largest value of  $n$  for which all  $\binom{n}{k}$  are squarefree for  $0 < k < n$ . This follows from the observation that if  $p$  is prime and  $p^\alpha \nmid \binom{n}{k}$  for any  $k$  then  $p^\beta \mid n + 1$ , where

$$p^\beta \geq \frac{n+1}{p^\alpha - 1}.$$

Thus,  $2^2 \nmid \binom{n}{k}$  for any  $k$  implies  $2^\beta \mid n + 1$  where  $2^\beta \geq \frac{n+1}{3}$ . Also,  $3^2 \nmid \binom{n}{k}$  for any  $k$  implies  $3^\gamma \mid n + 1$  where  $3^\gamma \geq \frac{n+1}{8}$ . Together these imply that  $d = 2^\beta 3^\gamma \mid n + 1$  where  $d \geq (n+1)^2/24$ . Since  $d$  cannot exceed  $n+1$  then  $n+1 \leq 24$  is forced, and the desired result follows.

For given  $n$  let  $f(n)$  denote the largest integer such that for some  $k$ ,  $\binom{n}{k}$  is divisible by the  $f(n)^{\text{th}}$  power of a prime. We can prove that  $f(n) \rightarrow \infty$  as  $n \rightarrow \infty$  (this is not hard) and very likely  $f(n) > c \log n$  but we are very far from being able to prove this. Similarly, if  $F(n)$  denotes the largest integer so that for all  $k$ ,  $1 \leq k < n$ ,  $\binom{n}{k}$  is divisible by the  $F(n)^{\text{th}}$  power of some prime, then it is quite likely that  $\overline{\lim} F(n) = \infty$ , but we have not proved this.

Let  $P(x)$  and  $p(x)$  denote the greatest and least prime factors of  $x$ , respectively. Probably

$$p\left(\binom{n}{k}\right) > \max(n-k, k^{1+\epsilon})$$

but this seems very deep (for related results see the papers of Ramachandra and others [11], [12]).

J. L. Selfridge and P. Erdős conjectured and Ecklund [1] proved that  $p\left(\binom{n}{k}\right) < \frac{n}{2}$  for  $k > 1$ , with the unique exception of  $p\left(\binom{7}{3}\right) = 5$ . Selfridge and Erdős [9] proved that

$$p\left(\binom{n}{k}\right) < \frac{c \cdot n}{k^{c_2}}$$

and they conjecture

$$p\left(\binom{n}{k}\right) < \frac{n}{k} \text{ for } n > k^2.$$

Finally, let  $d\left(\binom{n}{k}\right)$  denote the greatest divisor of  $\binom{n}{k}$  not exceeding  $n$ . Erdős originally conjectured that  $d\left(\binom{n}{k}\right) > n - k$  but this was disproved by Schinzel and Erdős [13]. Perhaps it is true however, that  $d_n > cn$  for a suitable constant  $c$ .

For problems and results of a similar nature the reader may consult [2], [3], [6], [7] or [10].

#### REFERENCES

1. E. Ecklund, Jr., "On Prime Divisors of the Binomial Coefficients," *Pac. J. of Math.*, 29 (1969), pp. 267-270.
2. E. Ecklund, Jr., and R. Eggleton, "Prime Factors of Consecutive Integers," *Amer. Math. Monthly*, 79 (1972) pp. 1082-1089.
3. E. Ecklund, Jr., P. Erdős and J. L. Selfridge, "A New Function Associated with the Prime Factors of  $\binom{n}{k}$ ," *Math. Comp.*, 28 (1974), pp. 647-649.
4. E. Ecklund, Jr., R. Eggleton, P. Erdős and J. L. Selfridge, "On the Prime Factorization of Binomial Coefficients," to appear.
5. P. Erdős, "On a Theorem of Sylvester and Schur," *J. London Math. Soc.*, 9 (1934), pp. 282-288.
6. P. Erdős, "Some Problems in Number Theory," *Computers in Number Theory, Proc. Atlas Symp.*, Oxford, 1969, Acad. Press (1971), pp. 405-414.

7. P. Erdős, "Über die Anzahl der Primzahlen von  $\binom{n}{k}$ ," *Archiv der Math.*, 24 (1973).
8. P. Erdős, "Problems and Results on Number Theoretic Properties of Consecutive Integers and Related Questions," *Proc. Fifth Manitoba Conf. on Numerical Mathematics* (1975), pp. 25–44.
9. P. Erdős and J. L. Selfridge, "Some Problems on the Prime Factors of Consecutive Integers," *Pullman Conf. on Number Theory*, (1971).
10. P. Erdős, R. L. Graham, I. Z. Ruzsa and E. G. Straus, "On the Prime Factors of  $\binom{2n}{n}$ ," *Math Comp.*, 29 (1975), pp. 83–92.
11. K. Ramachandra, "A Note on Numbers with a Large Prime Factor II," *J. Indian Math. Soc.*, 34 (1970), pp. 39–48.
12. K. Ramachandra, "A Note on Numbers with a Large Prime Factor III," *Acta Arith.*, 19 (1971), pp. 49–62.
13. A. Schinzel, "Sur un Probleme de P. Erdős," *Coll. Math.*, 5 (1952), pp. 198–204.

★★★★★