

Representations of Metaplectic Groups

Wee Teck Gan
University of California, San Diego
National University of Singapore

December 18, 2010

Local Langlands Correspondence (LLC)

Let F be a local field with Weil group

$$W_F \subset \text{Gal}(\overline{F}/F),$$

and Weil-Deligne group

$$WD_F = W_F \times \text{SL}_2(\mathbb{C})$$

Local Langlands Correspondence (LLC)

Let F be a local field with Weil group

$$W_F \subset \text{Gal}(\overline{F}/F),$$

and Weil-Deligne group

$$WD_F = W_F \times \text{SL}_2(\mathbb{C})$$

Let G be a connected reductive linear algebraic group over F .

Local Langlands Correspondence (LLC)

Let F be a local field with Weil group

$$W_F \subset \text{Gal}(\overline{F}/F),$$

and Weil-Deligne group

$$WD_F = W_F \times \text{SL}_2(\mathbb{C})$$

Let G be a connected reductive linear algebraic group over F .

The LLC for G gives a classification of

$$\text{Irr}(G) = \{\text{isom. classes of irreducible representations of } G(F)\}$$

in terms of arithmetic data.

L-parameters

Set

$$\Phi(G) = \{\text{isom. classes of pairs } (\phi, \eta)\}$$

L-parameters

Set

$$\Phi(G) = \{\text{isom. classes of pairs } (\phi, \eta)\}$$

with

$$\phi : WD_F \longrightarrow {}^L G$$

and

$$\eta \in \text{Irr}(A_\phi),$$

where

$$A_\phi = \pi_0(Z_{L G^0}(\phi)).$$

L-parameters

Set

$$\Phi(G) = \{\text{isom. classes of pairs } (\phi, \eta)\}$$

with

$$\phi : WD_F \longrightarrow {}^L G$$

and

$$\eta \in \text{Irr}(A_\phi),$$

where

$$A_\phi = \pi_0(Z_{L G^0}(\phi)).$$

Then LLC postulates a bijection

$$\mathcal{L} : \bigsqcup_{G'} \text{Irr}(G') \longleftrightarrow \Phi(G)$$

as G' ranges over pure inner forms of G .

Characterization

The bijection \mathcal{L} could be characterized in two ways:

Characterization

The bijection \mathcal{L} could be characterized in two ways:

- requiring it to preserve certain natural invariants, such as L -factors and ϵ factors;

Characterization

The bijection \mathcal{L} could be characterized in two ways:

- requiring it to preserve certain natural invariants, such as L -factors and ϵ factors;
- by stability properties of the characters of representations, and character identities with certain smaller groups ([theory of endoscopy](#)).

Characterization

The bijection \mathcal{L} could be characterized in two ways:

- requiring it to preserve certain natural invariants, such as L -factors and ϵ factors;
- by stability properties of the characters of representations, and character identities with certain smaller groups ([theory of endoscopy](#)).

The LLC is, on one hand, a generalization of [local class field theory](#), and on the other hand, a generalization of the [Cartan-Weyl theory of highest weights](#) for representations of connected compact Lie groups.

The LLC is known for

- $GL(n)$: Harris-Taylor, Henniart;

The LLC is known for

- $GL(n)$: Harris-Taylor, Henniart;
- $SL(n)$ and its inner forms: Gelbart-Knapp, Hiraga-Saito;

The LLC is known for

- $GL(n)$: Harris-Taylor, Henniart;
- $SL(n)$ and its inner forms: Gelbart-Knapp, Hiraga-Saito;
- $U(3)$ and its inner forms: Rogawski;
- $GSp(4)$ and its inner form, $Sp(4)$: joint with Takeda and Tantonio;

The LLC is known for

- $GL(n)$: Harris-Taylor, Henniart;
- $SL(n)$ and its inner forms: Gelbart-Knapp, Hiraga-Saito;
- $U(3)$ and its inner forms: Rogawski;
- $GSp(4)$ and its inner form, $Sp(4)$: joint with Takeda and Tanton;
- all classical groups $SO(n)$, $Sp(2n)$ and $U(n)$: pending book of Arthur, and work of Moeglin.

Thus, I will take the view that the LLC is essentially known.

Nonlinear Groups?

Can one extend the Langlands philosophy to the case of non-linear covering groups of $G(F)$?

Nonlinear Groups?

Can one extend the Langlands philosophy to the case of non-linear covering groups of $G(F)$?

There have been many attempts towards a partial extension, e.g.:

- Real case: Adams, Barbasch, Vogan, Renard, Herb,.....

Nonlinear Groups?

Can one extend the Langlands philosophy to the case of non-linear covering groups of $G(F)$?

There have been many attempts towards a partial extension, e.g.:

- Real case: Adams, Barbasch, Vogan, Renard, Herb,.....
- p -adic case: Savin, Kable, Weissman,

Nonlinear Groups?

Can one extend the Langlands philosophy to the case of non-linear covering groups of $G(F)$?

There have been many attempts towards a partial extension, e.g.:

- Real case: Adams, Barbasch, Vogan, Renard, Herb,.....
- p -adic case: Savin, Kable, Weissman,
- Global case: Waldspurger, Flicker, Kazhdan, Mezo, Ginzburg-Jiang-Rallis-Soudry,.....

Nonlinear Groups?

Can one extend the Langlands philosophy to the case of non-linear covering groups of $G(F)$?

There have been many attempts towards a partial extension, e.g.:

- Real case: Adams, Barbasch, Vogan, Renard, Herb,.....
- p -adic case: Savin, Kable, Weissman,
- Global case: Waldspurger, Flicker, Kazhdan, Mezo, Ginzburg-Jiang-Rallis-Soudry,.....

The time is perhaps ripe for a systematic extension of the Langlands program (local and global) to general covering groups. Today, we will look at a particular family of such groups: the **metaplectic groups**.

Shimura Correspondence

Cuspidal Hecke eigenform
of weight $(2k + 1)/2$
and level $\Gamma_0(4N)$



Cuspidal Hecke eigenform
of weight $2k$
and level $\Gamma_0(N)$

Shimura Correspondence

Cuspidal Hecke eigenform
of weight $(2k + 1)/2$
and level $\Gamma_0(4N)$



Cuspidal Hecke eigenform
of weight $2k$
and level $\Gamma_0(N)$

Established using converse theorem for classical modular forms.
Alternative approach via **Theta Correspondence** was pursued by
Niwa and Shintani.

Waldspurger's Results

Extending this, Waldspurger obtained a definitive treatment in the framework of automorphic representations.

Waldspurger's Results

Extending this, Waldspurger obtained a definitive treatment in the framework of automorphic representations.

Cuspidal Representations of $\widetilde{\mathrm{SL}}_2(\mathbb{A})$

↓ finite-to-one

Cuspidal Representations of $\mathrm{PGL}_2(\mathbb{A})$

Waldspurger's Results

Extending this, Waldspurger obtained a definitive treatment in the framework of automorphic representations.

Cuspidal Representations of $\widetilde{SL}_2(\mathbb{A})$

↓ finite-to-one

Cuspidal Representations of $PGL_2(\mathbb{A})$

Here, $\widetilde{SL}_2(\mathbb{A})$ denotes the double cover of $SL_2(\mathbb{A}) = Sp_2(\mathbb{A})$. The map depends on an additive character

$$\psi : F \backslash \mathbb{A} \rightarrow \mathbb{C}^\times.$$

Waldspurger's Local Results

The above result depends on a local analog:

Irreducible Genuine Representations
of $\widetilde{\mathrm{SL}}_2(F)$

↓ fibers of size 1 or 2

Irreducible Representations of $\mathrm{PGL}_2(F)$

Waldspurger's Local Results

The above result depends on a local analog:

Irreducible Genuine Representations
of $\widetilde{\mathrm{SL}}_2(F)$

↓ fibers of size 1 or 2

Irreducible Representations of $\mathrm{PGL}_2(F)$

Here an irreducible *genuine* representation of $\widetilde{\mathrm{SL}}_2(F)$ is one which does not factor to a representation of $\mathrm{SL}_2(F)$. The map depends on an additive character


$$\psi : F \rightarrow \mathbb{C}^\times$$

Refinement

If D denotes the quaternion division F -algebra, then PD^\times is a pure inner form of $\mathrm{PGL}_2(F)$ and the above map can be refined to a bijection:

Refinement

If D denotes the quaternion division F -algebra, then PD^\times is a pure inner form of $\mathrm{PGL}_2(F)$ and the above map can be refined to a bijection:



Remarks:

(i) The first (2-to-1) local map is simply the map Θ_ψ followed by the Jacquet-Langlands correspondence.

Remarks:

- (i) The first (2-to-1) local map is simply the map Θ_ψ followed by the Jacquet-Langlands correspondence.
- (ii) Observe that

$$\mathrm{PGL}_2 = \mathrm{SO}(V^+), \quad PD^\times = \mathrm{SO}(V^-)$$

where V^+ (resp. V^-) denotes the 3-dim split (resp. non-split) quadratic space of discriminant 1.

Remarks:

- (i) The first (2-to-1) local map is simply the map Θ_ψ followed by the Jacquet-Langlands correspondence.
- (ii) Observe that

$$\mathrm{PGL}_2 = \mathrm{SO}(V^+), \quad \mathrm{PD}^\times = \mathrm{SO}(V^-)$$

where V^+ (resp. V^-) denotes the 3-dim split (resp. non-split) quadratic space of discriminant 1.

So the map Θ_ψ is a classification of the genuine irreps of the double cover of the symplectic group of rank 1 by the irreps of the two special orthogonal groups of rank 1.

(iii) Coupled with the LLC, one has:

Genuine Irreps of $\widetilde{\mathrm{SL}}_2(F)$

$\updownarrow \mathcal{L}_\psi$

pairs $(\phi, \eta) : \eta \in \mathrm{Irr}(A_\phi)$

where

$$\phi : WD_F \rightarrow \mathrm{SL}_2(\mathbb{C}),$$

and

$$A_\phi = \pi_0(Z_{\mathrm{SL}_2(\mathbb{C})}(\phi)) = 1 \text{ or } \mathbb{Z}/2\mathbb{Z}.$$

Goal of this talk

- Extend the above results of Waldspurger to higher rank symplectic groups.
(Joint work with Savin, Ichino and Gross-Prasad.)

Goal of this talk

- Extend the above results of Waldspurger to higher rank symplectic groups.
(Joint work with Savin, Ichino and Gross-Prasad.)
- Highlight some other recent developments, especially with regards to a theory of endoscopy. (Work of Wenwei Li)

Notations: Metaplectic Groups

- $W = W_n =$ a $2n$ -dimensional symplectic vector space over a local field F ;
- $\mathrm{Sp}(W) =$ the associated symplectic group

Notations: Metaplectic Groups

- $W = W_n =$ a $2n$ -dimensional symplectic vector space over a local field F ;
- $\mathrm{Sp}(W)$ = the associated symplectic group
- $\mathrm{Mp}(W)$ = the unique 2-fold cover of $\mathrm{Sp}(W)$:

$$1 \longrightarrow \mu_2 \longrightarrow \mathrm{Mp}(W) \longrightarrow \mathrm{Sp}(W) \longrightarrow 1$$

Notations: Metaplectic Groups

- $W = W_n =$ a $2n$ -dimensional symplectic vector space over a local field F ;
- $\mathrm{Sp}(W) =$ the associated symplectic group
- $\mathrm{Mp}(W) =$ the unique 2-fold cover of $\mathrm{Sp}(W)$:

$$1 \longrightarrow \mu_2 \longrightarrow \mathrm{Mp}(W) \longrightarrow \mathrm{Sp}(W) \longrightarrow 1$$

- If $H \subset \mathrm{Sp}(W)$, then \tilde{H} is the preimage of H in $\mathrm{Mp}(W)$
- If $Z \cong \mu_2 =$ the center of $\mathrm{Sp}(W)$, then \tilde{Z} is the center of $\mathrm{Mp}(W)$.

Notations: Orthogonal Groups



V^+ = split quadratic space of
dim $2n + 1$ and disc. 1

Notations: Orthogonal Groups



V^+ = split quadratic space of
dim $2n + 1$ and disc. 1

V^- = non-split quadratic space of
dim $2n + 1$ and disc. 1

Notations: Orthogonal Groups



V^+ = split quadratic space of
dim $2n + 1$ and disc. 1

V^- = non-split quadratic space of
dim $2n + 1$ and disc. 1

- Set $\epsilon(V^\pm) = \pm$.
- Have associated orthogonal groups

$$O(V^\epsilon) = SO(V^\epsilon) \times \{\pm 1\}$$

Notations: Orthogonal Groups



V^+ = split quadratic space of
dim $2n + 1$ and disc. 1

V^- = non-split quadratic space of
dim $2n + 1$ and disc. 1

- Set $\epsilon(V^\pm) = \pm$.
- Have associated orthogonal groups

$$O(V^\epsilon) = SO(V^\epsilon) \times \{\pm 1\}$$

- An irred rep π of $SO(V)$ has 2 extensions π^+ and π^- to $O(V)$. The sign in π^ϵ simply encodes the central character:

$$\epsilon = \pi^\epsilon(-1).$$

Theorem A

Assume that F is p -adic with $p \neq 2$. Fix an additive character $\psi : F \rightarrow \mathbb{C}^\times$. Then there is a bijection

$$\begin{array}{c} \text{Irr}(\text{Mp}(W)) \\ \updownarrow \Theta_\psi \\ \text{Irr}(\text{SO}(V^+)) \sqcup \text{Irr}(\text{SO}(V^-)) \end{array}$$

Theorem A

Assume that F is p -adic with $p \neq 2$. Fix an additive character $\psi : F \rightarrow \mathbb{C}^\times$. Then there is a bijection

$$\begin{array}{c} \text{Irr}(\text{Mp}(W)) \\ \updownarrow \Theta_\psi \\ \text{Irr}(\text{SO}(V^+)) \sqcup \text{Irr}(\text{SO}(V^-)) \end{array}$$

Remark: when $F = \mathbb{R}$, the analogous thm was shown by Adams-Barbasch.

Theorem A

Assume that F is p -adic with $p \neq 2$. Fix an additive character $\psi : F \rightarrow \mathbb{C}^\times$. Then there is a bijection

$$\begin{array}{c} \text{Irr}(\text{Mp}(W)) \\ \updownarrow \Theta_\psi \\ \text{Irr}(\text{SO}(V^+)) \sqcup \text{Irr}(\text{SO}(V^-)) \end{array}$$

Remark: when $F = \mathbb{R}$, the analogous thm was shown by Adams-Barbasch.

Corollary B (LLC): If the LLC for $\text{SO}(V^\pm)$ holds (e.g. for $n = 1$ or 2), one has a bijection

$$\begin{array}{c} \text{Irr}(\text{Mp}(W)) \\ \updownarrow \mathcal{L}_\psi \\ \{(\phi, \eta) : \eta \in \text{Irr}(A_\phi)\} \end{array}$$

Sketch Proof of Thm A

The map Θ_ψ is defined using **theta correspondence**.

Sketch Proof of Thm A

The map Θ_ψ is defined using **theta correspondence**.

- the group

$$\mathrm{Mp}(W) \times \mathrm{O}(V)$$

has a natural representation $\Omega_{V,W,\psi}$ (the Weil rep).

Sketch Proof of Thm A

The map Θ_ψ is defined using **theta correspondence**.

- the group

$$\mathrm{Mp}(W) \times \mathrm{O}(V)$$

has a natural representation $\Omega_{V,W,\psi}$ (the Weil rep).

- for an irrep σ of $\mathrm{Mp}(W)$, the maximal σ -isotypic quotient of Ω_ψ has the form

$$\sigma \boxtimes \Theta_{V,W,\psi}(\sigma),$$

for some smooth representation $\Theta_{V,W,\psi}(\sigma)$ of $\mathrm{O}(V)$ (the *big* theta lift of σ).

Sketch Proof of Thm A

The map Θ_ψ is defined using **theta correspondence**.

- the group

$$\mathrm{Mp}(W) \times \mathrm{O}(V)$$

has a natural representation $\Omega_{V,W,\psi}$ (the Weil rep).

- for an irrep σ of $\mathrm{Mp}(W)$, the maximal σ -isotypic quotient of Ω_ψ has the form

$$\sigma \boxtimes \Theta_{V,W,\psi}(\sigma),$$

for some smooth representation $\Theta_{V,W,\psi}(\sigma)$ of $\mathrm{O}(V)$ (the *big* theta lift of σ).

- (Kudla) $\Theta_{V,W,\psi}(\sigma)$ is zero or of finite length.

Facts about theta correspondence

Let $\theta_{V,W,\psi}$ (the *small* theta lift) be the maximal semisimple quotient of $\Theta_{V,W,\psi}(\sigma)$.

Facts about theta correspondence

Let $\theta_{V,W,\psi}$ (the *small* theta lift) be the maximal semisimple quotient of $\Theta_{V,W,\psi}(\sigma)$. Then:

- (Howe, Waldspurger) If $p \neq 2$, then $\theta_{V,W,\psi}(\sigma)$ is zero or irreducible.

Facts about theta correspondence

Let $\theta_{V,W,\psi}$ (the *small* theta lift) be the maximal semisimple quotient of $\Theta_{V,W,\psi}(\sigma)$. Then:

- (Howe, Waldspurger) If $p \neq 2$, then $\theta_{V,W,\psi}(\sigma)$ is zero or irreducible.
- the above discussion holds with the roles of V and W switched.

Key Proposition

Thm A follows from:

(1) For $\sigma \in \text{Irr}(\text{Mp}(W))$, exactly one of

$$\Theta_{V^+, W, \psi}(\sigma) \quad \text{or} \quad \Theta_{V^-, W, \psi}(\sigma)$$

is nonzero.

The "at most one" part is a difficult theorem of Kudla-Rallis.

Key Proposition

Thm A follows from:

(1) For $\sigma \in \text{Irr}(\text{Mp}(W))$, exactly one of

$$\Theta_{V^+, W, \psi}(\sigma) \quad \text{or} \quad \Theta_{V^-, W, \psi}(\sigma)$$

is nonzero.

The "at most one" part is a difficult theorem of Kudla-Rallis.

(2) For $\pi \in \text{Irr}(\text{SO}(V))$, exactly one extension π^ϵ of π satisfies

$$\Theta_{V, W, \psi}(\pi^\epsilon) \neq 0.$$

Part (1) was also shown by C. Zorn.

Hence, if $p \neq 2$, one defines Θ_ψ by the composite:

$$\begin{array}{ccc} \text{Irr}(\text{Mp}(W)) & \xrightarrow{\theta_{V,W,\psi}} & \text{Irr}(\text{O}(V^+)) \sqcup \text{Irr}(\text{O}(V^-)) \\ & & \downarrow \text{rest} \\ & & \text{Irr}(\text{SO}(V^+)) \sqcup \text{Irr}(\text{SO}(V^-)) \end{array}$$

Hence, if $p \neq 2$, one defines Θ_ψ by the composite:

$$\begin{array}{ccc} \text{Irr}(\text{Mp}(W)) & \xrightarrow{\theta_{V,W,\psi}} & \text{Irr}(\text{O}(V^+)) \sqcup \text{Irr}(\text{O}(V^-)) \\ & & \downarrow \text{rest} \\ & & \text{Irr}(\text{SO}(V^+)) \sqcup \text{Irr}(\text{SO}(V^-)) \end{array}$$

Key Prop implies that the composite is bijective, proving Thm A.

Refinement of Key Prop

- for $\sigma \in \text{Irr}(\text{Mp}(W))$, which $\Theta_{V^\epsilon, W, \psi}(\sigma)$ is nonzero?
- for $\pi \in \text{Irr}(\text{SO}(V))$, which extension π^ϵ has nonzero theta lift to $\text{Mp}(W)$?

Refinement of Key Prop

- for $\sigma \in \text{Irr}(\text{Mp}(W))$, which $\Theta_{V^\epsilon, W, \psi}(\sigma)$ is nonzero?
- for $\pi \in \text{Irr}(\text{SO}(V))$, which extension π^ϵ has nonzero theta lift to $\text{Mp}(W)$?

Theorem C (**Epsilon Dichotomy**):

(i) Given $\pi \in \text{Irr}(\text{SO}(V))$, π^ϵ has nonzero theta lift to $\text{Mp}(W)$ if and only if

$$\epsilon = \epsilon(V) \cdot \epsilon(1/2, \pi)$$

Refinement of Key Prop

- for $\sigma \in \text{Irr}(\text{Mp}(W))$, which $\Theta_{V^\epsilon, W, \psi}(\sigma)$ is nonzero?
- for $\pi \in \text{Irr}(\text{SO}(V))$, which extension π^ϵ has nonzero theta lift to $\text{Mp}(W)$?

Theorem C (**Epsilon Dichotomy**):

(i) Given $\pi \in \text{Irr}(\text{SO}(V))$, π^ϵ has nonzero theta lift to $\text{Mp}(W)$ if and only if

$$\epsilon = \epsilon(V) \cdot \epsilon(1/2, \pi)$$

(ii) Given $\sigma \in \text{Irr}(\text{Mp}(W))$, σ has nonzero theta lift to $\text{O}(V^\epsilon)$ if and only if

$$\begin{aligned}\epsilon &= z_\psi(\sigma) \cdot \epsilon(1/2, \sigma, \psi) \\ &= z_\psi(\sigma) \cdot \epsilon(1/2, \Theta_\psi(\sigma)).\end{aligned}$$

Here, the epsilon factors are the standard epsilon factors defined by the doubling zeta integral of PS-Rallis (Lapid-Rallis).

Question

Does the LLC map \mathcal{L}_ψ satisfy the usual properties of the Langlands parametrization?

Question

Does the LLC map \mathcal{L}_ψ satisfy the usual properties of the Langlands parametrization?

- If $\sigma \in \text{Irr}(\text{Mp}(W))$ has L -parameter ϕ , then is it the case that π is discrete series iff ϕ does not factor through a proper Levi?

Question

Does the LLC map \mathcal{L}_ψ satisfy the usual properties of the Langlands parametrization?

- If $\sigma \in \text{Irr}(\text{Mp}(W))$ has L -parameter ϕ , then is it the case that π is discrete series iff ϕ does not factor through a proper Levi?
- If σ is a Langlands quotient of an induced representation, is its L -parameter obtained from that of the inducing data in the usual way?

Question

Does the LLC map \mathcal{L}_ψ satisfy the usual properties of the Langlands parametrization?

- If $\sigma \in \text{Irr}(\text{Mp}(W))$ has L -parameter ϕ , then is it the case that π is discrete series iff ϕ does not factor through a proper Levi?
- If σ is a Langlands quotient of an induced representation, is its L -parameter obtained from that of the inducing data in the usual way?
- If η is trivial, does (ϕ, η) correspond to a ψ -generic rep of $\text{Mp}(W)$?

Question

Does the LLC map \mathcal{L}_ψ satisfy the usual properties of the Langlands parametrization?

- If $\sigma \in \text{Irr}(\text{Mp}(W))$ has L -parameter ϕ , then is it the case that π is discrete series iff ϕ does not factor through a proper Levi?
- If σ is a Langlands quotient of an induced representation, is its L -parameter obtained from that of the inducing data in the usual way?
- If η is trivial, does (ϕ, η) correspond to a ψ -generic rep of $\text{Mp}(W)$?
- Are natural invariants such as various L -factors and ϵ -factors preserved by \mathcal{L}_ψ ?

Parabolic Subgroups of $O(V)$

The parabolic subgroups $Q = L \cdot U$ of $O(V)$ have Levi factors

$$L = \mathrm{GL}(n_1) \times \dots \times \mathrm{GL}(n_r) \times \mathrm{O}(V_0)$$

with

$$2(n_1 + \dots + n_r) + \dim V_0 = \dim V.$$

Parabolic Subgroups of $O(V)$

The parabolic subgroups $Q = L \cdot U$ of $O(V)$ have Levi factors

$$L = \mathrm{GL}(n_1) \times \dots \times \mathrm{GL}(n_r) \times \mathrm{O}(V_0)$$

with

$$2(n_1 + \dots + n_r) + \dim V_0 = \dim V.$$

Given reps τ_i of $\mathrm{GL}(n_i)$ and π_0 of $\mathrm{O}(V_0)$, can form principal series rep

$$I_Q(\tau_1, \dots, \tau_r, \pi_0) = \mathrm{Ind}_Q^{\mathrm{O}(V)} \tau_1 \boxtimes \dots \tau_r \boxtimes \pi_0.$$

If this is a standard module, its unique irred quotient is denoted by

$$J_Q(\tau_1, \dots, \tau_r, \pi_0).$$

Parabolic Subgroups of $Mp(W)$

The parabolic subgroups $P = M \cdot N$ of $Sp(W)$ have Levi subgroups:

$$M \cong GL(n_1) \times \dots \times GL(n_r) \times Sp(W_0)$$

with

$$2n_1 + \dots + 2n_r + \dim W_0 = \dim W.$$

The restriction of the metaplectic cover to N splits uniquely, so one has:

$$\tilde{P} = \tilde{M} \cdot N.$$

The double cover \tilde{M} of M can be described as follows.

The Group \tilde{M}

The restriction of the metaplectic cover to $GL(m)$ is a double cover described by:

$$\tilde{GL}(m) = GL(m) \times \{\pm 1\}$$

with multiplication

$$(g_1, \epsilon_1) \cdot (g_2, \epsilon_2) = \\ (g_1 \cdot g_2, \epsilon_1 \cdot \epsilon_2 \cdot (\det g_1, \det g_2)).$$

The Group \tilde{M}

The restriction of the metaplectic cover to $GL(m)$ is a double cover described by:

$$\tilde{GL}(m) = GL(m) \times \{\pm 1\}$$

with multiplication

$$(g_1, \epsilon_1) \cdot (g_2, \epsilon_2) = \\ (g_1 \cdot g_2, \epsilon_1 \cdot \epsilon_2 \cdot (\det g_1, \det g_2)).$$

Then the preimage \tilde{M} of M in $Mp(W)$ is

$$\tilde{M} = \tilde{GL}(n_1) \times_{\mu_2} \dots \times_{\mu_2} \tilde{GL}(n_r) \times_{\mu_2} Mp(W_0).$$

In particular, to form genuine principal series reps of $Mp(W)$, we need genuine reps of $\tilde{GL}(m)$.

Representations of $\tilde{GL}(m)$

The genuine representation theory of $\tilde{GL}(m)$ turns out to be the same as the representation theory of $GL(m)$, because:

Representations of $\tilde{GL}(m)$

The genuine representation theory of $\tilde{GL}(m)$ turns out to be the same as the representation theory of $GL(m)$, because:

(i) The determinant map \det has a lifting:

$$\begin{array}{ccc} \tilde{GL}(m) & \xrightarrow{\det} & \tilde{GL}(1) \\ \downarrow & & \downarrow \\ GL(m) & \xrightarrow{\det} & GL(1) \end{array}$$

defined by

$$\det(g, \epsilon) = (\det(g), \epsilon)$$

Representations of $\tilde{GL}(m)$

The genuine representation theory of $\tilde{GL}(m)$ turns out to be the same as the representation theory of $GL(m)$, because:

(i) The determinant map \det has a lifting:

$$\begin{array}{ccc} \tilde{GL}(m) & \xrightarrow{\det} & \tilde{GL}(1) \\ \downarrow & & \downarrow \\ GL(m) & \xrightarrow{\det} & GL(1) \end{array}$$

defined by

$$\det(g, \epsilon) = (\det(g), \epsilon)$$

(ii) Using ψ , one may define a genuine character χ_ψ of $\tilde{GL}(1)$ (Weil). By composition, one obtains a genuine 1-dim character χ_ψ of $\tilde{GL}(m)$.

Representations of $\tilde{GL}(m)$

The genuine representation theory of $\tilde{GL}(m)$ turns out to be the same as the representation theory of $GL(m)$, because:

(i) The determinant map \det has a lifting:

$$\begin{array}{ccc} \tilde{GL}(m) & \xrightarrow{\det} & \tilde{GL}(1) \\ \downarrow & & \downarrow \\ GL(m) & \xrightarrow{\det} & GL(1) \end{array}$$

defined by

$$\det(g, \epsilon) = (\det(g), \epsilon)$$

(ii) Using ψ , one may define a genuine character χ_ψ of $\tilde{GL}(1)$ (Weil). By composition, one obtains a genuine 1-dim character χ_ψ of $\tilde{GL}(m)$.

(iii) Tensoring with χ_ψ gives

$$\text{Irr}(GL(m)) \rightarrow \text{Irr}(\tilde{GL}(m)).$$

Principal Series Reps of $Mp(W)$

Given reps τ_i of $GL(n_i)$ and σ_0 of $Mp(W_0)$, one has the induced rep

$$I_{P,\psi}(\tau_1, \dots, \tau_r, \sigma_0) = \text{Ind}_{\tilde{P}}^{Mp(W)} \chi_\psi \tau_1 \boxtimes \dots \boxtimes \chi_\psi \tau_r \boxtimes \sigma_0.$$

If this is a standard module, its unique irred quotient is denoted by

$$J_{P,\psi}(\tau_1, \dots, \tau_r, \sigma_0)$$

Central Sign of $\sigma \in \text{Irr}(\text{Mp}(W))$

Recall that the center of $\text{Mp}(W)$ is the preimage \tilde{Z} of the center $Z \cong \mu_2$ of $\text{Sp}(W)$. So \tilde{Z} is a group of size 4 and has 2 irred genuine characters.

Central Sign of $\sigma \in \text{Irr}(\text{Mp}(W))$

Recall that the center of $\text{Mp}(W)$ is the preimage \tilde{Z} of the center $Z \cong \mu_2$ of $\text{Sp}(W)$. So \tilde{Z} is a group of size 4 and has 2 irred genuine characters.

To distinguish between them, note that Z is the subgroup

$$\{\pm I_n\} \subset \text{GL}(n) \quad (\text{the Siegel Levi}).$$

Thus, the character χ_ψ of the Siegel Levi restricts to a genuine character of \tilde{Z} .

Central Sign of $\sigma \in \text{Irr}(\text{Mp}(W))$

Recall that the center of $\text{Mp}(W)$ is the preimage \tilde{Z} of the center $Z \cong \mu_2$ of $\text{Sp}(W)$. So \tilde{Z} is a group of size 4 and has 2 irred genuine characters.

To distinguish between them, note that Z is the subgroup

$$\{\pm I_n\} \subset \text{GL}(n) \quad (\text{the Siegel Levi}).$$

Thus, the character χ_ψ of the Siegel Levi restricts to a genuine character of \tilde{Z} .

Every genuine irrep σ of $\text{Mp}(W)$ has a central character, and we set

$$z_\psi(\sigma) = \begin{cases} +1, & \text{if its central character is } \chi_\psi; \\ -1, & \text{otherwise.} \end{cases}$$

Theorem D

Suppose that $\sigma \in \text{Irr}(\text{Mp}(W))$ and $\pi = \Theta_\psi(\sigma)$ (so $p \neq 2$).

Theorem D

Suppose that $\sigma \in \text{Irr}(\text{Mp}(W))$ and $\pi = \Theta_\psi(\sigma)$ (so $p \neq 2$).

- σ is d.s. iff π is d.s. Moreover, the formal degrees of π and σ are equal (wrt suitable measures).

Theorem D

Suppose that $\sigma \in \text{Irr}(\text{Mp}(W))$ and $\pi = \Theta_\psi(\sigma)$ (so $p \neq 2$).

- σ is d.s. iff π is d.s. Moreover, the formal degrees of π and σ are equal (wrt suitable measures).
- σ is tempered iff π is tempered. Indeed, if

$$\sigma \subset I_{P,\psi}(\tau_1, \dots, \tau_r, \sigma_0)$$

with τ_i and σ_0 unitary d.s., then

$$\pi \subset I_Q(\tau_1, \dots, \tau_r, \pi_0)$$

with

$$\pi_0 = \Theta_\psi(\sigma_0).$$

Theorem D

Suppose that $\sigma \in \text{Irr}(\text{Mp}(W))$ and $\pi = \Theta_\psi(\sigma)$ (so $p \neq 2$).

- σ is d.s. iff π is d.s. Moreover, the formal degrees of π and σ are equal (wrt suitable measures).
- σ is tempered iff π is tempered. Indeed, if

$$\sigma \subset I_{P,\psi}(\tau_1, \dots, \tau_r, \sigma_0)$$

with τ_i and σ_0 unitary d.s., then

$$\pi \subset I_Q(\tau_1, \dots, \tau_r, \pi_0)$$

with

$$\pi_0 = \Theta_\psi(\sigma_0).$$

- If $\sigma = J_{P,\psi}(\tau_1, \dots, \tau_r, \sigma_0)$ is a Langlands quotient, then

$$\pi = J_Q(\tau_1, \dots, \tau_r, \Theta_\psi(\sigma_0)).$$

- If π is generic, then σ is ψ -generic. If σ is tempered and ψ -generic, then π is generic.

- If π is generic, then σ is ψ -generic. If σ is tempered and ψ -generic, then π is generic.
- If π and σ are d.s., and τ is a supercuspidal rep of $GL(r)$, then

$$\mu(s, \pi \times \tau, \psi) = \mu(s, \sigma \times \tau, \psi).$$

(Plancherel measures here)

- If π is generic, then σ is ψ -generic. If σ is tempered and ψ -generic, then π is generic.
- If π and σ are d.s., and τ is a supercuspidal rep of $GL(r)$, then

$$\mu(s, \pi \times \tau, \psi) = \mu(s, \sigma \times \tau, \psi).$$

(Plancherel measures here)

- For any character χ of $GL(1)$,

$$\begin{cases} L(s, \pi \times \chi) = L(s, \sigma \times \chi, \psi) \\ \epsilon(s, \pi \times \chi, \psi) = \epsilon(s, \sigma \times \chi, \psi) \end{cases}$$

(standard factors defined by doubling zeta integral)

- If π is generic, then σ is ψ -generic. If σ is tempered and ψ -generic, then π is generic.
- If π and σ are d.s., and τ is a supercuspidal rep of $GL(r)$, then

$$\mu(s, \pi \times \tau, \psi) = \mu(s, \sigma \times \tau, \psi).$$

(Plancherel measures here)

- For any character χ of $GL(1)$,

$$\begin{cases} L(s, \pi \times \chi) = L(s, \sigma \times \chi, \psi) \\ \epsilon(s, \pi \times \chi, \psi) = \epsilon(s, \sigma \times \chi, \psi) \end{cases}$$

(standard factors defined by doubling zeta integral)

- if π is generic, then for any irrep τ of $GL(r)$,

$$\begin{cases} L(s, \pi \times \tau) = L(s, \sigma \times \tau, \psi) \\ \epsilon(s, \pi \times \tau, \psi) = \epsilon(s, \sigma \times \tau, \psi) \end{cases}$$

(LHS: Shahidi, RHS: Szpruch)

Dependence on ψ

For $c \in F^\times$, let

$$\psi_c(x) = \psi(cx).$$

How are Θ_{ψ_c} and Θ_ψ related?

Dependence on ψ

For $c \in F^\times$, let

$$\psi_c(x) = \psi(cx).$$

How are Θ_{ψ_c} and Θ_ψ related?

More precisely, suppose that

$$\mathcal{L}_\psi(\sigma) = (\phi, \eta)$$

and

$$\mathcal{L}_{\psi_c}(\sigma) = (\phi_c, \eta_c).$$

How are (ϕ, η) and (ϕ_c, η_c) related?

Dependence on ψ

For $c \in F^\times$, let

$$\psi_c(x) = \psi(cx).$$

How are Θ_{ψ_c} and Θ_ψ related?

More precisely, suppose that

$$\mathcal{L}_\psi(\sigma) = (\phi, \eta)$$

and

$$\mathcal{L}_{\psi_c}(\sigma) = (\phi_c, \eta_c).$$

How are (ϕ, η) and (ϕ_c, η_c) related?

Suppose that

$$\phi = \bigoplus_i n_i \phi_i,$$

then

$$A_\phi = \bigoplus_{i: \phi_i \text{ is symplectic}} \mathbb{Z}/2\mathbb{Z} a_i.$$

Theorem E

(i) $\phi_c = \phi \otimes \chi_c$;

It follows by (i) that we have canonical identification of component groups:

$$A_\phi = A_{\phi_c} = \bigoplus_i \mathbb{Z}/2\mathbb{Z}a_i,$$

so that it makes sense to compare η and η_c .

Theorem E

(i) $\phi_c = \phi \otimes \chi_c$;

It follows by (i) that we have canonical identification of component groups:

$$A_\phi = A_{\phi_c} = \bigoplus_i \mathbb{Z}/2\mathbb{Z}a_i,$$

so that it makes sense to compare η and η_c .

(ii) the characters η and η_c are related by:

$$\eta_c(a_i)/\eta(a_i) = \epsilon(1/2, \phi_i) \cdot \epsilon(1/2, \phi_i \otimes \chi_c) \cdot \chi_c(-1)^{\frac{1}{2} \dim \phi_i} \in \{\pm 1\}.$$

Theorem E

(i) $\phi_c = \phi \otimes \chi_c$;

It follows by (i) that we have canonical identification of component groups:

$$A_\phi = A_{\phi_c} = \bigoplus_i \mathbb{Z}/2\mathbb{Z}a_i,$$

so that it makes sense to compare η and η_c .

(ii) the characters η and η_c are related by:

$$\eta_c(a_i)/\eta(a_i) = \epsilon(1/2, \phi_i) \cdot \epsilon(1/2, \phi_i \otimes \chi_c) \cdot \chi_c(-1)^{\frac{1}{2} \dim \phi_i} \in \{\pm 1\}.$$

The proof of Thm. E uses the **Gross-Prasad conjecture** for special orthogonal groups recently proved by Wadsworth.

Work of Dani Szpruch

Work of Dani Szpruch

The **Langlands-Shahidi method** is a theory which defines local L-factors and ϵ -factors for generic representations of a linear reductive group G .

Work of Dani Szpruch

The **Langlands-Shahidi method** is a theory which defines local L-factors and ϵ -factors for generic representations of a linear reductive group G .

In his thesis, Szpruch has extended this theory to the case of $\mathrm{Mp}(W)$. For ψ -generic representations σ of $\mathrm{Mp}(W)$, and irreducible representations ρ of $\mathrm{GL}_r(F)$, he defines

$$L(s, \sigma \times \rho, \psi) \quad \text{and} \quad \epsilon(s, \sigma \times \rho, \psi).$$

Relation with Character Theory

One hallmark of the LLC is its relation with character theory; in particular the notion of stability and the theory of endoscopy.

Relation with Character Theory

One hallmark of the LLC is its relation with character theory; in particular the notion of stability and the theory of endoscopy.

- If

$$\Pi_{\psi, \phi} = \{\sigma : \mathcal{L}_{\psi}(\sigma) = (\phi, *)\}$$

is a tempered L-packet, then the sum

$$\sum_{\sigma \in \Pi_{\psi, \phi}} \text{Trace}(\sigma)$$

is supposed to be a "stable" distribution;

Relation with Character Theory

One hallmark of the LLC is its relation with character theory; in particular the notion of stability and the theory of endoscopy.

- If

$$\Pi_{\psi, \phi} = \{\sigma : \mathcal{L}_{\psi}(\sigma) = (\phi, *)\}$$

is a tempered L-packet, then the sum

$$\sum_{\sigma \in \Pi_{\psi, \phi}} \text{Trace}(\sigma)$$

is supposed to be a "stable" distribution;

- other linear combinations of $\text{Trace}(\sigma)$'s are supposed to be "transfers" of stable distribution from "endoscopic groups".

Relation with Character Theory

One hallmark of the LLC is its relation with character theory; in particular the notion of stability and the theory of endoscopy.

- If

$$\Pi_{\psi, \phi} = \{\sigma : \mathcal{L}_{\psi}(\sigma) = (\phi, *)\}$$

is a tempered L-packet, then the sum

$$\sum_{\sigma \in \Pi_{\psi, \phi}} \text{Trace}(\sigma)$$

is supposed to be a "stable" distribution;

- other linear combinations of $\text{Trace}(\sigma)$'s are supposed to be "transfers" of stable distribution from "endoscopic groups".

Basic notions in "... " needs to be developed.

Work of Wenwei Li

Extending earlier works of Adams-Schulze and Renard, Wenwei Li has recently developed a definitive theory of endoscopy for $Mp(W)$, and much more!

Work of Wenwei Li

Extending earlier works of Adams-Schulze and Renard, Wenwei Li has recently developed a definitive theory of endoscopy for $\mathrm{Mp}(W)$, and much more!

- For each pair (a, b) with $a + b = n$, he defined transfer factors

$$\Delta_{a,b} : \mathrm{Mp}(W) \times (\mathrm{SO}(2a + 1) \times \mathrm{SO}(2b + 1)) \rightarrow \mathbb{C}.$$

Work of Wenwei Li

Extending earlier works of Adams-Schulze and Renard, Wenwei Li has recently developed a definitive theory of endoscopy for $\mathrm{Mp}(W)$, and much more!

- For each pair (a, b) with $a + b = n$, he defined transfer factors

$$\Delta_{a,b} : \mathrm{Mp}(W) \times (\mathrm{SO}(2a + 1) \times \mathrm{SO}(2b + 1)) \rightarrow \mathbb{C}.$$

- showed existence of transfer mappings

$$\begin{array}{c} C_c^\infty(\mathrm{Mp}(W)) \\ \downarrow t_{a,b} \\ C_c^\infty(\mathrm{SO}(2a + 1) \times \mathrm{SO}(2b + 1)), \end{array}$$

giving rise to a map

$$\mathcal{D}_{st}(\mathrm{SO}(2a+1) \times \mathrm{SO}(2b+1))$$

$$t_{a,b}^* \downarrow$$

$$\mathcal{D}_{gen}(\mathrm{Mp}(W)),$$

between spaces of distributions.

$$\mathcal{D}_{st}(\mathrm{SO}(2a+1) \times \mathrm{SO}(2b+1))$$

$$t_{a,b}^* \downarrow$$

$$\mathcal{D}_{gen}(\mathrm{Mp}(W)),$$

between spaces of distributions.

- Established the ordinary and weighted fundamental lemmas for the transfer $t_{a,b}$.

$$\mathcal{D}_{st}(\mathrm{SO}(2a+1) \times \mathrm{SO}(2b+1))$$

$$t_{a,b}^* \downarrow$$

$$\mathcal{D}_{gen}(\mathrm{Mp}(W)),$$

between spaces of distributions.

- Established the ordinary and weighted fundamental lemmas for the transfer $t_{a,b}$.

In other words, the elliptic endoscopic groups of $\mathrm{Mp}(W)$ are the groups

$$\mathrm{SO}(2a+1) \times \mathrm{SO}(2b+1),$$

with $a + b = n$.

Automorphic Forms of $Mp(W)$

Now we come to the global story.

We will consider the discrete spectrum

$$L^2_{disc} = L^2_{gen, disc}(\mathrm{Sp}(W)(F) \backslash \mathrm{Mp}(W)(\mathbb{A}))$$

of genuine (square-integrable) automorphic forms on $Mp(W)$.

Problem: describe its decomposition into irreducible reps, in the spirit of Arthur's conjecture.

The Tempered Part

Let's focus on the tempered part $L^2_{temp,disc}$. For a fixed additive automorphic character ψ , one expects that

$$L^2_{temp,disc} = \bigoplus_{\phi} L^2_{\phi,\psi}$$

with

$$\phi = \bigoplus_{i=1}^r \phi_i : L_F \longrightarrow \mathrm{Sp}_{2n}(\mathbb{C}),$$

(where L_F is the hypothetical Langlands group) is multiplicity-free and each ϕ_i is symplectic.

The Tempered Part

Let's focus on the tempered part $L^2_{temp, disc}$. For a fixed additive automorphic character ψ , one expects that

$$L^2_{temp, disc} = \bigoplus_{\phi} L^2_{\phi, \psi}$$

with

$$\phi = \bigoplus_{i=1}^r \phi_i : L_F \longrightarrow \mathrm{Sp}_{2n}(\mathbb{C}),$$

(where L_F is the hypothetical Langlands group) is multiplicity-free and each ϕ_i is symplectic.

Following Arthur, can reformulate the notion ϕ without recourse to L_F . One can take ϕ to be a collection $\{\pi_1, \dots, \pi_r\}$ of cuspidal representations of $\mathrm{GL}(n_i)$ such that

- $n_1 + \dots + n_r = n$;
- $L^2(s, \pi_i, \wedge^2)$ has a pole at $s = 1$.

Global Packet associated to ϕ

For a given ϕ , what is $L_{\phi, \psi}^2$?

Global Packet associated to ϕ

For a given ϕ , what is $L_{\phi, \psi}^2$?

Given $\phi = \phi_1 \oplus \dots \oplus \phi_r$, one obtains

- local L-parameters ϕ_v and thus local L-packets

$$\Pi_{\phi_v, \psi_v} = \{\sigma_{\eta_v} : \eta_v \in \text{Irr}(A_{\phi_v})\}$$

Global Packet associated to ϕ

For a given ϕ , what is $L_{\phi,\psi}^2$?

Given $\phi = \phi_1 \oplus \dots \oplus \phi_r$, one obtains

- local L-parameters ϕ_v and thus local L-packets

$$\Pi_{\phi_v,\psi_v} = \{\sigma_{\eta_v} : \eta_v \in \text{Irr}(A_{\phi_v})\}$$

- taking tensor products of these local packets, get global packet

$$\Pi_{\phi,\psi} = \{\sigma_{\eta} = \bigotimes_v \sigma_{\eta_v} : \eta \in \text{Irr}(\prod_v A_{\phi_v})\}$$

Global Packet associated to ϕ

For a given ϕ , what is $L_{\phi, \psi}^2$?

Given $\phi = \phi_1 \oplus \dots \oplus \phi_r$, one obtains

- local L-parameters ϕ_v and thus local L-packets

$$\Pi_{\phi_v, \psi_v} = \{\sigma_{\eta_v} : \eta_v \in \text{Irr}(A_{\phi_v})\}$$

- taking tensor products of these local packets, get global packet

$$\Pi_{\phi, \psi} = \{\sigma_{\eta} = \bigotimes_v \sigma_{\eta_v} : \eta \in \text{Irr}(\prod_v A_{\phi_v})\}$$

- a global component group

$$A_{\phi} = \mathbb{Z}/2\mathbb{Z}a_1 \times \dots \times \mathbb{Z}/2\mathbb{Z}a_r$$

and a diagonal map

$$\Delta : A_{\phi} \rightarrow \prod_v A_{\phi_v}$$

Langlands-Arthur's Multiplicity Formula

One conjectures that

$$L^2_{\phi,\psi} = \bigoplus_{\eta} m_{\eta} \cdot \sigma_{\eta}$$

where the multiplicity m_{η} is defined by

Langlands-Arthur's Multiplicity Formula

One conjectures that

$$L_{\phi,\psi}^2 = \bigoplus_{\eta} m_{\eta} \cdot \sigma_{\eta}$$

where the multiplicity m_{η} is defined by

$$m_{\eta} = \langle \Delta^*(\eta), \epsilon_{\psi} \rangle,$$

and ϵ_{ψ} is a quadratic character of A_{ϕ} given by

$$\epsilon_{\psi}(a_i) = \epsilon(1/2, \phi_i).$$

Langlands-Arthur's Multiplicity Formula

One conjectures that

$$L_{\phi, \psi}^2 = \bigoplus_{\eta} m_{\eta} \cdot \sigma_{\eta}$$

where the multiplicity m_{η} is defined by

$$m_{\eta} = \langle \Delta^*(\eta), \epsilon_{\psi} \rangle,$$

and ϵ_{ψ} is a quadratic character of A_{ϕ} given by

$$\epsilon_{\psi}(a_i) = \epsilon(1/2, \phi_i).$$

This conjecture amounts to a description of $L_{disc, temp}^2$ in terms of the tempered discrete spectrum of $SO(2n+1)$.

When $n=1$, this is the main result of Waldspurger's work on the Shimura correspondence.

Towards a stable trace formula for $Mp(W)$

The work of Wenwei Li has given a approach to proving the global conjecture via the **trace formula**.

Towards a stable trace formula for $Mp(W)$

The work of Wenwei Li has given a approach to proving the global conjecture via the **trace formula**.

- He has begun the study of the trace formula for general covering groups.
- For $Mp(W)$, he has begun the stabilization of the trace formula.

Towards a stable trace formula for $\mathrm{Mp}(W)$

The work of Wenwei Li has given a approach to proving the global conjecture via the **trace formula**.

- He has begun the study of the trace formula for general covering groups.
- For $\mathrm{Mp}(W)$, he has begun the stabilization of the trace formula.

The stable trace formula should look like:

$$\mathrm{TF}_{\mathrm{Mp}(W)}(f) = \sum_{a+b=n} c_{a,b} \cdot \mathrm{STF}_{\mathrm{SO}(2a+1) \times \mathrm{SO}(2b+1)}(t_{a,b}(f))$$

where $c_{a,b} = 1/4$ if $ab \neq 0$ and $1/2$ otherwise.

Using an elliptic stable trace formula, together with the global theta correspondence and some extra work, one should obtain

- a proof of stability properties of local L-packets and character identities for $\mathrm{Mp}(W)$;
- the multiplicity formula for cuspidal representations of $\mathrm{Mp}(W)$ with sufficiently many supercuspidal local components

This is joint work with W. W. Li.

Conclusion

- Essentially all aspects of the Langlands philosophy extend beautifully to the metaplectic groups $Mp(W)$. One hopes that a general theory in the style of the Langlands program can be developed for general covering groups.

Conclusion

- Essentially all aspects of the Langlands philosophy extend beautifully to the metaplectic groups $Mp(W)$. One hopes that a general theory in the style of the Langlands program can be developed for general covering groups.
- Several recent progresses indicate that the time is ripe for a systematic development of such a theory. A systematic study of the trace formula of covering groups will be the key, but further explicit examples would be very welcome.