

# MATH 142A: Introduction to Analysis

[math-old.ucsd.edu/~ynemish/teaching/142a](http://math-old.ucsd.edu/~ynemish/teaching/142a)

Today: Ordered field  
> Q&A: January 7

Next: Ross § 4

Week 1:

- visit course website
- homework 0 (due Friday, January 7)
- join Piazza

## Fields

$\mathbb{N} \subsetneq \mathbb{Z} \subsetneq \mathbb{Q} \subsetneq \mathbb{R}$  (proper subsets)

Let  $F$  be a set with two binary operations

$+ : F \times F \rightarrow F$  (addition) and  $\cdot : F \times F \rightarrow F$  (multiplication)

Consider the following properties :

A1.  $a + (b+c) = (a+b) + c \quad \forall a, b, c \in F$  (associativity)

$(1:2):2 \neq 1:(2:2), \quad 2^3 = 2^{(2^2)} \neq (2^2)^3 \leftarrow \text{not associative}$

A2.  $a+b = b+a \quad \forall a, b \in F$  (commutativity) [" $\forall$ " means "for all"]

$3-2 \neq 2-3 \leftarrow \text{not commutative}$

A3.  $\exists 0 \in F$  s.t.  $a+0=a \quad \forall a \in F$  (neutral element)

$0 \notin \mathbb{N}$  [ $\exists$  means "there exists"]

A4.  $\forall a \in F \quad \exists (-a) \in F$  s.t.  $a + (-a) = 0$  (additive inverse of  $a$ )

$$\mathbb{Q}_{\geq 0} := \{r \in \mathbb{Q} : r \geq 0\}$$

$$-1 \notin \mathbb{Q}_{\geq 0}$$

Fields (cont)

M1.  $a(bc) = (ab)c \quad \forall a, b, c \in F$  (associativity)

M2.  $ab = ba \quad \forall a, b \in F$  (commutativity)

M3.  $\exists 1 \in F$  s.t.  $a \cdot 1 = a \quad \forall a \in F$  (neutral element)

M4.  $\forall a \in F$  s.t.  $a \neq 0 \quad \exists a^{-1} \in F$  s.t.  $a a^{-1} = 1$  (multiplicative inverse)

$$F = \{M \in \mathbb{R}^{2 \times 2} : \det M \neq 0\}$$

$$\begin{pmatrix} 0 & 1 \\ 1 & 1 \end{pmatrix} \begin{pmatrix} 1 & 1 \\ 1 & 0 \end{pmatrix} \neq \begin{pmatrix} 1 & 1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} 0 & 1 \\ 1 & 1 \end{pmatrix}$$

DL  $a(b+c) = ab+ac \quad \forall a, b, c \in F$

Definition (Field) Set  $F$  with more than one element and binary operations  $+$  and  $\cdot$  satisfying A1-A4, M1-M4, DL is called a field.

A1-A4, M1-M4 and DL are called the field axioms

Remark  $\mathbb{Q}, \mathbb{R}$  are fields,  $\mathbb{N}, \mathbb{Z}$  are not fields (with usual  $+, \cdot$ )

## Consequences of field axioms

Theorem 3.1 Let  $\mathbb{F}$  with operations  $+$  and  $\cdot$  be a field.

Then for any  $a, b, c \in \mathbb{F}$

$$(i) a+c = b+c \Rightarrow a=b \quad (iv) (-a)(-b) = ab$$

$$(ii) a \cdot 0 = 0 \quad (v) ac = bc \wedge c \neq 0 \Rightarrow a = b$$

$$(iii) (-a)b = -ab \quad (vi) ab = 0 \Rightarrow a=0 \vee b=0$$

Proof. (i)  $a+c = b+c \Rightarrow (a+c)+(-c) = (b+c)+(-c)$

$$(a+c)+(-c) \stackrel{A_1}{=} a+ (c+(-c)) \stackrel{A_4}{=} a+0 = a, (b+c)+(-c) \stackrel{A_1}{=} b+ (c+(-c)) \stackrel{A_4}{=} b+0 = b$$

which implies that  $a=b$

$$(ii) a \cdot 0 = a \cdot (0+0) \stackrel{DL}{=} a \cdot 0 + a \cdot 0 \quad | \quad \Rightarrow a \cdot 0 + a \cdot 0 = 0 + a \cdot 0 \stackrel{(i)}{\Rightarrow} a \cdot 0 = 0 \\ a \cdot 0 = a \cdot 0 + 0 = 0 + a \cdot 0 \quad | \quad \blacksquare$$

Prop If  $0_1$  and  $0_2$  are (additive) neutral elements, then  $0_1 = 0_2$ .

Proof.  $0_1 \stackrel{A_3}{=} 0_1 + 0_2 \stackrel{A_2}{=} 0_2 + 0_1 \stackrel{A_3}{=} 0_2 \quad \blacksquare$

## Ordered fields

Definition Set  $S$  with a (binary) relation  $\leq$  is called  
**linearly ordered** if

(01)  $\forall a, b \in S$  either  $a \leq b$  or  $b \leq a$

(02)  $\forall a, b \in S$  ( $a \leq b \wedge b \leq a \Rightarrow a = b$ ) [antisymmetry]

(03)  $\forall a, b, c \in S$  ( $a \leq b \wedge b \leq c \Rightarrow a \leq c$ ) [transitivity]

Definition Let  $F$  be a set with operations  $+$  and  $\cdot$  and  
order relation  $\leq$ .  $F$  is called an **ordered field** if

- $F$  with  $+$  and  $\cdot$  is a **field**
- $F$  with  $\leq$  is **linearly ordered**
- (04)  $a \leq b \Rightarrow a+c \leq b+c \quad \forall a, b, c \in F$
- (05)  $a \leq b \wedge 0 \leq c \Rightarrow ac \leq bc$

## Properties of ordered fields

Theorem 3.2 Let  $\mathbb{F}$  be an ordered field with operations  $+$ ,  $\cdot$  and order relation  $\leq$ . Then  $\forall a, b, c \in \mathbb{F}$

- (i)  $a \leq b \Rightarrow -b \leq -a$  (v)  $0 < 1$
- (ii)  $a \leq b \wedge c \leq 0 \Rightarrow bc \leq ac$  (vi)  $0 < a \Rightarrow 0 < a^{-1}$
- (iii)  $0 \leq a \wedge 0 \leq b \Rightarrow 0 \leq ab$  (vii)  $0 < a < b \Rightarrow 0 < b^{-1} < a^{-1}$
- (iv)  $0 \leq a^2$  [ $a^2 = a \cdot a$ ] [“ $a < b$ ” means ‘ $a \leq b \wedge a \neq b$ ’]

Proof. (i)  $a \leq b \stackrel{\text{O4}}{\Rightarrow} a + ((-a) + (-b)) \leq b + ((-a) + (-b)) \stackrel{\text{A1-A4}}{\Rightarrow} -b \leq -a$

(ii)  $0+0=0 \Rightarrow -0=0$ , therefore  $c \leq 0 \stackrel{\text{(i)}}{\Rightarrow} 0 \leq -c$ . Then  
 $a \leq b \wedge 0 \leq -c \stackrel{\text{O5}}{\Rightarrow} a(-c) \leq b(-c) \stackrel{\text{T3.1}}{\Rightarrow} -ac \leq -bc \stackrel{\text{(i)}}{\Rightarrow} bc \leq ac$

(iv) By O1 either  $a \leq 0$  or  $0 \leq a$ .  $0 \leq a \stackrel{\text{O5}}{\Rightarrow} 0 \cdot a \leq a \cdot a \Rightarrow 0 \leq a^2$   
 $a \leq 0 \Rightarrow 0 \leq (-a)(-a) \stackrel{\text{T3.1}}{\Rightarrow} 0 \leq a^2$

■

## Absolute value

Let  $\mathbb{F}$  be an ordered field

Def 3.3. Let  $a \in \mathbb{F}$ . We call  $|a| := \begin{cases} a & \text{if } 0 \leq a \\ -a & \text{if } a \leq 0 \end{cases}$

the **absolute value** of  $a$ .

Def 3.4 Let  $a, b \in \mathbb{F}$ . We call  $\text{dist}(a, b) := |a - b|$

the **distance** between  $a$  and  $b$   $[a - b := a + (-b)]$

Thm 3.5 (i)  $0 \leq |a| \quad \forall a \in \mathbb{F}$

(ii)  $|ab| = |a||b| \quad \forall a, b \in \mathbb{F}$

(iii)  $|a+b| \leq |a| + |b| \quad \forall a, b \in \mathbb{F}$  (Triangle inequality)

Proof (i) Follows from the definition and Thm 3.2 (i).

(ii) Exercise (check 4 cases)

### Proof (cont) (iii)

Step 1:  $\forall c \in F, 0 \leq c \Rightarrow -|c| \leq c \leq |c|$

Proof:  $0 \leq c \Rightarrow |c| = c \wedge -c \leq 0 \stackrel{0l}{\Rightarrow} -|c| \leq 0 \leq c \leq |c|$

Step 2:  $\forall c \in F, c \leq 0 \Rightarrow -|c| \leq c \leq |c|$

Proof:  $c \leq 0 \Rightarrow (|c| = -c) \wedge (-|c| = c) \wedge (0 \leq |c|) \Rightarrow -|c| \leq c \leq 0 \leq |c|$

Step 3:  $-|a| \leq a \leq |a|, -|b| \leq b \leq |b|$

Follows from Step 1 and Step 2.

Step 4:  $-|a|-|b| \leq a-|b| \leq a+b \leq |a|+b \leq |a|+|b|$

$$\Rightarrow \begin{cases} a+b \leq |a|+|b| \\ -(a+b) \leq -(-|a|-|b|) = |a|+|b| \end{cases} \Rightarrow |a+b| \leq |a|+|b|$$

■

Corollary  $\forall a, b, c \in F \quad \text{dist}(a, c) \leq \text{dist}(a, b) + \text{dist}(b, c)$

Proof. Exercise (Hint: Define  $x = a-b, y = b-c$ )