

MATH180C: Introduction to Stochastic Processes II

[Lecture A00: math.ucsd.edu/~ynemish/teaching/180cA](http://math.ucsd.edu/~ynemish/teaching/180cA)

[Lecture B00: math.ucsd.edu/~ynemish/teaching/180cB](http://math.ucsd.edu/~ynemish/teaching/180cB)

Today: Brownian motion

Next: PK 8.1-8.2

Week 9:

- **CA P E S**
- homework 7 (due Friday, May 27)
- HW6 regrades are active on Gradescope until May 28, 11 PM
- Friday May 27 office hour: AP&M 7321

Brownian motion. Definition

Def. **Brownian motion** with diffusion coefficient σ^2 is a continuous time stochastic process $(B_t)_{t \geq 0}$ satisfying

- (i) $B(0) = 0$, $B(t)$ is continuous as a function of t
- (ii) For all $0 \leq s < t < \infty$ $B(t) - B(s)$ is a Gaussian random variable with mean 0 and variance $\sigma^2(t-s)$
- (iii) The increments of B are independent: if $0 = t_0 < t_1 < \dots < t_n$ then $\{B(t_i) - B(t_{i-1})\}_{i=1}^n$ are independent (Gaussian) r.v.s.

$\sigma^2 = 1 \leftarrow$ standard BM

BM as a Gaussian process

Def. Stochastic process $(X_t)_{t \geq 0}$ is called a Gaussian process

if for any $0 \leq t_1 < t_2 < \dots < t_n$

$(X_{t_1}, \dots, X_{t_n})$ is a Gaussian vector, or equivalently

for any $c_1, \dots, c_n \in \mathbb{R}$

$c_1 X_{t_1} + c_2 X_{t_2} + \dots + c_n X_{t_n}$ is a Gaussian r.v.

Recall that the distribution of a Gaussian vector is uniquely defined by its mean and covariance matrix.

Similarly, each Gaussian process is uniquely described by

$$\mu(t) = E(X_t) \quad \text{and} \quad \Gamma(s, t) = \text{Cov}(X_s, X_t) \geq 0$$

↑ covariance function

BM as a Gaussian process

Proposition BM is a Gaussian process with

$$\mu(t) = 0 \quad \text{and} \quad \Gamma(s, t) = \min\{s, t\} = s \wedge t$$

Proof. For any $0 = t_0 < t_1 < t_2 < \dots < t_n$, $B_{t_j} - B_{t_{j-1}}$ are indep.

Gaussian, thus

$$\sum_{i=1}^n c_i B_{t_i} = \sum_{i=1}^n c_i \sum_{j=1}^i (B_{t_j} - B_{t_{j-1}}) = \sum_{j=1}^n \sum_{i=j}^n c_i (B_{t_j} - B_{t_{j-1}})$$

is also Gaussian.

By definition $\mu(t) = E(B_t) = 0$. Let $s < t$.

$$\begin{aligned} \text{Then } \Gamma(s, t) &= \text{Cov}(B_s, B_t) \\ &= \text{Cov}(B_s, B_s + (B_t - B_s)) \\ &= \text{Cov}(B_s, B_s) + \text{Cov}(B_s, B_t - B_s) \\ &= s + 0 = s = \min\{s, t\} \quad \blacksquare \end{aligned}$$

Some properties of BM

Proposition. Let $(B_t)_{t \geq 0}$ be a standard BM. Then

- (i) For any $s > 0$, the process $(B_{t+s} - B_s, t \geq 0)$ is a BM independent of $(B_u, 0 \leq u \leq s)$.
- (ii) The process $(-B_t, t \geq 0)$ is a BM
- (iii) For any $c > 0$, the process $(cB_{\frac{t}{c}}, t \geq 0)$ is a BM
- (iv) The process $(X_t)_{t \geq 0}$ defined by $X_0 = 0$, $X_t = tB_{\frac{1}{t}}$ for $t > 0$ is a BM.

Proof (i) Define $X_t = B_{t+s} - B_s$. Then $X_0 = 0$, $X_{t_2} - X_{t_1} = B_{t_2+s} - B_{t_1+s}$
 \Rightarrow independent Gaussian increments, $E(X_{t_2} - X_{t_1}) = 0$, $\text{Var}(X_{t_2} - X_{t_1}) = t_2 - t_1$

$(X_t)_{t \geq 0}$ has continuous paths $\Rightarrow (X_t)_{t \geq 0}$ is a BM

(iv) X_t is Gaussian, for $s < t$ $\text{Cov}(sB_{\frac{1}{s}}, tB_{\frac{1}{t}}) = st \cdot \min\{\frac{1}{s}, \frac{1}{t}\} = s$


Proof of $\lim_{t \rightarrow 0} X_t = 0$ is more technical, thus omitted. \blacksquare

Construction of BM

BM can be constructed as a limit of properly rescaled random walks.

Let $\{\xi_k\}_{k=1}^{\infty}$ be a sequence of i.i.d. r.v.s, $E(\xi_i) = 0$,
 $\text{Var}(\xi_i) = \sigma^2 < \infty$. Denote $S_m = \sum_{k=1}^m \xi_k$ and define

$$X_t^n = \frac{1}{\sigma\sqrt{n}} \left(S_{[nt]} + (nt - [nt]) \xi_{[nt]+1} \right)$$



Theorem (Donsker) $(X_t^n)_{t \geq 0}$ converges in distribution
to the standard BM.