

MATH 285: Stochastic Processes

math-old.ucsd.edu/~ynemish/teaching/285

Today: Branching processes

- Homework 4 is due on Friday, February 11, 11:59 PM

Galton - Watson Branching Process

Consider a population whose evolution (reproduction) is determined by the following rules

(i) Each individual produces k offsprings with probability


$$p_k, \text{ where } \sum_{k=0}^{\infty} p_k = 1$$

(ii) All individuals reproduce independently

Denote by X_n the size of the n -th generation. The number of individuals in the n -th generation

depends only on the number of individuals in generation $n-1 \Rightarrow (X_n)$ is a Markov chain

If $X_n = 0$ for some n , then $\forall m > n \quad X_m = 0$ - extinction.



Galton - Watson Branching Process

Q: What is the probability that the population never goes extinct? $\mathbb{P}[X_n \geq 1 \text{ } \forall n \in \mathbb{N}] = ?$

Direct computation: Let $\{Y_i\}_{i=1}^{\infty}$ be i.i.d. random variables with $Y_i \in \{0, 1, \dots\}$ and $\mathbb{P}[Y_i = k] = p_k$. Then

$$p(i,j) =$$

Distribution of $Y_1 + \dots + Y_i$ is given by the i-fold convolution, which is hard to work with.

Instead, we study one particular quantity

and develop a method to establish if $q(i) = 1$ or $q(i) < 1$

Galton - Watson Branching Process

Denote by the expected number of offsprings for each individual ,

Then

$$E_i[X_n] =$$

and by the Markov property

$$E_i[X_n | X_{n-1} = k] =$$

Thus

$$E_i[X_n] =$$

=

But $E_i[X_n] =$

Galton - Watson Branching Process

(1) If $\mu < 1$, then for any i the population gets extinct almost surely

$$q(i) =$$

- $\mathbb{P}_i[X_n \geq 1] \leq i\mu^n \Rightarrow \lim_{n \rightarrow \infty} \mathbb{P}_i[X_n \geq 1] = 0 \Rightarrow$

- $\mathbb{P}_i[X_n = 0 \text{ for some } n] =$

$$(X_n = 0 \Rightarrow X_{n+1} = 0) \xrightarrow{\text{monotonicity}}$$

$$\mathbb{P}_i\left[\bigcup_{n=0}^{\infty} \{X_n = 0\}\right] = .$$

Def 15.1 Let $(X_n)_{n \geq 0}$ be a branching process with offspring distribution p_0, p_1, \dots and mean μ . We call (X_n) subcritical if $\mu < 1$; critical if $\mu = 1$; supercritical if $\mu > 1$.

Galton - Watson Branching Process

Subcritical GW branching process gets extinct with

probability 1. What about the super-/critical regime?

Observation: denote $q_n(i) = \mathbb{P}_i[X_n=0]$

(2) $q_n(i) =$

$X_n=0$ iff for each of i


independent subprocesses $X_n^{(j)}=0$

We saw that

$$q(i) := \mathbb{P}_i[\exists n : X_n=0] =$$

$$q(i) = (q(1))^i \text{ and it is enough to compute } q(1) =: q.$$

Q: How to compute $q = \mathbb{P}_i[\exists n : X_n=0]$?



Probability generating function

Def Let Y be a random variable with values in $\{0, 1, 2, \dots\}$. We call the function

$$\varphi_Y(s) := E[s^Y] = \sum_{k=0}^{\infty} s^k P[Y=k]$$

The probability generating function of Y .

Properties:

(1) $\varphi_Y(s)$ is analytic on $(-1, 1)$; $\varphi_Y^{(n)}(0) = n! P[Y=k]$

(2) $\varphi_Y(1) = 1$; $\varphi_Y(0) = P[Y=0]$

(3) For $|s| < 1$, $\varphi_Y'(s) = \sum_{k=1}^{\infty} k s^{k-1} P[Y=k]$; if $E[Y] < \infty$, then $\varphi_Y'(1) = E[Y]$

(4) For $|s| < 1$, $\varphi_Y''(s) = \sum_{k=2}^{\infty} k(k-1) s^{k-2} P[Y=k]$; in particular, if $P[Y \geq 2] > 0$, then $\varphi_Y(s)$ is (strictly) convex on $(0, 1)$

Galton - Watson Branching Process

Theorem 15.2 Let $(X_n)_{n \geq 0}$ be a branching process with offspring distribution p_0, p_1, \dots . Let φ be the probability generating function of this distribution $\varphi(s) = \sum_{k=0}^{\infty} p_k s^k$. Then the extinction probability q is given by

$$q =$$

Proof.

(i) $q(i) = q^i$

(ii)

Using first step analysis

$$q = \mathbb{P}_1[\exists n : X_n = 0] =$$

=

Galton - Watson Branching Process

(iii) $q \in [0,1]$, $\varphi(1) = 1$

(iv) Let $\hat{q} = \min\{s \in [0,1] : \varphi(s) = s\}$. Then $\forall n q_n \leq \hat{q}$

Induction:

Suppose $q_{n-1} \leq \hat{q}$. Then

$$q_n = P_1[X_n=0] =$$

=

Thus $q_n \leq$

mathematical induction $q_n \leq \hat{q}$ for all $n \in \mathbb{N}$

By the principle of

(v) $q = \varphi(q)$ and $q \in [0,1] \Rightarrow$

$| \Rightarrow$

(vi) $\forall n q_n \leq \hat{q} \Rightarrow$

Galton - Watson Branching Process

Q: When does $q < 1$? When does $s = \varphi(s)$ for $s \in [0, 1]$?

Remark If $p_1 = 1$, then $P_i[X_n = 1] = 1$.

Corollary 15.3 Suppose $p_1 \neq 1$. Then $q = 1$ if the process is critical or subcritical, and $q < 1$ if the process is supercritical.

Proof. Subcritical: discussed before.

Supercritical: $\mu > 1$. Denote $f(s) = \varphi(s) - s$. Then

- $f'(1) = \varphi'(1) - 1 = \mu - 1 > 0$
- $f(0) = p_0$, $f(1) = \varphi(1) - 1 = 0$
- f is continuous on $[0, 1]$



Galton - Watson Branching Process

Critical: $\mu = 1$

- $\mu = 1 \Rightarrow p_0 \neq 1$ (otherwise $\mu = \sum_{k=0}^{\infty} k p_k = 0$)
 - $\mu = 1 \Rightarrow p_0 \neq 0$ (otherwise $\sum_{k=1}^{\infty} k p_k = 1 \Rightarrow p_1 = 1$)
 - $p_1 \neq 1 \Rightarrow \sum_{k=2}^{\infty} p_k > 0$ (otherwise $\mu = \sum_{k=0}^{\infty} k p_k = 0 \cdot p_0 + 1 \cdot p_1 = p_1 < 1$)
 - $\varphi'(1) = 1$
 - if $t \in (0, 1)$, then
- $\varphi'(t) = \sum_{k=1}^{\infty} k t^{k-1} p_k =$
- Take any $s \in (0, 1)$, $\int_s^1 \varphi'(t) dt =$
 \Rightarrow

