

MATH180C: Introduction to Stochastic Processes II

<https://mathweb.ucsd.edu/~ynemish/teaching/180c>

Today: Introduction. Birth processes

Next: PK 6.2-6.3

Week 1:

- visit course web site
- join Piazza

Stochastic (random) processes

Def. Let (Ω, \mathcal{F}, P) be a probability space.

Stochastic process is a collection $(X_t : t \in T)$ of random variables $X_t : \Omega \rightarrow S \subset \mathbb{R}$ (all defined on the same probability space)

- often t represents time, but can be 1-D space
- T is called the index set, S is called the state space
- $X : \Omega \times T \rightarrow S$ ($X_t(\omega) \in S$)
- for any fixed ω , we get a realization of all random variables $(X_t(\omega) : t \in T) \leftarrow$ sample path trajectory
- stochastic process $X_{\cdot}(\omega) : T \rightarrow S$ — random function

Stochastic processes. Classification

Questions:

- What is T
- What is S
- Relations between X_{t_1} and X_{t_2} for $t_1 \neq t_2$?
- Properties of the trajectory

Discrete time

$T = \mathbb{N}, \mathbb{Z}, \text{finite set}$

↑ random vector

Continuous time

$T = \mathbb{R}, [0, +\infty), [0, 1]$

Real-valued

$S = \mathbb{R}$

Integer-valued

$S = \mathbb{Z}$


Nonnegative ...

$S \subset [0, +\infty)$

Continuous, right-continuous (cadlag) sample path

Examples of stochastic processes

- Gaussian processes: for any $t \in T$, X_t has normal distrib.
- Stationary processes: distribution doesn't change in time
- Processes with stationary and independent increments (Levy)
- Poisson process: increments are independent and Poisson(-)
- **Markov processes**: "distribution in the future depends only on the current state, but does not depend on the past"




Examples of stochastic processes

- **Martingales** : $\mathbb{E}[X_{n+1} | X_n, \dots, X_1, X_0] = X_n$ ("fair game")

- **Brownian motion** (Wiener process) is a continuous-time st. proc.

Gaussian, martingale, has stationary and independent increments, Markov, $\text{Var}[W_t] = t$
 $\text{Cov}[W_t, W_s] = \min\{s, t\}$, its sample path is everywhere continuous and nowhere differentiable

- diffusion processes (stochastic differential equations)



- ...

Continuous time MC

Continuous Time Markov Chains

Def (Discrete-time Markov chain)

Let $(X_n)_{n \geq 0}$ be a discrete time stochastic process taking values in $\mathbb{Z}_+ = \{0, 1, 2, \dots\}$ (for convenience). $(X_n)_{n \geq 0}$ is called Markov chain if for any $n \in \mathbb{N}$ and $i_0, i_1, \dots, i_{n-1}, i, j \in \mathbb{Z}_+$

$$P(X_{n+1} = j \mid X_0 = i_0, X_1 = i_1, \dots, X_{n-1} = i_{n-1}, X_n = i) = P(X_{n+1} = j \mid X_n = i)$$

Def (Continuous-time Markov chain)

Let $(X_t)_{t \geq 0} = (X_t : 0 \leq t < \infty)$ be a continuous time process taking values in \mathbb{Z}_+ . $(X_t)_{t \geq 0}$ is called Markov chain if for any $n \in \mathbb{N}$, $0 \leq t_0 < t_1 < \dots < t_{n-1} < s, t > 0$, $i_0, i_1, \dots, i_{n-1}, i, j \in \mathbb{Z}_+$

$$P(X_{t+s} = j \mid X_{t_0} = i_0, X_{t_1} = i_1, \dots, X_{t_{n-1}} = i_{n-1}, X_s = i)$$

Markov property

$$= P(X_{t+s} = j \mid X_s = i)$$

Poisson process


Def. A Poisson process of intensity (rate) $\lambda > 0$ is an integer-valued stochastic process $(X_t)_{t \geq 0}$ for which

1) for each time points $t_0 = 0 < t_1 < \dots < t_n$, the process increments $X_{t_1} - X_{t_0}, X_{t_2} - X_{t_1}, \dots, X_{t_n} - X_{t_{n-1}}$

are independent random variables

2) for $s \geq 0$ and $t > 0$, the random variable $X_{s+t} - X_s$ has the Poisson distribution $\mathbb{P}[X_{s+t} - X_s = k] = \frac{(\lambda t)^k}{k!} e^{-\lambda t}$, $k = 0, 1, \dots$

3) $X_0 = 0$



Example: Poisson process as MC

Is Poisson process a continuous time MC?

Poisson process:

✓ continuous time

✓ discrete state

(*)

Let $(X_t)_{t \geq 0}$ be a Poisson process, let $i_0 \leq i_1 \leq \dots \leq i_{n-1} \leq i \leq j$

$$\begin{aligned} & P(X_{s+t} = j \mid X_{t_0} = i_0, X_{t_1} = i_1, \dots, X_{t_{n-1}} = i_{n-1}, X_s = i) \quad X_{t+s} - X_s = j - i \\ &= \frac{P(X_{t_0} = i_0, X_{t_1} - X_{t_0} = i_1 - i_0, X_{t_2} - X_{t_1} = i_2 - i_1, \dots, X_s - X_{t_{n-1}} = i - i_{n-1})}{P(X_{t_0} = i_0, \dots, X_s - X_{t_{n-1}} = i - i_{n-1})} \\ &= P(X_{t+s} - X_s = j - i) \\ &= P(X_{t+s} = j \mid X_s = i) \end{aligned}$$