

# MATH180C: Introduction to Stochastic Processes II

<https://mathweb.ucsd.edu/~ynemish/teaching/180c>

Today: Martingales

Next: PK 2.5, Durrett 5.1-5.2

Week 8:

- HW6 due Monday, May 22 on Gradescope
- Midterm 2 on Wednesday, May 24

# Martingales

Definition. A stochastic process  $(X_n, n \geq 0)$  is a martingale if for  $n = 0, 1, \dots$

(a)

(b)

After taking the expectation of both sides of (b) we get that

$(X_n)_{n \geq 0}$  is a martingale  $\Rightarrow$

- submartingale :
- supermartingale :

## Examples of martingales

(i) Let  $X_1, X_2, \dots$  be independent RV's with  $E(|X_k|) < \infty$   
and  $E(X_k) = 0$ . Define  $S_n = X_1 + \dots + X_n$ ,  $S_0 = 0$ .

Then

$\Rightarrow$

(ii) Let  $X_1, X_2, \dots$  be independent RV with  $X_k \geq 0$ ,  $E(|X_k|) < \infty$   
and  $E(X_k) = 1$ . Define  $M_n = X_1 X_2 \dots X_n$ ,  $M_0 = 1$ .

Then

$\Rightarrow$

## Example

### Stock prices in a perfect market

Let  $X_n$  be the closing price at the end of day  $n$  of a certain publicly traded security such as a share or stock. Many scholars believe that in a perfect market these price sequences should be martingales. (see PK page 73 for more details).

## History and gambling

Let  $(X_n)_{n \geq 0}$  be a stochastic process describing your total winnings in  $n$  games with unit stake.

Think of  $X_n - X_{n-1}$  as your net winnings per unit stake in game  $n$ ,  $n \geq 1$ , in a series of games, played at times  $n=1, 2, \dots$ .

In the martingale case

Some early works of martingales was motivated by gambling. Note that there exists a betting strategy called the "martingale system"  $\leftarrow$  doubling bets after losses

## Some basic properties

Let  $(X_n)_{n \geq 0}$  be a martingale.

•

Proof

Exercise:

• Markov inequality: If  $X_n \geq 0 \quad \forall n$ , then for any  $\lambda > 0$

$\Rightarrow$

## Maximal inequality for nonnegative martingales

Thm. Let  $(X_n)_{n \geq 0}$  be a martingale with nonnegative values.

For any  $\lambda > 0$  and  $m \in \mathbb{N}$

(1)

and

(2)

Proof. We prove (1), (2) follows by taking the limit  $m \rightarrow \infty$ .

Take the vector  $(X_0, X_1, \dots, X_m)$  and partition the sample space wrt the index of the first r.v. rising above  $\lambda$

Compute

using the above partition

## Proof of the maximal inequality

$$E(X_m) =$$

$\geq$

Compute  $E(X_m \mathbb{1}_{X_0 < \lambda, \dots, X_{n-1} < \lambda, X_n \geq \lambda})$  by conditioning on

$X_0, X_1, \dots, X_{n-1}, X_n$ :

$$E(X_m \mathbb{1}_{X_0 < \lambda, \dots, X_{n-1} < \lambda, X_n \geq \lambda})$$

=

=

=

Sum for all  $n$

$$E(X_m) \geq$$



## Example

A gambler begins with a unit amount of money and faces a series of independent fair games. In each game the gambler bets fraction  $p$  of his current fortune, wins with probability  $\frac{1}{2}$ , loses with probability  $\frac{1}{2}$ . Estimate the probability that the gambler ever doubles the initial fortune.

Denote by  $Z_n, n \geq 0$ , the gambler's fortune after  $n$ -th game.

Denote

Then

## Martingale transform

In the previous example the stake in  $n$ -th game is  $p Z_{n-1}$ . What if we choose another strategy?

Def Let  $(X_n)_{n \geq 0}$  be a nonnegative martingale, and let  $(C_n)_{n \geq 0}$  be a stochastic process with  $C_n = f_n(X_0, \dots, X_{n-1})$ . Then the stochastic process

is called the

- Think of
- $X_k - X_{k-1}$  as the winning per unit stake in  $k$ -th game
  - $C_k$  as your stake in  $k$ -th game  
decision is made based on the previous history
  - $(C \cdot X)_n$  as total winnings up to time  $n$

## Martingale transform

Prop. Let  $Z_n = X_0 + (C \cdot X)_n$ . Let  $C_k > 0$  bounded if  $Z_{k-1} > 0$  and  $C_k = 0$  if  $Z_{k-1} = 0$ . Then  $(Z_n)_{n \geq 0}$  is a martingale

Proof:  $E(Z_{n+1} | Z_0, \dots, Z_n) =$   
=

Note that

If  $Z_n > 0$ , then  $C_1 > 0, \dots, C_n > 0$ ,

and

$$E(Z_{n+1} | Z_0, \dots, Z_n) =$$
  
=

If  $Z_n = 0$ , then  $C_{n+1} = 0$  and  $E(Z_{n+1} | Z_0, \dots, Z_n) = 0 = Z_n$

## Convergence of nonnegative martingales

Thm.

If  $(X_n)_{n \geq 0}$  is a nonnegative (super)martingale, then  
with probability 1

and

Example

An urn initially contains one red ball and one green ball. Choose a ball and return it to the urn together with another ball of the same color. Repeat. Denote by  $X_n$  the fraction of red ball after  $n$  iterations.

## Example (cont.)

(i)  $(X_n)_{n \geq 0}$  is a martingale

Denote by  $R_n$  the number of red balls after  $n$ -th iteration

$$R_n =$$

Then

$$E(X_{n+1} | X_0, \dots, X_n) =$$

=

(ii)  $X_n$  is nonnegative  $\Rightarrow$

(iii) Compute the distribution of  $X_\infty$

$$P(X_n = \frac{k}{n+2}) = \frac{1}{n+1} \quad \text{for } k \in \{1, 2, \dots, n+1\}$$

$$P(X_\infty \leq x) = x, \quad x \in (0, 1) \Rightarrow X_\infty \sim \text{Unif}(0, 1)$$