

MATH180C: Introduction to Stochastic Processes II

<https://mathweb.ucsd.edu/~ynemish/teaching/180c>

Today: Brownian motion

Next: PK 8.3- 8.4

Week 10:

- homework 8 (due Friday, June 9)

Reflection principle

Thm. Let $(B_t)_{t \geq 0}$ be a standard BM. Then

for any $t \geq 0$ and $x > 0$

$$\cancel{(S_t)_{t \geq 0} \stackrel{(d)}{=} (B_t)_{t \geq 0}}$$

$$P\left(\overbrace{\max_{0 \leq u \leq t} B_u}^{S_t} \geq x\right) = P(|B_t| \geq x) = 2 \cdot P(B_t \geq x)$$

Let $\tau_x = \min\{t : B_t = x\}$.

Thm. $F_{\tau_x}(t) = \sqrt{\frac{2}{\pi}} \int_{x/\sqrt{t}}^{\infty} e^{-\frac{v^2}{2}} dv$, $f_{\tau_x}(t) = \frac{x}{\sqrt{2\pi}} t^{-3/2} e^{-\frac{x^2}{2t}}$

Zeros of BM

Denote by $\theta(t, t+s)$ the probability that $B_u = 0$ on $(t, t+s)$

$$\theta(t, t+s) :=$$

Thm. For any $t, s > 0$

$$\theta(t, t+s) =$$

Proof Compute $P(B_u = 0 \text{ for some } u \in (t, t+s])$ by conditioning on the value of B_t .

$$\theta(t, t+s) =$$

(*)

Define $\tilde{B}_u = B_{t+u} - B_t$. Then

$$P(B_u = 0 \text{ on } (t, t+s] \mid B_t = x) =$$

(**)

Zeros of BM

Plugging **(**)** into **(*)** gives

$$\begin{aligned}\Theta(t, t+s) &= \int_{-\infty}^{+\infty} P(B_u = x \text{ for some } u \in (0, s]) \frac{1}{\sqrt{2\pi t}} e^{-\frac{x^2}{2t}} dx \\ &= \int_0^{+\infty} P(B_u = x \text{ for some } u \in (0, s]) \frac{1}{\sqrt{2\pi t}} e^{-\frac{x^2}{2t}} dx \\ &\quad + \int_0^{\infty} P(B_u = -x \text{ for some } u \in (0, s]) \frac{1}{\sqrt{2\pi t}} e^{-\frac{x^2}{2t}} dx \\ &= \end{aligned}$$

Finally, $P(B_u = x > 0 \text{ for some } u \in (0, s]) =$

$$\textbf{(**)} = \int_0^{\infty} \sqrt{\frac{2}{\pi t}} e^{-\frac{x^2}{2t}} \left(\int_0^s \frac{x}{\sqrt{2\pi}} y^{-3/2} e^{-\frac{x^2}{2y}} dy \right) dx =$$

Zeros of BM

$$\int_0^{\infty} x e^{-\frac{x^2}{2} \left(\frac{1}{t} + \frac{1}{y} \right)} dx =$$

$$\Rightarrow (*) =$$

Now use the change of variable $z = \sqrt{\frac{y}{t}}$, $dy = 2t dz$

$$\begin{aligned} (*) &= \frac{\sqrt{t}}{\pi} \int_0^{\sqrt{s/t}} \frac{1}{t(1+z^2)\sqrt{t}z} \cdot 2t dz = \frac{2}{\pi} \int_0^{\sqrt{s/t}} \frac{1}{1+z^2} dz = \frac{2}{\pi} \arctan\left(\sqrt{\frac{s}{t}}\right) \\ &= \frac{2}{\pi} \arccos\left(\sqrt{\frac{t}{s+t}}\right) \end{aligned}$$

↑ exercise

Remark Let $T_0 := \inf\{t > 0 : B_t = 0\}$. Then $P(T_0 = 0) = 1$

There is a sequence of zeros of $B_t(\omega)$ converging to 0.

To understand the structure of the set of zeros \rightarrow Cantor set

Behavior of BM as $t \rightarrow \infty$

Thm. Let $(B_t)_{t \geq 0}$ be a (standard) BM. Then

$$P\left(\sup_{t \geq 0} B_t = +\infty, \inf_{t \geq 0} B_t = -\infty\right) = 1$$


(BM "oscillates with increasing amplitude")

Proof. Denote $Z = \sup_{t \geq 0} B_t$. Then for any $c > 0$

$$cZ =$$

By property (iii), cB_{t/c^2} is a standard BM, so cZ has the same distribution as $Z \Rightarrow P(Z=0) = p, P(Z=\infty) = 1-p$

$$p = P(Z=0)$$

$\Rightarrow P(Z=0) = 0, P(Z=\infty) = 1$. Similarly for $\inf_{t \geq 0} B_t$ 

Sample paths of $(B_t)_t$ are not differentiable

Thm. $P(B_t \text{ is not differentiable at zero}) = 1$

Proof. $P(\sup_{t \geq 0} B_t = \infty, \inf_{t \geq 0} B_t = -\infty) = 1. \quad (\star)$

Consider $\tilde{B}_t = t B_{1/t}$. $(\tilde{B}_t)_{t \geq 0}$ is a BM (by property (iv))

By (\star) , for any $\varepsilon > 0 \exists t < \varepsilon, s < \varepsilon$ such that

$\tilde{B}_t > 0, \tilde{B}_s < 0 \Rightarrow$ only differentiable if $\tilde{B}'_0 = 0$

But if $\tilde{B}'_0 = 0$, then

for some $t > 0$ and all $0 < s < t$,

which implies that

for all $0 < s < t$, which

contradicts to (\star) \blacksquare

Thm $P((B_t)_{t \geq 0} \text{ is nowhere differentiable}) = 1$

Reflected BM

Def. Let $(B_t)_{t \geq 0}$ be a standard BM. The stochastic

process

$$|B_t| = \begin{cases} B(t), & \text{if } B(t) \geq 0 \\ -B(t), & \text{if } B(t) < 0 \end{cases}$$

is called reflected BM.

Think of a movement in the vicinity of a boundary.

Moments: $E(R_t) =$

$$\text{Var}(R_t) = E(B_t^2) - (E(|B_t|))^2 =$$


Transition density: $P(R_t \leq y | R_0 = x) =$

=

$$\Rightarrow P_t(x, y) =$$


Thm (Lévy) Let $M_t = \max_{0 \leq u \leq t} B_u$. Then $(M_t - B_t)_{t \geq 0}$ is a reflected BM.

Reflected BM



Brownian bridge

Brownian bridge is constructed from a BM by conditioning on the event $\{B(0)=0, B(1)=0\}$.



Thm 1. Brownian bridge is a continuous Gaussian process on $[0,1]$ with mean 0 and covariance function $\Gamma(s,t) =$

Brownian motion with drift

Def Let $(B_t)_{t \geq 0}$ be a standard BM. Then for $\mu \in \mathbb{R}$ and $\sigma > 0$ the process $(X_t)_{t \geq 0}$ with $X_t =$, $t \geq 0$ is called the Brownian motion with drift μ and variance parameter σ^2 .


Remark BM with drift μ and variance parameter σ is a stochastic process $(X_t)_{t \geq 0}$ satisfying

- 1) $X_0 = 0$, $(X_t)_{t \geq 0}$ has continuous sample paths
- 2) $(X_t)_{t \geq 0}$ has independent increments
- 3) For $t > s$ $X_t - X_s \sim$

In particular, $X_t \sim$

$\Rightarrow X_t$ is not centered,
not symmetric w.r.t. the origin

Brownian motion with drift



Gambler's ruin problem for BM with drift

Let $(X_t)_{t \geq 0}$ be a BM with drift $\mu \in \mathbb{R}$ and variance parameter $\sigma^2 > 0$. Fix $a < x < b$ and denote

$T = T_{ab} = \min\{t \geq 0 : X_t = a \text{ or } X_t = b\}$, and

$u(x) = P(X_T = b \mid X_0 = x)$.

Theorem.

(i) $u(x) =$

(ii) $E(T_{ab} \mid X_0 = x) =$

No proof

Example

Fluctuations of the price of a certain share is modeled by the BM with drift $\mu = 1/10$ and variance $\sigma^2 = 4$. You buy a share at 100\$ and plan to sell it if its price increases to 110\$ or drops to 95\$.

- (a) What is the probability that you will sell at profit?
(b) What is the expected time until you sell the share?

Denote by $(X_t)_{t \geq 0}$ a BM with drift $\frac{1}{10}$ and variance 4,
 $x =$, $b =$, $a =$. Then $2\mu/\sigma^2 =$ and

(a) $P(X_T = 110 | X_0 = 100) =$

(b) $E(T | X_0 = 100) =$

Maximum of a BM with negative drift

Thm Let $(X_t)_{t \geq 0}$ be a BM with drift $\mu < 0$, variance σ^2 and $X_0 = 0$. Denote $M = \max_{t \geq 0} X_t$. Then

Proof. $X_0 = 0$, therefore $M \geq 0$. For any $b > 0$

$$P(M > b) =$$

=

=

$$P(M > b) =$$

Geometric BM

Def. Stochastic process $(Z_t)_{t \geq 0}$ is called a geometric Brownian motion with drift parameter d and variance σ^2 if $X_t = \frac{Z_t}{Z_0}$ is a BM with drift $\mu = d - \frac{1}{2}\sigma^2$ and variance σ^2 .

In other words, $Z_t = z_0 e^{(d - \frac{1}{2}\sigma^2)t + \sigma B_t}$, where $(B_t)_{t \geq 0}$ is a standard BM and $z > 0$ is the starting point $Z_0 = z$.

If $0 \leq t_1 < t_2 < \dots < t_n$, then $\frac{Z_{t_i}}{Z_{t_{i-1}}}$

Since B has independent increments

$\frac{Z_{t_1}}{Z_{t_0}}, \frac{Z_{t_2}}{Z_{t_1}}, \dots, \frac{Z_{t_n}}{Z_{t_{n-1}}}$ are independent and

$$\frac{Z_{t_n}}{Z_{t_0}} =$$

← "relative change of price = product of independent relative changes"

Expectation of Geometric BM

Let $(Z_t)_{t \geq 0}$ be geometric BM with parameters α and σ .

Then

$$E(Z_t | Z_0 = z) =$$

$$E(e^{\sigma B_t}) =$$

$$\Rightarrow E(Z_t | Z_0 = z) = z e^{(\alpha - \frac{1}{2}\sigma^2)t} e^{t \frac{\sigma^2}{2}} =$$

Remark

It can be shown that for $0 < \alpha < \frac{1}{2}\sigma^2$ $Z_t \rightarrow 0$ as $t \rightarrow \infty$

At the same time, for $\alpha > 0$ $E(Z_t) \rightarrow \infty$.

Variance of geometric BM

$$E(Z_t^2 | Z_0 = z) =$$

=

$$\text{Var}(Z_t | Z_0 = z) =$$

Theorem

Let $(Z_t)_{t \geq 0}$ be geometric BM with parameters α and σ^2 .

Then

$$(i) \quad E(Z_t | Z_0 = z) = z e^{\alpha t}$$

$$(ii) \quad \text{Var}(Z_t | Z_0 = z) = z^2 e^{2\alpha t} (e^{\sigma^2 t} - 1)$$

Gambler's ruin for geometric BM

Let $(Z_t)_{t \geq 0}$ be geometric BM with parameters α and σ^2 .

Let $A < 1 < B$, and denote $T = \min \{ t : \frac{Z_t}{Z_0} = A \text{ or } \frac{Z_t}{Z_0} = B \}$.

Theorem

$$P\left(\frac{Z_T}{Z_0} = B\right) =$$

Example Fluctuations of the price are modeled by a geometric BM with drift $\alpha = 0.1$ and variance $\sigma^2 = 4$. You buy a share at 100\$ and plan to sell it if its price increases to 110\$ or drops to 95\$.

Take $A =$, $B =$, $2\alpha/\sigma^2 =$, $1 - 2\alpha/\sigma^2 =$

$$P(X_T = 110 | X_0 = 100) =$$