

Remark: For any \mathcal{O}_X -module \mathcal{F} and collection

$\{s_i \mid i \in I\} \subset \Gamma(X, \mathcal{F})$, we can define a morphism of

\mathcal{O}_X -modules: $\varphi: \mathcal{O}_X^{\oplus I} \longrightarrow \mathcal{F}$

which on any open $U \subset X$ sends $\sum_{i \in I} f_i \in (\mathcal{O}_X^{\oplus I})(U)$

to $\sum_{i \in I} f_i s_i|_U \in \mathcal{F}(U)$.

We have that \mathcal{F} is generated by $\{s_i \mid i \in I\}$ iff

φ is surjective.

Another notion we need: Zeros of sections of (quasi-)coherent sheaves.

Let \mathcal{F} be a (quasi-)coherent sheaf on a noetherian scheme X and $s \in \Gamma(X, \mathcal{F})$. We define the scheme of zeros of s ,

denoted $Z(s)$, as follows (this is a closed subscheme of X):

s defines a morphism of \mathcal{O}_X -modules

$$s: \mathcal{F}^* \longrightarrow \mathcal{O}_X$$

on any $U \subset X$ open $\mathcal{F}^*(U) \ni l \longmapsto l(s|_U) \in \mathcal{O}_X(U)$
 $l \in \mathcal{F}^*(U) = \text{Hom}_{\mathcal{O}_U}(\mathcal{F}|_U, \mathcal{O}_U) \quad \text{"} \quad \mathcal{O}_U(U)$

The image of s is a coherent \mathcal{O}_X -submodule of \mathcal{O}_X ,

i.e., a coherent sheaf of ideals on X , the associated closed subscheme is, by definition, $Z(s)$.

Claim 1: The support $\text{Supp } Z(s)$ of $Z(s)$ (i.e., the underlying closed subset of $Z(s)$) is the set of points $x \in X$ s.t. the image of $s_x: \mathcal{F}_x^* \rightarrow \mathcal{O}_{X,x}$ is contained

in the maximal ideal $\mathfrak{m}_x \subset \mathcal{O}_{X,x}$.

Proof:
$$s: \mathcal{F}^* \longrightarrow \mathcal{O}_X$$

$$0 \longrightarrow \mathcal{I}_{Z(s)} \longrightarrow \mathcal{O}_X \longrightarrow i_* \mathcal{O}_{Z(s)} \longrightarrow 0$$

where $i: Z(s) \hookrightarrow X$

$$x \in Z(s) \iff (i_* \mathcal{O}_{Z(s)})_x \neq 0$$

$$\iff \mathcal{I}_{Z(s),x} \neq \mathcal{O}_{X,x}$$

$$\iff \mathcal{I}_{Z(s),x} \subset \mathfrak{m}_x \subset \mathcal{O}_{X,x}$$

$$\iff s(\mathcal{F}_x^*) \subset \mathfrak{m}_x.$$

Claim 2: Suppose \mathcal{F} is locally free of finite rank n .

Then, for $x \in X$, \mathcal{F}_x is a free $\mathcal{O}_{X,x}$ -module of rank n .

$$\mathcal{F}_x \cong \mathcal{O}_{X,x}^{\oplus n}$$

$$\Rightarrow \mathcal{F}_x^* = \text{Hom}_{\mathcal{O}_{X,x}}(\mathcal{F}_x, \mathcal{O}_{X,x}) \cong \text{Hom}_{\mathcal{O}_{X,x}}(\mathcal{O}_{X,x}^{\oplus n}, \mathcal{O}_{X,x})$$

$$\cong \mathcal{O}_{X,x}^{\oplus n}$$

The condition $s_x(\mathcal{F}_x^*) \subset \mathcal{M}_x$ means $s(x) \in \mathcal{M}_x \mathcal{F}_x$.

Indeed: if we pass to the quotient by \mathcal{M}_x :

$$\mathcal{F}_x / \mathcal{M}_x \mathcal{F}_x \cong \mathcal{O}_{X,x} / \mathcal{M}_x \cong k(x)$$

$$\text{and } \mathcal{F}_x^* / \mathcal{M}_x \mathcal{F}_x^* \cong \left(\mathcal{F}_x / \mathcal{M}_x \mathcal{F}_x \right)^* \cong \left(k(x) \right)^*$$

$s_x(\mathcal{F}_x^*) \subset \mathcal{M}_x \mathcal{F}_x^* \iff s_x$ induces the 0 linear map on $\left(k(x) \right)^*$

$$\iff s_x = 0 \in k(x) = \mathcal{F}_x / \mathcal{M}_x \mathcal{F}_x$$

$$\iff s(x) \in \mathcal{M}_x \mathcal{F}_x.$$

Note: This fails if \mathcal{F} is not locally free:

e.g.: \mathcal{F} = skyscraper sheaf supported on a proper closed subset of X .

Then $\mathcal{F}^* = \mathcal{O}_X$ the zero sheaf on X

Homogeneous ideals of subsets of projective space:

Assume $S := R[X_0, \dots, X_n]$, R comm. ring

$$X := \text{Proj } S = \mathbb{P}_R^n$$

We saw that $\Gamma(X, \mathcal{O}_X(d)) = S_d \quad \forall d \in \mathbb{Z}$

i.e., $\Gamma_*(\mathcal{O}_X) = S$.

Def: (1) For a subset $Y \subset X$, define

$$I_{Y,d} := \{s \in S_d \mid Y \subset Z(s)\}$$

= the set of homogeneous polynomials of degree d
vanishing on $Y := \{s \in S_d \mid \forall x \in Y, s(x) \in \mathfrak{m}_x \text{ (} \mathbb{Q}_x(d) \cong \mathfrak{m}_x \text{)}\}$

(2) For a closed subscheme $Y \subset X$ with ideal sheaf \mathcal{I}_Y ,

define $I_{Y,d} := \{s \in S_d \mid \mathcal{I}_Z(s) \subset \mathcal{I}_Y\}$
 $= \{s \in S_d \mid Y \subset Z(s) \text{ as subschemes}\}$

Ex: If Y is reduced, the two definitions are equal.

Def: The homogeneous ideal of a subset or closed subscheme $Y \subset X = \mathbb{P}_R^n$ is

$$I_Y := \bigoplus_{d \in \mathbb{Z}} I_{Y,d} \subset S$$

The homogeneous coordinate ring of Y is $S(Y) := S / I(Y)$.

Remark: Choose $s \in S_d$ and suppose $Y \subset X$ is a closed subscheme. We have $s \in I_{Y,d}$ iff $\mathcal{I}_Z(s) \subset \mathcal{I}_Y$.

$\mathcal{I}_Z(s)$ is the image of $s: \mathcal{O}_X(d)^* \rightarrow \mathcal{O}_X$

recall that $\mathcal{O}_X(d)^* \cong \mathcal{O}_X(-d)$, so

$\mathcal{I}_Z(s)$ is the image of $s: \mathcal{O}_X(-d) \rightarrow \mathcal{O}_X$

$\mathcal{I}_Z(s) \subset \mathcal{I}_Y \iff s: \mathcal{O}_X(-d) \rightarrow \mathcal{O}_X$
 $\searrow \quad \swarrow$
 \mathcal{I}_Y factors

twist by $\mathcal{O}_X(d)$: $\mathcal{O}_X(-d) \otimes \mathcal{O}_X(d) \xrightarrow{s} \mathcal{O}_X(d)$

\iff
 $\mathcal{O}_X \xrightarrow{s} \mathcal{O}_X(d)$
 $\searrow \quad \nearrow$
 $\mathcal{I}_Y(d)$

$\searrow \quad \nearrow$
 $\mathcal{I}_Y(d)$

Fact: $\mathcal{O}_X \xrightarrow{s} \mathcal{O}_X(d)$ is "multiplication" by s :

$$\forall U \subset X \quad \begin{array}{ccc} \mathcal{O}_X(U) & \xrightarrow{s} & \mathcal{O}_X(d)(U) \\ \downarrow \varphi & \longmapsto & \downarrow \varphi \circ s|_U \in \mathcal{O}_X(d)(U) \end{array}$$

s factors through $\mathcal{I}_Y(d) \iff s \in \Gamma(X, \mathcal{I}_Y(d))$

More generally: $\forall \mathcal{O}_X$ -module \mathcal{F} ,

$$\Gamma(X, \mathcal{F}) \stackrel{\circ s}{=} \text{Hom}_{\mathcal{O}_X}(\mathcal{O}_X, \mathcal{F})$$

$$\longmapsto \left(\varphi \longmapsto \varphi \circ s|_U \text{ on any } U \right)$$

$$\varphi(1) \longleftarrow \varphi$$

$$\Rightarrow s \in \Gamma(X, \mathcal{I}_Y(d)) \iff s: \mathcal{O}_X \rightarrow \mathcal{I}_Y(d)$$

Conclusion: $s \in \mathcal{I}_{Y,d} \iff s \in \Gamma(X, \mathcal{I}_Y(d))$.

$$\text{i.e., } I_{Y,d} = \Gamma(X, \mathcal{I}_Y(d))$$

Note that $\mathcal{I}_Y \hookrightarrow \mathcal{O}_X \Rightarrow \mathcal{I}_Y(d) \hookrightarrow \mathcal{O}_X(d)$

$$\Rightarrow \Gamma(\mathcal{I}_Y(d)) \hookrightarrow \Gamma(\mathcal{O}_X(d)) \hookrightarrow I_{Y,d}$$

$$\begin{aligned} \Rightarrow I_Y &= \bigoplus_{d \in \mathbb{Z}} I_{Y,d} = \bigoplus_{d \in \mathbb{Z}} \Gamma(\mathcal{I}_Y(d)) \\ &=: \Gamma_* (\mathcal{I}_Y) \end{aligned}$$

Recall: If S is finitely generated by S_1 as an S_0 -alg. and \mathcal{F} is quasi-coherent, then $\widetilde{\Gamma_* (\mathcal{F})} \xrightarrow{\sim} \mathcal{F}$

This is true for $S = R[X_0, \dots, X_n]$ and $\mathcal{F} = \mathcal{I}_Y$,

$$\text{so } \widetilde{\Gamma_* (\mathcal{I}_Y)} \xrightarrow{\sim} \mathcal{I}_Y, \text{ i.e., } \widetilde{I_Y} \xrightarrow{\sim} \mathcal{I}_Y.$$

Main Lemma: There is a natural isomorphism

$$g: Y \xrightarrow{\cong} \text{Proj } S(Y)$$

Proof: The quotient morphism $\varphi: S \longrightarrow S(Y) = S/I(Y)$

induces a morphism $f: \text{Proj } S(Y) \longrightarrow \text{Proj } S$.

(f is defined everywhere because φ is surjective)

We will show that f factors through an isom.

$$\text{Proj } S(Y) \xrightarrow{\cong} Y \hookrightarrow X := \text{Proj } S$$

We have the exact sequence

$$0 \longrightarrow I_Y \longrightarrow S \longrightarrow S(Y) \longrightarrow 0 \text{ (by def.)}$$

Lemma 1: The \sim functor is exact.