

Equidistribution of affine random walks on some nilmanifolds.

Based on joint works with Weikun He and Elon Lindenstrauss

Tsviqa Lakrec

Hebrew University of Jerusalem

- 1 Random walks on tori and Heisenberg nilmanifold
- 2 Quantitative statement
- 3 More general statement

Random walk associated to a group action

Consider an action $G \curvearrowright X$.

Let $\mu \in \mathcal{P}(G)$ be a probability measure. Let $x \in X$.

Definition

The **random walk** on X induced by μ and starting at x is the sequence of random variable $(g_n g_{n-1} \cdots g_1 x)_{n \geq 1}$ where $(g_n)_{n \geq 1}$ is i.i.d. of law μ .

The law of $g_n g_{n-1} \cdots g_1 x$ is $\mu^{*n} * \delta_x$, i.e. for any function $f: X \rightarrow \mathbb{C}$,

$$\mathbb{E}[f(g_n g_{n-1} \cdots g_1 x)] = \int_X f d(\mu^{*n} * \delta_x).$$

We are interested in the convergence in law, i.e. the convergence of $\mu^{*n} * \delta_x$ in the weak-* topology on X .

Let $X = N/\Lambda$ be a **compact nilmanifold**. That is,

- 1 N is a connected simply-connected nilpotent Lie group,
- 2 $\Lambda \subset N$ is a lattice in N , i.e. it is discrete in N and
- 3 the Haar measure on N induces a finite N -invariant measure on N/Λ .

We renormalize this measure to be a probability measure and denote it by m_X .

Group of affine transformations

On $X = N/\Lambda$, we consider the action of its automorphism group

$$\text{Aut}(X) = \{ \gamma \in \text{Aut}(N) \mid \gamma(\Lambda) = \Lambda \}$$

and that of its affine transformations

$$\text{Aff}(X) = \text{Aut}(X) \ltimes N,$$

Here, for $\gamma \in \text{Aut}(X)$ and $n \in N$, $(\gamma, n) \in \text{Aut}(X) \ltimes N$ is the map

$$\forall x \in N, \quad x\Lambda \mapsto n\gamma(x)\Lambda.$$

For $g = (\gamma, n)$, we call γ the automorphism part and denote $\theta(g) = \gamma$.

Examples

- 1 Let $d \geq 1$, $N = \mathbb{R}^d$ and $\Lambda = \mathbb{Z}^d$. Then $X = \mathbb{T}^d = \mathbb{R}^d / \mathbb{Z}^d$,
 $\text{Aut}(X) = \text{GL}_d(\mathbb{Z})$.
- 2 Let N be the Heisenberg group, $N = \mathbb{R}^3$ endowed with the law

$$(x, y, z) \cdot (x', y', z') = (x + x', y + y', z + z' + xy').$$

Let $\Lambda = \{ (x, y, z) \in N \mid x, y, z \in \mathbb{Z} \}$.

Then $\text{Aut}(X) = \text{GL}_2(\mathbb{Z}) \ltimes \mathbb{Z}^2$ is the set of transformations

$$(x, y, z) \mapsto (ax+by, cx+dy, \det(g)(z - \frac{xy}{2}) + \frac{1}{2}(ax+by)(cx+dy) + \alpha x + \beta y)$$

where $g = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \text{GL}_2(\mathbb{Z})$ and $(\alpha, \beta) \in \frac{1}{2}\mathbb{Z}^2$ satisfies some parity condition.

Question

Let X be a compact nilmanifold. Given $\mu \in \mathcal{P}(\text{Aff}(X))$ and $x \in X$,

- 1 does $\mu^{*n} * \delta_x \xrightarrow{*} m_X$?
- 2 If it does, how fast is the convergence?

Expected answer : Yes, unless there is obvious obstruction.

Remark

Let $H = \langle \text{Supp}(\mu) \rangle$, the subgroup generated by the support.
If $\overline{Hx} \neq X$, then $\mu^{*n} * \delta_x \not\xrightarrow{*} m_X$.

Note that $\text{Aff}(X) \curvearrowright X$ is transitive. Hence $X = \text{Aff}(X)/\text{Aut}(X) \ltimes \Lambda$ is a homogeneous space. Let \mathfrak{g} be the Lie algebra of $\text{Aff}(X)$.

Theorem (Benoist-Quint)

Let $H \subset \text{Aff}(X)$ be a subgroup and $x \in X$.

Assume that the Zariski closure of $\text{Ad}(H)$ in $\text{GL}(\mathfrak{g})$ is semisimple without compact factor.

Then \overline{Hx} is a finite homogeneous union of affine submanifolds.

Orbit closures, case of a torus

Theorem (Guivarc'h-Starkov & Muchnik)

Let Γ be a subgroup of $GL_d(\mathbb{Z}) = \text{Aut}(\mathbb{T}^d)$. Assume

- 1 the action of the Γ on \mathbb{Q}^d is strongly irreducible.
- 2 the Zariski closure Γ in $GL_d(\mathbb{R})$ is semisimple without compact factor.

For every $x \in \mathbb{T}^d$, the orbit Γx is either finite or dense.

Definition

We say Γ acts **strongly irreducibly** on \mathbb{Q}^d if it does not preserve any finite nontrivial union of proper subspaces of \mathbb{Q}^d .

Theorem (Bourgain-Furman-Lindenstrauss-Mozes, He-Saxcé)

Let $\mu \in \mathcal{P}(\mathrm{GL}_d(\mathbb{Z}))$ having a finite β -exponential moment for some $\beta > 0$. Let $\Gamma = \langle \mathrm{Supp}(\mu) \rangle$. Assume that the action of the Γ on \mathbb{R}^d is strongly irreducible.

Then for any $x \in \mathbb{T}^d$, $\mu^{*n} * \delta_x \xrightarrow{*} m_{\mathbb{T}^d}$ unless $x \in \mathbb{Q}^d / \mathbb{Z}^d$ (i.e. Γx is finite).

Definition

For some $\beta > 0$, we say μ *has a finite β -exponential moment* if

$$\int_{\mathrm{SL}_d(\mathbb{Z})} \|g\|^\beta d\mu(g) < +\infty.$$

Recall that $\theta: \text{Aff}(X) \rightarrow \text{Aut}(X)$ denotes the projection.

Theorem (He-Lindenstrauss-L.)

Let $\mu \in \mathcal{P}(\text{GL}_d(\mathbb{Z}) \ltimes \mathbb{R}^d)$ having a finite support.

Let $H = \langle \text{Supp}(\mu) \rangle$ and $\Gamma = \theta(H)$.

Assume Γ satisfies the assumptions in the BFLM theorem.

*Then for any $x \in \mathbb{T}^d$, $\mu^{*n} * \delta_x \rightarrow^* \mathfrak{m}_{\mathbb{T}^d}$ unless Hx is finite.*

Very similar result was obtained by Boyer, under different assumptions.

Equidistribution in law, Heisenberg nilmanifold

Let $X = N/\Lambda$ with N being the $(2k + 1)$ -dimensional Heisenberg group. Note that (N, N) is the one dimensional center and $N/(N, N)\Lambda$ is a $2k$ -dimensional torus.

Denote by $\pi: X \rightarrow N/(N, N)\Lambda$ the projection.

Theorem (H-Lindenstrauss-L.)

Let $\mu \in \mathcal{P}(\text{Aff}(X))$ with a finite support.

Let $H = \langle \text{Supp}(\mu) \rangle$ and $\Gamma = \theta(H)$.

Assume that the action of Γ on $N/(N, N)$ satisfies the assumptions in the BFLM theorem.

Then for any $x \in X$, $\mu^{*n} * \delta_x \rightarrow^* m_X$ unless $\pi(Hx)$ is finite.

If $\mu \in \mathcal{P}(\text{Aut}(X))$, then finite support can be relaxed to having a finite exponential moment.

- 1 Random walks on tori and Heisenberg nilmanifold
- 2 Quantitative statement**
- 3 More general statement

Wasserstein distance

We fix a Riemannian distance on $X = N/\Lambda$.

For $\alpha \in (0, 1)$, let $\mathcal{C}^{0,\alpha}(X)$ denote the space of α -Hölder continuous functions on X , equipped with the norm

$$\|f\|_{0,\alpha} = \|f\|_{\infty} + \sup_{x \neq y \in X} \frac{|f(x) - f(y)|}{d(x,y)^{\alpha}}.$$

Definition

Let ν and η be Borel measures on X . The α -Wasserstein distance between them is

$$\mathcal{W}_{\alpha}(\nu, \eta) = \sup_{f \in \mathcal{C}^{0,\alpha}(X): \|f\|_{0,\alpha} \leq 1} \left| \int_X f \, d\nu - \int_X f \, d\eta \right|.$$

Quantitative statement, Lyapunov exponent

Let $\mu \in \mathcal{P}(\text{Aff}(X))$ with a finite support.

Let $H = \langle \text{Supp}(\mu) \rangle$ and $\Gamma = \theta(H)$.

Definition

Denote by $\lambda_{1,N/(N,N)}(\theta_*\mu)$ the **top Lyapunov exponent** of the random walk induced by $\theta_*\mu$ on the Euclidean space $N/(N, N)$.

$$\lambda_{1,N/(N,N)}(\theta_*\mu) = \lim_{n \rightarrow +\infty} \frac{1}{n} \int \log \|\theta(g)\|_{N/(N,N)} d\mu^{*n}(g)$$

where $\|\cdot\|_{N/(N,N)}$ denotes any operator norm on $\text{End}(N/(N, N))$.

Theorem (He-Lindenstrauss-L.)

Assume $N = \mathbb{R}^d$ or a Heisenberg group. Assume that the action of Γ on $N/(N, N)$ satisfies the assumptions in the BFLM theorem.

Given $\lambda \in (0, \lambda_{1, N/(N, N)}(\mu))$ and $\alpha \in (0, \beta)$, there exists $C \geq 1$ such that :
If for some $x \in X$, $t \in (0, \frac{1}{2})$ and $m \geq C \log \frac{1}{t}$,

$$\mathcal{W}_\alpha(\mu^{*m} * \delta_x, \mathfrak{m}_X) > t,$$

then there exists $x' \in X$ and a finite set $F \subset \text{Aff}(X)$ such that

$$d(x, x') + \max_{g \in \text{Supp}(\mu)} d(g, F) \leq e^{-\lambda m},$$

and the projection of the orbit $\langle F \rangle x'$ in $N/(N, N)\Lambda$ is finite of cardinality less than t^{-C} .

If $\mu \in \mathcal{P}(\text{Aut}(X))$ instead, then this can be reformulated with infinite support instead.

- 1 Random walks on tori and Heisenberg nilmanifold
- 2 Quantitative statement
- 3 More general statement

Growth rate and spectral radius

Let H be a group and μ a probability measure on H .

Definition

Consider an action of H on a group Z by automorphisms. Let $\theta_Z: H \rightarrow \text{Aut}(Z)$ denote the homomorphism. Define

$$\tau_Z(\mu) = \lim_{m \rightarrow +\infty} \frac{1}{m} \log \# \theta_Z(\text{Supp}(\mu^{*m})).$$

This definition needs to be slightly modified when μ is not finitely supported.

Definition

If $(X, m_X) \rightarrow (Y, m_Y)$ is a factor map of a H -spaces ($p.m.p$ action on both), define

$$\sigma_{X,Y}(\mu) = - \lim_{m \rightarrow +\infty} \frac{1}{m} \log \|P(\mu)^m\|_{L^2(X, m_X) \ominus L^2(Y, m_Y)},$$

Definition

Let $\lambda > 0$, $C > 1$, $\alpha \in (0, 1]$, $\mu \in \mathcal{P}(\text{Aut}(X))$, and let $H = \langle \text{Supp}(\mu) \rangle$. We say that the μ -induced random walk on X satisfies **(C, λ, α) -quantitative equidistribution** if the following holds for any integer $m \geq 1$ and any $t \in (0, \frac{1}{2})$. Assume

$$m \geq C \log \frac{1}{t} \quad \text{and} \quad \mathcal{W}_\alpha(\mu^{*m} * \delta_x, \mathfrak{m}_X) > t.$$

Then there exists a point $x' \in X$ such that

- 1 $d(x, x') \leq e^{-\lambda m}$
- 2 $\pi(Hx')$ lies in a proper closed H -invariant subset of T of height $\leq t^{-C}$.

Theorem (He-Lindenstrauss-L.)

Let $\mu \in \mathcal{P}(\text{Aut}(X))$ be with a finite β -exponential moment, Γ as before. Assume that there exists a rational Γ -invariant connected central subgroup $Z \subset N$ such that

$$\tau_Z(\mu) < 2\sigma_{X,Y}(\mu)$$

where $Y = N/(\Lambda Z)$ is the factor nilmanifold obtained by quotienting out Z .

If the μ -induced random walk on Y satisfies (λ, α) -quantitative equidistribution for some $\lambda > 0$ and $\alpha \in (0, \beta]$ then the μ -induced random walk on X satisfies (λ', α) -quantitative equidistribution for any $\lambda' \in (0, \lambda)$.

- 1 N is a step 2 nilpotent group and the action of Γ on its center $Z(N)$ is virtually nilpotent.
- 2 Action of $\Gamma \subset \mathrm{SL}_{2d}(\mathbb{Z})$ on \mathbb{T}^{2d} consisting of $d \times d$ -block triangular matrices. For example, let μ be the law of

$$\left(\begin{array}{c|c} A & I_d \\ \hline 0 & D \end{array} \right).$$

where A and D are independent random variables. Given any A , we can choose D to ensure

$$\tau_{\mathbb{R}^d \oplus 0}(\mu) < 2\sigma_{\mathbb{T}^{2d}, 0 \oplus \mathbb{T}^d}(\mu).$$

Thank you for listening!