

**Math 270B: Numerical Analysis (Part B)**  
**Winter quarter 2023**

**Homework Assignment 5**

**Due: 10:00 am, Wednesday, February 15, 2023.**

1. Given  $q_1, \dots, q_m \in \mathcal{P}$ . Prove that they are linearly independent in  $\mathcal{P}$  if and only if the Gram matrix  $G(q_1, \dots, q_m) = [G_{ij}]_{i,j=1}^m$  is symmetric positive definite, where

$$G_{ij} = \int_a^b p_i(x)p_j(x) dx, \quad i, j = 1, \dots, m.$$

2. Use the Gram-Schmidt orthogonalization to construct the orthogonal polynomials  $Q_0$ ,  $Q_1$ , and  $Q_2$  on  $[0, 1]$  from  $q_0(x) = 1$ ,  $q_1(x) = x$ , and  $q_2(x) = x^2$ .
3. Let  $\{Q_n\}_{n=0}^\infty$  be orthonormal polynomials on  $[a, b]$ . Define  $K_n(x, t) = \sum_{k=0}^n Q_k(x)Q_k(t)$  for all  $n \geq 0$  and all  $x, t \in \mathbb{R}$ . Show that

$$p_n(x) = \int_a^b p_n(t)K_n(x, t) dt \quad \forall p_n \in \mathcal{P}_n \text{ and } x \in \mathbb{R}.$$

4. Let  $u, v \in C^2([0, 1])$  be such that  $u(0) = u(1) = v(0) = v(1) = 0$ . Let  $\lambda, \mu \in \mathbb{R}$  be such that  $\lambda \neq \mu$ . Assume that  $-u'' + u = \lambda u$  and  $-v'' + v = \mu v$  on  $[0, 1]$ . Prove that  $u$  and  $v$  are orthogonal in  $L^2(0, 1)$ .
5. Let  $P_n(x) = (2^n n!)^{-1} [(x^2 - 1)^n]^{(n)}$  ( $n = 0, 1, \dots$ ) be the Legendre polynomials. Let  $r \geq 1$  be an integer. Show that

$$\int_{-1}^1 P_m^{(r)}(x)P_n^{(r)}(x)(1-x^2)^r dx = 0 \quad \text{if } m, n \geq r \text{ and } m \neq n.$$