

SDP and SOS

Eight Exercises for Pablo Parrilo's Lectures
LMIPO, UC San Diego, March 22-23, 2010

1. Consider a univariate polynomial of degree d , that is bounded by one in absolute value on the interval $[-1, 1]$. How large can its leading coefficient be? Give an SOS formulation for this problem, and solve it numerically for $d = 2, 3, 4, 5$. Can you guess what the general solution is as a function of d ? Can you characterize the optimal polynomial?
2. Consider a given univariate rational function $r(x)$, for which we want to find a good polynomial approximation $p(x)$ of fixed degree d on the interval $[-2, 2]$.
 - (a) Write an SOS formulation to compute the best polynomial approximation of $r(x)$ in the supremum norm.
 - (b) Same as before, but now $p(x)$ is also required to be convex.
 - (c) Same as before, but $p(x)$ is required to be a convex lower bound of $r(x)$ (i.e., $p(x) \leq r(x)$ for all $x \in [-2, 2]$).
 - (d) Let $r(x) = \frac{1-2x+x^2}{1+x+x^2}$. Find the solution of the previous subproblems (for $d = 4$), and plot them.
3. Given a set $\mathcal{S} \subseteq \mathbb{R}^n$ that strictly contains the origin, we define the *polar set* $\mathcal{S}^\circ \subseteq \mathbb{R}^n$ as:

$$\mathcal{S}^\circ := \{y \in \mathbb{R}^n \mid y^T x \leq 1, \quad \forall x \in \mathcal{S}\}.$$

- (a) Let \mathcal{S} be the feasible set of an SDP, i.e.,

$$\mathcal{S} = \{x \in \mathbb{R}^m \mid \sum_{i=1}^m x_i A_i \preceq C\},$$

where A_i and C are given symmetric matrices, and $C \succ 0$. Find a convenient description of \mathcal{S}° . Can you optimize a linear function over \mathcal{S}° ?

4. Consider the polynomial $p(x) = x^4 + 2ax^2 + b$. For what values of (a, b) is this polynomial non-negative? Draw the region of nonnegativity in the (a, b) plane. Where does the discriminant of p vanish? How do you explain this?
5. Consider the polynomial system $\{x + y^3 = 2, x^2 + y^2 = 1\}$.
 - (a) Is it feasible over \mathbb{C} ? How many solutions are there?
 - (b) Is it feasible over \mathbb{R} ? If not, give a Positivstellensatz-based infeasibility certificate of this fact.

6. In general, the SOS decomposition of a univariate polynomial is not unique. Given a specific basis of $\mathbb{R}[x]$ (for instance, the standard monomial basis we have been considering), a “natural” choice can be obtained by finding a matrix Q satisfying

$$p(x) = [x]_d^T Q [x]_d, \quad Q \succeq 0$$

and such that the determinant of Q is as large as possible. This “central solution” can be computed by solving a convex optimization problem, since $\log \det Q$ is a concave function on the region where Q is positive semidefinite. In fact, this convex optimization problem can be reformulated a semidefinite programming problem.

- (a) Compute numerically the central solution for the polynomial $p(x) = x^6 - 6x^5 + 16x^4 - 24x^3 + 22x^2 - 12x + 4$.
- (b) Show, using the KKT optimality conditions, that in general the inverse of the optimal matrix Q is a Hankel matrix.
7. Consider a random variable X , with an unknown probability distribution supported on the set Ω , and for which we know its first $d + 1$ moments (μ_0, \dots, μ_d) . We want to find bounds on the probability of an event $S \subseteq \Omega$, i.e., want to bound $P(X \in S)$. We assume S and Ω are given intervals. Consider the following optimization problem in the decision variables c_k :

$$\min \sum_{k=0}^d c_k \mu_k \quad \text{s.t.} \quad \begin{cases} \sum_{k=0}^d c_k x^k \geq 1 & \forall x \in S \\ \sum_{k=0}^d c_k x^k \geq 0 & \forall x \in \Omega. \end{cases} \quad (1)$$

- (a) Show that any feasible solution of (1) gives a valid upper bound on $P(X \in S)$. How would you solve this problem?
- (b) Assume that $\Omega = [0, 5]$, $S = [4, 5]$, and we know that the mean and variance of X are equal to 1 and $1/2$, respectively. Give upper and lower bounds on $P(X \in S)$. Are these bounds tight? Can you find the worst-case distributions?
8. Consider linear maps between symmetric matrices, i.e., of the form $\Lambda : \mathcal{S}^n \rightarrow \mathcal{S}^m$. A map is said to be a *positive map* if it maps the PSD cone \mathcal{S}_+^n into the PSD cone \mathcal{S}_+^m (i.e., it preserves positive semidefinite matrices).

- (a) Show that any linear map of the form $A \mapsto \sum_i P_i^T A P_i$, where $P_i \in \mathbb{R}^{n \times m}$, is positive. These maps are known as *decomposable* maps.
- (b) Show that the linear map $C : \mathcal{S}^3 \rightarrow \mathcal{S}^3$ (due to M.-D. Choi) given by:

$$C : A \mapsto \begin{bmatrix} 2a_{11} + a_{22} & 0 & 0 \\ 0 & 2a_{22} + a_{33} & 0 \\ 0 & 0 & 2a_{33} + a_{11} \end{bmatrix} - A.$$

is a positive map, but is not decomposable.

Hint: Consider the polynomial defined by $p(x, y) := y^T \Lambda(xx^T)y$. How can you express positivity and decomposability of the linear map Λ in terms of the polynomial p ?